

ORIGINAL ARTICLE

Open Access



Enhanced laccase-mediated transformation of diclofenac and flufenamic acid in the presence of bisphenol A and testing of an enzymatic membrane reactor

Veronika Hahn^{1,2*}, Mareike Meister^{1,2}, Stephan Hussy³, Arno Cordes⁴, Günther Enderle³, Akuma Saningong⁵ and Frieder Schauer¹

Abstract

The inadequate removal of pharmaceuticals and other micropollutants in municipal wastewater treatment plants, as evidenced by their detection of these substances in the aquatic environment has led to the need for sustainable remediation strategies. Laccases possess a number of advantages including a broad substrate spectrum. To identify promoting or inhibitory effects of reaction partners in the remediation processes we tested not only single compounds—as has been described in most studies—but also mixtures of pollutants. The reaction of diclofenac (DCF) and flufenamic acid (FA), mediated by *Trametes versicolor* laccase resulted in the formation of products, which were more hydrophilic than the respective reactant (reactant concentration of 0.1 mM; laccase activity 0.5 U/ml). Analyses (HPLC, LC/MS) showed that the product **1a** and **1b** for DCF and FA, respectively, to be a *para*-benzoquinone imine derivative. The formation of **1a** was enhanced by the addition of bisphenol A (BPA). After 6 days 97% more product was formed in the mixture of DCF and BPA compared with DCF tested alone. Product **1a** was also detected in experiments with micropollutant-supplemented secondary effluent. Within 24 h 67% and 100% of DCF and BPA were transformed, respectively (25 U/ml). Experiments with a membrane reactor (volume 10 l; phosphate buffer, pH 7) were in good agreement with the results of the laboratory scale experiments (50 ml). EC₅₀-values were also determined. The data support the use of laccases for the removal or detoxification of recalcitrant pollutants. Thus, the enzyme laccase may be a component of an additional environmentally friendly process for the treatment stage of wastewater remediation.

Keywords: Quinone, Bioremediation, Detoxification, Pharmaceuticals, Biodegradation, *Vibrio fischeri*

Introduction

The daily use of anti-inflammatory drugs such as diclofenac (DCF) or flufenamic acid (FA) in creams or tablets, as well as of plastic materials which release bisphenol A (BPA), contributes to pollution with these environmentally dangerous compounds. BPA is used in plastic industries; consequently the release from different sources is conceivable such as packages for food

and drink, varnishes, paintings and glues (Barnabe et al. 2009). The main entries for pharmaceuticals such as DCF or FA in ground and surface water are sludge or effluents of wastewater treatment plants (WWTP) where the pollutants are insufficiently degraded. Another source is dung and manure of farm animals which were treated with these substances (Ternes 1998; Deblonde et al. 2011; Samaras et al. 2013; Carmona et al. 2014).

Only 17–69% of DCF is removed in sewage treatment plants (Ternes 1998; Zwiener et al. 2000; Heberer 2002a; Heberer and Feldmann 2005; Deblonde et al. 2011; Samaras et al. 2013). FA is also insufficiently removed in

*Correspondence: veronikahahn@gmx.at

¹ Institute of Microbiology, Ernst-Moritz-Arndt-University Greifswald, Friedrich-Ludwig-Jahn-Str. 15, 17487 Greifswald, Germany
Full list of author information is available at the end of the article

WWTP. Moreover, Gracia-Lor et al. (2012) and Carmona et al. (2014) described for DCF and FA higher concentrations in the effluent than in the corresponding influent of Spanish WWTP. The authors proposed a release as result of a cleavage of transformation products, deconjugation of metabolites and desorption from organic material.

DCF, FA and BPA were found repeatedly in surface water. Ternes (1998) reported for DCF a median of 0.15 µg/l and a maximum of 1.20 µg/l for German river and stream water. In Berlin, up to 0.5–1 µg/l and 0.38 µg/l were found in the Teltowkanal and in ground water, respectively (Heberer 2002b). FA was detected in river and tap water with average concentrations of 21 and 16 ng/l, respectively (Carmona et al. 2014). BPA was also found in surface and ground water (Loos et al. 2009, 2010). In small streams of the German region called Hessesches Ried BPA concentrations of up to 1.92 µg/l were detected (Quednow and Püttmann 2008).

DCF, FA as well as BPA may cause toxic effects. In rainbow trout DCF accumulates in the bile to a factor of approximately 580, and additionally the tested environmentally relevant concentrations caused necrosis in the kidney (Mehinto et al. 2010). The endocrine disrupting compound BPA leads to the induction of feminization in aquatic organisms, such as frogs (Levy et al. 2004; Bhandari et al. 2015). Because of the potential risk for human health, DCF was included in a watch list of emerging pollutants by the EU (Commission 2013, 2015). Nadanaciva et al. (2013) working with zebrafish, showed a high toxicity for FA, but it was not possible to locate the morphological changes because the difference between the FA concentration with no effect where no death occur (NOEC) and the concentration causing 100% lethality (LC_{100}) was too small. Thus, for FA as well as DCF the same NOEC of 10 µM was determined whereas the LC_{100} was 30 µM for FA and 300 µM for DCF, allowing the determination of morphological changes, such as liver browning, only for DCF (Nadanaciva et al. 2013). Furthermore, in vitro test showed a considerable effect of FA on liver mitochondria (Nadanaciva et al. 2013).

The repeated detection of pharmaceutical residues in the aquatic environment demonstrates the urgent need for efficient wastewater treatment processes. In this regard, the enzyme laccase has been considered as a tool for a sustainable remediation strategy and for additional purification steps in wastewater treatment. Laccases [E.C. 1.10.3.2] can oxidize a broad range of compounds such as phenols or amines (Keilin and Mann 1939; Bollag et al. 1988; Thurston 1994; Mikolasch and Schauer 2009) by one-electron-reactions (Nakamura 1960; Solomon et al. 2001, 2008; Munk et al. 2017). In these laccase-mediated oxidations, radicals are formed which can undergo two possible reactions. The first involves binding

to other compounds, while the second involves cleavage (Hahn et al. 2014). An important advantage of laccases, in particular for remediation purposes, is their need for atmospheric oxygen as the only co-substrate. Thus, oxygen is reduced to water.

Despite the well known removal of water contaminants such as BPA or other phenols, DCF and estrogens by free or immobilized laccase of *Trametes versicolor* (Auriol et al. 2008; Catapane et al. 2013; Ammann et al. 2014) has until now in most studies been tested only single compounds. The transformation in mixtures of micropollutants was described in most cases without the distinction of the influencing effect of the single compounds on the degradation in these mixtures (Tran et al. 2010; Nguyen et al. 2014; Asif et al. 2017b). To determine the effect of multiple contaminants on transformation rate and product formation we employed mixtures of pollutants. In addition, removal of FA was tested due to limited data.

The laccase of *T. versicolor* was employed for the transformation of DCF, FA and BPA and combinations of the substances were also tested. The removal of these test substances as well as product formation was determined by HPLC analyses. The structural characterization of products was performed by LC/MS.

The results of this study will be used for the development of an enzyme-based membrane reactor designated as an additional or tertiary treatment stage in sewage treatment plants.

Materials and methods

Chemicals

All chemicals used were of analytical grade and were used as received.

Diclofenac sodium salt, flufenamic acid, bisphenol A, 4'-hydroxydiclofenac and ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt) were purchased from Sigma-Aldrich Chemie GmbH (Steinheim, Germany). 5-Hydroxydiclofenac was obtained from Toronto Research Chemicals (Toronto, Canada). Potassium dihydrogen phosphate and di-sodium hydrogen phosphate dihydrate were purchased from Carl-Roth GmbH and Co. KG (Karlsruhe, Germany).

Enzyme

The laccase from *T. versicolor* was obtained from ASA Spezialenzyme GmbH (Wolfenbüttel, Germany). The company markets the enzyme under the name "laccase C". It was produced by a fed-batch fermentation process. The fungal biomass was separated by centrifugation and the enzyme-containing supernatant was concentrated by ultrafiltration. After addition of stabilizing agents the retentate of the ultrafiltration step was lyophilized. The laccase is active within pH 3.0–7.5 (pH

optimum of 5). It was used as received (activity > 800 U/g; substrate:syringaldazine).

For transformation experiments the laccase was used in Sørensen's phosphate buffer at pH 7 in accordance with the pH 7–7.5 of the secondary wastewater.

Experimental procedures for transformation in phosphate buffer or secondary effluent

For the transformation assays the respective compounds (initial reactant concentration: 0.1 mM) were incubated in a final volume of 50 ml in 500-ml flasks. The reaction solution was phosphate buffer or secondary effluent. The laccase activity was 0.5 U/ml for reactions in phosphate buffer and, additionally, 2.5 and 25 U/ml in wastewater. The secondary effluent from a wastewater treatment plant (85,000 inhabitant equivalents Greifswald, Germany) was 0.22 μm filtered and spiked with the respective compound.

In controls the respective compounds were incubated in phosphate buffer or secondary effluent without laccase. Reaction mixtures were incubated with agitation at 150 rpm at 25 °C in the dark.

Reactor setup and operation

The laboratory reactor was equipped with an Atec ultrafiltration system to avoid loss of laccase (Fig. 1). From the different membranes tested for that purpose a tubular

ceramic membrane element with a pore size of 10 nm gave the best results (Membrane: gamma- Al_2O_3 , support: alpha Al_2O_3 , manufacturer:Inopor). Like Atec's commercial Miditube filtration systems the laboratory unit uses a unique design allowing almost independent control of cross flow over the membrane surface, and trans membrane pressure. Filtration with low trans membrane pressures leads only to the formation of thin layers on the ultrafiltration membranes extending the time interval to back flush.

Depending on the temperature (20–27 °C) the permeate flow of 26–36 l/(m² h bar) was constant during experiment if working pressure was kept low (1.1 bar). The ratio of bleeding to permeate flow was adjusted to 15:1. System was run continuously during duration of the experiment (4 h).

The respective substances (initial reactant concentration: 0.1 mM) were tested in a final volume of 10 l in phosphate buffer. The laccase activity employed was 0.5 U/ml.

The decrease of reactants and the products formed were analyzed by HPLC.

Measurement of laccase activity

The activity of laccase was determined spectrophotometrically at 420 nm with ABTS as substrate (Bourbonnais and Paice 1990) using the method described by Jonas




Fig. 1 Scheme of the reactor equipped with an Atec ultrafiltration system

et al. (1997) at pH 5. 1 U is defined as the turnover of 1 $\mu\text{mol}/\text{ml min}$

Analytical HPLC (high-pressure liquid chromatography)

For routine analysis, the reaction mixtures were analyzed using an HPLC system LC-10AT VP (Shimadzu, Germany) consisting of a FCV-10AL VP pump, SPD-M10A VP diode array detector, and a SCL-10A VP control unit controlled by Class-VP version 6.12 SP5. Substances were separated on an endcapped, 5- μm , LiChroCART[®] 125-4 RP18 column (Merck, Darmstadt, Germany) run at a flow rate of 1 ml/min. The solvent system used consisted of a gradient of methanol (eluent A) and 0.1% phosphoric acid (eluent B), starting from an initial ratio of 10% A and 90% B and reaching 100% methanol within 14 min. Elution with methanol was continued for a further 6 min.

Structural characterization of products by LC/MS (liquid chromatography/mass spectrometry)

The reaction mixtures and isolated products were characterized using a LC/MS system. The atmospheric pressure ionization (API) mass spectrometry experiments were performed on an Agilent Series 1200 HPLC system with diode array detector and an Agilent 6120 quadrupole mass spectrometer (Waldbronn, Germany). The MS was run with the electrospray ionization (API-ES) source in positive mode (dry and nebulizer gas: nitrogen; nebulizer pressure: 45 psig; drying gas flow: 10 l/min; drying gas temperature: 350 °C; capillary voltage: 4 kV; fragmentor voltage: 75 V). HPLC separation was performed on a Zorbrax SB-C18 (2.1 \times 50 mm, 1.8 μm) column (Agilent, Waldbronn, Germany), at a flow rate of 0.07 ml/min. The solvent system consisted of a gradient of acetonitrile (eluent A) and 0.1% aqueous ammonium formate (eluent

B), starting from an initial ratio of 10% A and 90% B and reaching 100% methanol within 7 min. Elution with methanol was continued for a further 6 min.

Experimental methods; UV-vis data and MS spectra for products **1a_{I-III}**, **2**, and **1b_{I,II}** (Additional file 1: Table S1–S6) are available in the electronic additional file.

Determination of toxicity using *Vibrio fischeri* (DIN EN ISO 11348, (DIN 38412-L34, DIN 38412-L341))

This marine bacterium is bioluminescent and emits light with a wavelength of 490 nm. The bioluminescence is reduced in the presence of toxic agents.

Samples were taken before and after laccase treatment from a membrane reactor (initial concentration of reactants: 0.1 mM, volume: 10 l; phosphate buffer; incubation time: 4 h).

For the determination of EC₂₀ and EC₅₀ the samples were inoculated with *Vibrio fischeri* (LCK 482, Hach Lange GmbH, Düsseldorf, Germany) and incubated for 30 min. The reduction of bioluminescence was measured photometrically (LUMISTox 300-Version 3.03 Hach Lange GmbH, Düsseldorf, Germany). The results were compared with a control containing physiological saline solution instead of the sample.

Results

Transformation of DCF, FA and BPA including structural characterization of the products formed

DCF, FA and BPA were transformed by laccase of *T. versicolor* to varying extents (concentration of reactants: 0.1 mM; Fig. 2). The reactant BPA was completely transformed within 24 h in the laccase-mediated reactions irrespective of whether it was tested individually or in combination with DCF or FA. The influence of added




Fig. 2 Laccase-mediated transformation [%] of DCF or FA treated individually or in combination with BPA in phosphate buffer (concentration of reactants 0.1 mM)

BPA on the transformation of the other two compounds was significant within 3 h. In the reaction with BPA the acids DCF and FA were 2.7 and 7 times faster transformed in the presence of BPA. After 24 h DCF transformation reached approximately 65% in both reactions whereas FA reached 9% (without BPA) and 15% (with BPA).

For all reactions different low- and high-molecular weight products were detected but due to the small quantities and difficult analyses (at least for the high-molecular weight products) only a selection of products were structurally characterized.

The laccase-mediated reaction of DCF leads to the formation of product **1a_I** which was detected after an incubation period of 6 days. In combination of DCF and BPA the product (**1a_{II}**) was detected already after 3 h. Thus, within 6 days only 3% of **1a_I** was formed in the reaction of DCF (**1a_I**: 0.39 µg/ml; 0.0013 mM) compared to the mixture of DCF and BPA (**1a_{II}**: 13.34 µg/ml; 0.0430 mM).










The product **1a_I** was more hydrophilic (*R_f* 12.23 min) than DCF (*R_f* 14.18 min) and showed UV-vis absorption maxima at 202, 267 and 462 nm (Table 1). The LC/MS (API-ES positive mode) analyses of **1a_I** resulted in the detection of [M + H]⁺ ion multiplets at *m/z* (rel.

intensity) 310.0, 312.0 and 313.8 (90:74:5; Additional file 1: Table S1) implying an hydroxylation and oxidation of DCF.

The UV-vis absorption maximum at 267 nm as well as the ion multiplets at *m/z* 310.0, 312.0 and 314.0 (Additional file 1: Table S3) were reported by Shen et al. (1999) for the *p*-benzoquinone imine derivative (**1a_{III}**) of 5-hydroxydiclofenac (5HDCF) formed during incubation of DCF with liver microsomes. Thus, the reactions of DCF (alone or in a mixture with BPA) or 5HDCF resulted in the same product **1a**. For the laccase-mediated DCF transformation a hydroxylation at C5 with subsequent formation of a *p*-benzoquinone imine **1a_{I,II}** was presumed.

The 4'-hydroxydiclofenac (4'HDCF) was incubated with laccase to exclude the 4'-position for the hydroxylation. The reaction resulted in the formation of the corresponding quinone imine (**2**). The LC/MS (API-ES positive mode) analyses of **2** resulted also in the detection of [M + H]⁺ ion multiplets at *m/z* (rel. intensity) 310.0, 312.0, 314.0 (100, 62, 18; Table 1; Additional file 1: Table S4) but the retention time of 12.41 min as well as the UV-vis absorption maxima at 202, 278, 524 nm were different to those of **1a_{I,III}**.

Table 1 Structural data of products 1a_{I-III} formed during laccase-mediated reactions of DCF alone or in combination with BPA as well as 5HDCF; data for product 2 from the reaction of 4'HDCF and for products 1b_{I,II} from the reaction of FA alone or in combination with BPA are provided

Substrate	DCF	DCF and BPA	5HDCF	4'HDCF	FA	FA and BPA
						
Product ^a		 [M] 310.14		 [M] 310.14	 [M] 295.22	
	1a_I	1a_{II}	1a_{III}	2	1b_I	1b_{II}
<i>R_f</i> (HPLC) (min)	12.23	12.22	12.29	12.41	12.09	12.00
UV-vis λ _{max}	202, 267, 462	203, 266, 463	204, 266, 463	202, 278, 524	203, 274, 440 nm	203, 277, 434 nm
MS (AP-ESI pos. ion mode): <i>m/z</i> (%)	[M + H] ⁺ 310.0, 312.0, 313.8 (90, 74, 5), [M + Na] ⁺ 332.0 (9), [M + K] ⁺ 347.8 (32)	[M + H] ⁺ 310.0, 311.9, 313.9 (100, 55, 10), [M + Na] ⁺ 331.9 (9), [M + K] ⁺ 347.9 (5)	[M + H] ⁺ 310.0, 312.0, 314.0 (100, 58, 7), [M + Na] ⁺ 331.9 (6)	[M + H] ⁺ 310.0, 312.0, 314.0 (100, 62, 18), [M + Na] ⁺ 331.9 (7)	[M + H] ⁺ 296.1 (100), [M + Na] ⁺ 318.0 (12), [M + K] ⁺ 334.0 (29)	[M + H] ⁺ 296.0 (100)
MS (AP-ESI neg. ion mode): <i>m/z</i> (%)	Not detected	Not detected	Not detected	Not detected	[M-H] ⁻ 294.1 (81)	[M-H] ⁻ 294.2 (100)

According to the HPLC and LC/MS data products **1a_{I-III}** and **1b_{I,II}** have the same chemical structure, respectively

^a Only Products of DCF and FA are shown

Similar to the reaction of DCF the reaction of FA alone and in combination with BPA resulted after 10 days in a more hydrophilic product (**1b_I** R_f 12.09 min or **1b_{II}** R_f 12.00 min) than FA (R_f 14.99 min). The LC/MS analyses (API-ES positive and negative mode) for the laccase-mediated reaction of FA resulted in the detection of $[M + H]^+$ (rel. intensity) 296.1 (100), $[M + Na]^+$ 318.0, $[M + K]^+$ 334.0 and $[M - H]^-$ 294.1 (Additional file 1: Table S5). The C4-position for the hydroxylation was assumed because of the similarities with the reaction of DCF, though no authentic standard was available. The amount of products (**1b_{I,II}**) formed was too small to determine a dependency between product yield and BPA addition. The concentration of BPA may not have been high enough to promote the formation of **1b_{II}**.

DCF transformation alone and in combination with BPA in a membrane reactor and determination of toxicity

Experiments with a membrane reactor (volume of 10 l) were carried out to provide information about the efficiency of laccase-mediated transformation of pollutants.

In samples of laboratory scale, and in the filtrate of a membrane reactor, DCF transformation was enhanced by the presence of BPA (Fig. 3). Thus, in both situations the removal of DCF was approximately 20% higher after 4 h in the mixture of DCF and BPA than without BPA. Furthermore, **1a_I** was not detected when DCF was tested alone due to the short incubation time of 4 h. In the combination of DCF and BPA the product **1a_{II}** was formed as in the 50 ml assay. The loss of laccase during the 4 h incubation in the reactor was negligible.




Fig. 3 Transformation of DCF tested individually (black lines) and in mixture with BPA (dashed lines) in 500-ml flasks (laboratory scale; reaction volume: 50 ml; filled triangles) and in filtrate of a membrane reactor (reaction volume: 10 l; open triangle) in phosphate buffer (pH 7) and a laccase activity of 0.5 U/ml (concentration of reactants: 0.1 mM)

The reaction assays of the membrane reactor were tested for their toxicity against *Vibrio fischeri* before and after incubation with laccase (Table 2). In all tested assays the toxicity decreased with decreasing concentration of the pollutant/s caused by laccase-mediated transformation. After an incubation time of 4 h 32% of DCF was removed whereas in the mixture with BPA 52% of DCF was transformed. The reactions with BPA resulted in a complete transformation.

For DCF and BPA the toxicity was reduced by 42 and 76%, respectively (according to EC_{50} -values). In the reaction of DCF with BPA the toxicity was diminished by 69% after laccase treatment. The residual toxicity is probably caused by products formed or—in case of DCF—by the high concentration of reactant, which is left in the reaction assay. Nevertheless, the data show a clear decrease in toxicity due to laccase treatment. A longer incubation time would be expected to lead to higher transformation of DCF, which may cause a further decrease in toxicity.

DCF transformation alone and in combination with BPA in municipal wastewater

A laccase activity of 0.5 or 2.5 U/ml was used for the experiments with the secondary effluent. However, this activity, as seen in the previous experiments, resulted in no transformation of DCF. 25 U/ml in wastewater was needed to achieve 67% transformation of DCF and complete removal of BPA within 24 h. In the mixture of DCF and BPA 33% and 65% were transformed, respectively.

Within 24 h **1a_I** was only detected in the reaction with DCF alone (**1a_I** 7.03 μ g/ml; 0.0227 mM) in accordance with the faster transformation of DCF in this reaction compared to the reaction in a mixture with BPA.

Discussion

Laccase-mediated reactions resulted in the transformation of micropollutants which are problematic for aquatic environment. The transformation efficiency of BPA, DCF, and FA mediated by *T. versicolor* laccase in the lab scale reactions (reaction volume 50 ml; phosphate buffer) was: BPA > DCF > FA. The laccase-mediated oxidation can be hindered by different factors such as steric problems through the size of the substrate-binding site (Tadesse et al. 2008; Galli et al. 2011).

Moreover, it is believed that laccase can only oxidize those substrates which have a redox potential below that of the enzyme (Xu et al. 2000). Thus, laccases were divided according to their redox potential into low (0.4–0.5 V, determined against a normal hydrogen electrode) and high redox potential (0.7–0.8 V) laccases (Xu 1996; Xu et al. 1998; Li et al. 1999; Frascioni et al. 2010). The laccase of *T. versicolor* belongs to the high redox potential laccases (0.785 V, Reinhammar 1972). The low redox

Table 2 Results of the toxicity test (test organism: *Vibrio fischeri*), including concentration of DCF and BPA, before and after laccase treatment in a membrane reactor (volume: 10 l; phosphate buffer; incubation time: 4 h)

Reaction	Concentration				Toxicity test (<i>Vibrio fischeri</i>)			
	DCF		BPA		EC ₅₀		EC ₂₀	
	mg/l	mM	mg/l	mM	mg/l	mM	mg/l	mM
DCF								
Before laccase treatment	31.81	0.10			11.36	0.0357	5.53	0.0174
After laccase treatment	21.70	0.07			19.50	0.0613	9.59	0.0302
BPA								
Before laccase treatment			22.83	0.10	5.24	0.0230	1.36	0.0060
After laccase treatment			0	0	22.16	0.0971	3.01	0.0132
DCF and BPA								
Before laccase treatment	31.81	0.10	22.83	0.10	3.91	0.0143	1.50	0.0055
After laccase treatment	15.27	0.05	0	0	12.41	0.0454	6.24	0.0225

potential of mono- and dihydroxylated aromatic compounds such as guaiacol or hydroquinone (0.5–0.6 V) allows an easy oxidation by laccase (Mai et al. 2001). The kind of substituents has also an influence on the oxidizability (Tadesse et al. 2008; Hahn et al. 2014). Thus, electron-withdrawing groups such as fluoro or chloro and carboxyl groups which are part of DCF and FA reduce the possibility for oxidation whereas electron-donating hydroxyl groups of BPA decrease the redox potential resulting in an easier oxidation.

The addition of BPA at least slightly enhanced the transformation of DCF and FA although complete transformation was not achieved. An incomplete biotransformation of DCF by laccase is not unexpected, and was also described for another laccase preparation of *T. versicolor* (Sigma-Aldrich Chemie GmbH, ref. 38429, Buchs, Switzerland) by Margot et al. (2013).

The reaction of DCF and BPA in 10 l phosphate buffer in the membrane reactor resulted also in enhanced transformation of DCF in the presence of BPA. This demonstrates that the scale up was successful. In both—the 50 ml and 10 l scale—experiments the removal of DCF was approximately 20% higher after 4 h in the mixture of DCF and BPA than without BPA.

Nair et al. (2013) described an efficient transformation of DCF by immobilized laccase of *Coriolopsis gallica* in a continuous stirred tank membrane reactor, though the reactor volume was only 50 ml (flow rate: 40 ml/h). Thus, 70% of the DCF was transformed within 80 h (initial concentration: 10 µM, laccase activity approximately 1 U/ml, McIlvaine buffer pH5). Nguyen et al. (2014) described the removal of BPA and DCF in an enzymatic membrane reactor using the laccase of *Myceliophthora thermophila*. Within 132 h 85% BPA and 60% DCF was continuously transformed (initial concentration 500 µg/l d, laccase

activity approximately 90 µM/min, Milli-Q-water). A further experiment with the same laccase and a mixture of BPA, DCF with 28 other pharmaceuticals such as naproxen or ibuprofen was described by Asif et al. (2017b). In the enzymatic reactor approximately 88% BPA and 45% DCF were transformed (initial concentration: 20 µg/l, laccase activity approximately 95–100 µM/min, Milli-Q-water).

The enhanced transformation of DCF and FA in the presence of BPA is striking. It suggests that BPA, or transformation products resulting from a cleavage of BPA, may act as mediators. Different authors described hydroxylated monoaromatic compounds such as 4-isopropenylphenol or 4-ethyl-2-methoxyphenol which were formed during laccase-mediated transformation of BPA (Fukuda et al. 2001; Chairin et al. 2013; Arca-Ramos et al. 2015). In a similar way, mediators can be formed in course of the degradation processes such as the depolymerization of lignin (or are produced by the fungus itself). Such mediators are small-molecular weight substances such as syringaldehyde or vanillin. These compounds may be oxidized by laccase and, in turn, can oxidize compounds which are not accessible for the laccase due to their high redox potential or to steric hindrance (Bourbonnais and Paice 1990; Baiocco et al. 2003; Wesenberg et al. 2003; Kunamneni et al. 2008; Tadesse et al. 2008; Mogharabi and Faramarzi 2014). Lloret et al. (2010) showed an improved transformation of DCF in the presence of different mediators such as syringaldehyde or 1-hydroxybenzotriazole.

Formation of product 1a

The formation of **1a** was promoted by the presence of BPA. Thus, only 3% of **1a**₁ was detected after 6d in the reaction of DCF alone, compared to the reaction in the presence of BPA. In contrast, within the first 4 h of

incubation, the transformation of DCF was approximately 20% higher in the reaction with BPA and nearly the same after 24 h. Thus, despite an only slightly enhanced transformation of DCF in a mixture with BPA the formation of **1a**_{I,II} is favored in the mixture suggesting an easier hydroxylation in the reaction with added BPA than in the reaction only with DCF.

In contrast with the reactions in phosphate buffer, **1a**_{I,II} was formed in the secondary effluent, in particular in the reaction with DCF alone, due to a slower removal of DCF in the mixture with BPA.

The product **1a**_{I,II} and **1b**_{I,II} is probably formed via a hydroxylation of DCF or FA by a nucleophilic attack of water on a laccase-generated cation radical, with subsequent oxidation resulting in a *para*-benzoquinone imine derivative. To the best of our knowledge, this is the first description of products **1a** and **1b** for laccase-catalyzed reactions. The laccase-mediated hydroxylation of *para*-dihydroxylated aromatic compounds was reported previously (Manda et al. 2007; Hahn et al. 2009) but interestingly the hydroxylation of DCF did not involve the well characterized substitution of a chloro group as previously described (Minard et al. 1981; Iimura et al. 1996; Kordon et al. 2010; Hahn et al. 2014). Hydroxylation on the chlorinated ring of DCF is probably hindered due to the electron withdrawing effects of the chloride atoms (Faber et al. 2012). The hydroxylation of DCF took place at the C5-position. Marco-Urrea et al. (2010) also described the formation of 5-hydroxydiclofenac in reactions with whole cells of *T. versicolor* cultures, whereas with isolated laccase only 4-(2,6-dichlorophenylamino)-1,3-benzenedimethanol was detected. In these laccase-mediated reactions again the C5 and additionally C3 was derivatized (Marco-Urrea et al. 2010).

The formation of the quinone imine derivative **1a** was also described for a reaction of DCF catalyzed by a peroxymonosulfate/Cobalt(II) system by a one-electron mechanism similar to laccase-mediated reactions (Ahmed et al. 2012). The C5-position for hydroxylation is favored both for the peroxymonosulfate/Cobalt(II) system, as well as for the electrochemical formation of the *para*-benzoquinone derivative (Ahmed et al. 2012; Faber et al. 2012).

DCF is metabolized by human cytochrome-P450 enzymes to a range of hydroxylated products, namely 4'-, 3'- and also 5-hydroxydiclofenac (so called phase I metabolites; Davies and Anderson 1997; Dorado et al. 2003). For humans, as well as fungi, the formation of 4'-hydroxydiclofenac is the primary transformation pathway of DCF, 3'- and 5-hydroxydiclofenac are only minor products (Davies and Anderson 1997; Webster et al. 1998; Dorado et al. 2003).

Detoxification in a membrane reactor

The experimental data provided for the reaction course of DCF and BPA in a reaction volume of 50 ml and 10 l demonstrate that the laboratory scale results can be transferred to higher reaction volumes, which are necessary for biotechnological applications in membrane reactors.

The determined toxicity of DCF and BPA alone or in a mixture decreased after the treatment with laccase in the membrane reactor due to decreased concentrations of the reactants. Nevertheless it has to be borne in mind, that the hydroxylated product 5HDCF (EC₅₀ 39.59 mg/ml, 0.1268 mM) possesses a higher, whereas the oxidized form of 5HDCF-the quinone imine (**1a**_{III}: EC₅₀ 14.37 mg/ml, 0.0463 mM), has a similar toxicity as DCF (EC₅₀ 11.36 mg/ml, 0.0357 mM).

Marco-Urrea et al. (2010) also observed a decrease in ecotoxicity (Microtox test with *V. fischeri*) with a concomitant removal of DCF and its transformation products in a culture with *T. versicolor*.

The laccase-mediated transformation of BPA was demonstrated previously by Fukuda et al. (2001). The reaction results in the low-molecular weight product 4-isopropenylphenol and in dimers and oligomers of BPA with a higher molecular weight than BPA (Fukuda et al. 2001; Uchida et al. 2001; Fukuda et al. 2004). Neither product group showed estrogenic activity (Fukuda et al. 2004).

Transformation of DCF and BPA in secondary effluent

In the tested municipal wastewater, the transformation of DCF and BPA was diminished in comparison with that in phosphate buffer. Although the values for the laccase-mediated transformation with 65% for DCF and 100% for BPA resemble that determined in phosphate buffer, these values were achieved only with 50-fold higher laccase activity than in phosphate buffer. In a mixture of DCF and BPA, the transformation was 2 and 1.5-fold lower, as compared to the individual compounds alone, whereas in phosphate buffer the transformation was higher (in case of DCF) or the same (in case of BPA).

A similar effect was observed by Nair et al. (2013). Thus, the incubation of DCF (in a mixture with BPA and 17- α -ethinylestradiol) in secondary effluent leads to 40% less transformation of DCF. The authors attributed the lower transformation to the presence of organic matter (Nair et al. 2013) which may act as radical scavengers (Ahmed et al. 2012). Nevertheless, the influence of inorganic compounds such as CO₃²⁻, HCO₃⁻ or Cl⁻ cannot be ruled out (Ahmed et al. 2012). Thus, chloride and fluoride anions have been described as laccase inhibitors (Xu 1996; Nagai et al. 2002; Ramírez-Cavazos et al. 2014). The effect of such wastewater constituents can be manifold.

Beside radical scavenging, the components may influence the structure of the laccase, resulting in insufficient binding of the substrate in the active site, and consequently in less transformation. A reaction of the pollutants with the components is also conceivable preventing binding on the active site of the enzyme. Beyond this the secondary effluent may contain other inhibitors. Thus, azide and cyanide anions as well as L-cysteine have been described as laccase inhibitors (Nagai et al. 2002; Ramírez-Cavazos et al. 2014). Kim and Nicell (2006) described the reduced conversion of BPA in the presence of different anions such as fluoride, sulfide, sulfite and cyanide. The influence of humic acid on the removal of pollutants is also not clear at the moment (Asif et al. 2017a). Finally, it cannot be ruled out that other laccase substrates are present in the secondary effluent, which lower the transformation capacity for DCF and BPA.

The removal of persistent pollutants during wastewater remediation remains a challenge for the future. Efforts to determine the effects of pollutants in a mixture during the water purification process should be enhanced, in particular for enzyme-mediated processes. The decrease of toxicity in the assays containing DCF, BPA or a mixture of both confirms the suitability of laccase-mediated reactions for micropollutant elimination. The enzyme laccase may be an important tool for a greener, and thereby environmentally friendly, remediation process.

Additional file

Additional file 1. Additional tables.

Abbreviations

ABTS: 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt; AP-ESI: atmospheric-pressure electrospray ionization; DCF: diclofenac; BPA: bisphenol A; FA: flufenamic acid; 4'HDCF: 4'-hydroxydiclofenac; 5HDCF: 5-hydroxydiclofenac; HPLC: high-pressure liquid chromatography; LC/MS: liquid chromatography/mass spectrometry; WWTP: wastewater treatment plants.

Authors' contributions

SH, AC, GE, AS, FS and VH initiated the project. SH, MM and VH performed the experiments and analyzed the data. All authors read and approved the final manuscript.

Author details

¹ Institute of Microbiology, Ernst-Moritz-Arndt-University Greifswald, Friedrich-Ludwig-Jahn-Str. 15, 17487 Greifswald, Germany. ² Leibniz Institute for Plasma Science and Technology (INP Greifswald e.V.), Felix-Hausdorff-Str. 2, 17489 Greifswald, Germany. ³ Atec Automatisierungstechnik GmbH, Emmi-Noether-Str. 6, 89231 Neu-Ulm, Germany. ⁴ ASA Spezialenzyme GmbH, Am Exer 19 C, 38302 Wolfenbüttel, Germany. ⁵ EurA Consult AG, Max-Eyth-Str. 2, 73479 Ellwangen, Germany.

Acknowledgements

We thank K. Wedig (municipal wastewater treatment plant of Greifswald, Germany) for support during determination of toxicity using *Vibrio fischeri*. R. Jack is gratefully acknowledged for help in preparing the manuscript.

This work was funded by the Federal Ministry for Economic Affairs and Energy with the central innovation programme for small and medium-sized enterprises (16KN017326) which is gratefully acknowledged.

Competing interests

All authors declare that they have no competing interests.

Availability of data and materials

Not applicable.

Consent for publication

Not applicable.

Ethics approval and consent to participate

This article does not contain any studies with human participants or animals performed by any of the authors.

Funding

This study was funded by the Federal Ministry for Economic Affairs and Energy with the central innovation programme for small and medium-sized enterprises (16KN017326).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 2 January 2018 Accepted: 27 January 2018

Published online: 24 February 2018

References

- Ahmed MM, Barbati S, Doumenq P, Chiron S (2012) Sulfate radical anion oxidation of diclofenac and sulfamethoxazole for water decontamination. *Chem Eng J* 197:440–447
- Ammann EM, Gasser CA, Hommes G, Corvini PF (2014) Immobilization of defined laccase combinations for enhanced oxidation of phenolic contaminants. *Appl Microbiol Biotechnol* 98(3):1397–1406. <https://doi.org/10.1007/s00253-013-5055-y>
- Arca-Ramos A, Eibes G, Feijoo G, Lema JM, Moreira MT (2015) Potentiality of a ceramic membrane reactor for the laccase-catalyzed removal of bisphenol A from secondary effluents. *Appl Microbiol Biotechnol* 99:9299–9308. <https://doi.org/10.1007/s00253-015-6826-4>
- Asif MB, Hai FI, Hou J, Price WE, Nghiem LD (2017a) Impact of wastewater derived dissolved interfering compounds on growth, enzymatic activity and trace organic contaminant removal of white rot fungi—a critical review. *J Environ Manag* 201:89–109
- Asif MB, Hai FI, Kang J, van de Merwe JP, Leusch FDL, Yamamoto K, Price WE, Nghiem LD (2017b) Degradation of trace organic contaminants by a membrane distillation—enzymatic bioreactor 7(879):1–15. <https://doi.org/10.3390/app7090879>
- Auriol M, Filali-Meknassi Y, Adams CD, Tyagi RD, Noguero TN, Pina B (2008) Removal of estrogenic activity of natural and synthetic hormones from a municipal wastewater: efficiency of horseradish peroxidase and laccase from *Trametes versicolor*. *Chemosphere* 70(3):445–452. <https://doi.org/10.1016/j.chemosphere.2007.06.064>
- Baiocco P, Barreca AM, Fabbri M, Galli C, Gentili P (2003) Promoting laccase activity towards non-phenolic substrates: a mechanistic investigation with some laccase-mediator systems. *Org Biomol Chem* 1(1):191–197. <https://doi.org/10.1039/B208951c>
- Barnabe S, Brar SK, Tyagi RD, Beauchesne I, Surampalli RY (2009) Pre-treatment and bioconversion of wastewater sludge to value-added products—fate of endocrine disrupting compounds. *Sci Total Environ* 407:1471–1488. <https://doi.org/10.1016/j.scitotenv.2008.11.015>
- Bhandari RK, Deem SL, Holliday DK, Jandegian CM, Kassotis CD, Nagel SC, Tillitt DE, Vom Saal FS, Rosenfeld CS (2015) Effects of the environmental estrogenic contaminants bisphenol A and 17 α -ethinyl estradiol on sexual

- development and adult behaviors in aquatic wildlife species. *Gen Comp Endocrinol* 214:195–219. <https://doi.org/10.1016/j.ygcen.2014.09.014>
- Bollag JM, Shuttleworth KL, Anderson DH (1988) Laccase-mediated detoxification of phenolic compounds. *Appl Environ Microbiol* 54(12):3086–3091
- Bourbonnais R, Paice MG (1990) Oxidation of non-phenolic substrates: an expanded role for laccase in lignin biodegradation. *FEBS Lett* 267(1):99–102. [https://doi.org/10.1016/0014-5793\(90\)80298-W](https://doi.org/10.1016/0014-5793(90)80298-W)
- Carmona E, Andreu V, Picó Y (2014) Occurrence of acidic pharmaceuticals and personal care products in Turia River Basin: from waste to drinking water. *Sci Total Environ* 484:53–63
- Catapano M, Nicolucci C, Menale C, Mita L, Rossi S, Mita DG, Diano N (2013) Enzymatic removal of estrogenic activity of nonylphenol and octylphenol aqueous solutions by immobilized laccase from *Trametes versicolor*. *J Hazard Mater* 248:337–346. <https://doi.org/10.1016/j.jhazmat.2013.01.031>
- Chairin T, Nitheranont T, Watanabe A, Asada Y, Khanongnuch C, Lumyong S (2013) Biodegradation of bisphenol A and decolorization of synthetic dyes by laccase from white-rot fungus, *Trametes polyzona*. *Appl Biochem Biotechnol* 169:539–545. <https://doi.org/10.1007/s12010-012-9990-3>
- Commission 24 August 2013. Directive 2013/39/EU of the European Parliament and of the Council of 12 August 2013 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy. Luxembourg: Off J Eur Union 226:1–17
- Commission 24 March 2015. Commission implementing decision (EU) 2015/495 of 20 March 2015 establishing a watch list of substances for Union-wide monitoring in the field of water policy pursuant to Directive 2008/105/EC of the European Parliament and of the Council. Luxembourg: Off J Eur Union 78:40–42
- Davies NM, Anderson KE (1997) Clinical pharmacokinetics of diclofenac. Therapeutic insights and pitfalls. *Clin Pharmacokinet* 33(3):184–213
- Deblonde T, Cossu-Leguille C, Hartemann P (2011) Emerging pollutants in wastewater: a review of the literature. *Int J Hyg Environ Health* 214(6):442–448. <https://doi.org/10.1016/j.ijheh.2011.08.002>
- Dorado P, Berecz R, Cáceres MC, Llerena A (2003) Analysis of diclofenac and its metabolites by high-performance liquid chromatography: relevance of CYP2C9 genotypes in diclofenac urinary metabolic ratios. *J Chromatogr B* 789:437–442
- Faber H, Melles D, Brauckmann C, Wehe CA, Wentker K, Karst U (2012) Simulation of the oxidative metabolism of diclofenac by electrochemistry/ (liquid chromatography/) mass spectrometry. *Anal Bioanal Chem* 403(2):345–354. <https://doi.org/10.1007/s00216-011-5665-0>
- Frasconi M, Favero G, Boer H, Koivula A, Mazzei F (2010) Kinetic and biochemical properties of high and low redox potential laccases from fungal and plant origin. *Biochim Biophys Acta* 1804:899–908
- Fukuda T, Uchida H, Takashima Y, Uwajima T, Kawabata T, Suzuki M (2001) Degradation of bisphenol A by purified laccase from *Trametes villosa*. *Biochem Biophys Res Commun* 284(3):704–706. <https://doi.org/10.1006/bbrc.2001.5021>
- Fukuda T, Uchida H, Suzuki M, Miyamoto H, Morinaga H, Nawata H, Uwajima T (2004) Transformation products of bisphenol A by a recombinant *Trametes villosa* laccase and their estrogenic activity. *J Chem Technol Biot* 79:1212–1218
- Galli C, Gentili P, Jolivald C, Madzak C, Vadalà R (2011) How is the reactivity of laccase affected by single-point mutations? Engineering laccase for improved activity towards sterically demanding substrates. *Appl Microbiol Biotechnol* 91:123–131. <https://doi.org/10.1007/s00253-011-3240-4>
- Gracia-Lor E, Sancho JV, Serrano R, Hernández F (2012) Occurrence and removal of pharmaceuticals in wastewater treatment plants at the Spanish Mediterranean area of Valencia. *Chemosphere* 87:453–462
- Hahn V, Mikolasch A, Wende K, Bartrow H, Lindequist U, Schauer F (2009) Synthesis of model morpholine derivatives with biological activities by laccase-catalysed reactions. *Biotechnol Appl Biochem* 54:187–195. <https://doi.org/10.1042/Ba20090219>
- Hahn V, Mikolasch A, Schauer F (2014) Cleavage and synthesis function of high and low redox potential laccases towards 4-morpholinoaniline and aminated as well as chlorinated phenols. *Appl Microbiol Biotechnol* 98(4):1609–1620. <https://doi.org/10.1007/s00253-013-4984-9>
- Heberer T (2002a) Occurrence, fate, and removal of pharmaceutical residues in the aquatic environment: a review of recent research data. *Toxicol Lett* 131(1–2):5–17
- Heberer T (2002b) Tracking persistent pharmaceutical residues from municipal sewage to drinking water. *J Hydrol* 266(3–4):175–189
- Heberer T, Feldmann D (2005) Contribution of effluents from hospitals and private households to the total loads of diclofenac and carbamazepine in municipal sewage effluents-modeling versus measurements. *J Hazard Mater* 122(3):211–218. <https://doi.org/10.1016/j.jhazmat.2005.03.007>
- Imura Y, Hartikainen P, Tatsumi K (1996) Dechlorination of tetrachloroquaiacol by laccase of white-rot basidiomycete *Coriolus versicolor*. *Appl Microbiol Biotechnol* 45:434–439
- Jonas U, Hammer E, Schauer F, Bollag JM (1997) Transformation of 2-hydroxy-dibenzofuran by laccases of the white rot fungi *Trametes versicolor* and *Pycnoporus cinnabarinus* and characterization of oligomerization products. *Biodegradation* 8(5):321–328. <https://doi.org/10.1023/a:1008220120431>
- Keilin D, Mann T (1939) Laccase, a blue copper-protein oxidase from the latex *Rhus succedanea*. *Nature* 143:23–24
- Kim Y-J, Nicell JA (2006) Impact of reaction conditions on the laccase-catalyzed conversion of bisphenol A. *Bioresour Technol* 97:1431–1442. <https://doi.org/10.1016/j.biortech.2005.06.017>
- Kordon K, Mikolasch A, Schauer F (2010) Oxidative dehalogenation of chlorinated hydroxybiphenyls by laccases of white-rot fungi. *Int Biodeterior Biodegrad* 64(3):203–209. <https://doi.org/10.1016/j.ibiod.2009.10.010>
- Kunamneni A, Camarero S, Garcia-Burgos C, Plou FJ, Ballesteros A, Alcalde M (2008) Engineering and applications of fungal laccases for organic synthesis. *Microb Cell Fact* 7:1–17
- Levy G, Lutz I, Krüger A, Kloas W (2004) Bisphenol A induces feminization in *Xenopus laevis* tadpoles. *Environ Res* 94(1):102–111. [https://doi.org/10.1016/S0013-9351\(03\)00086-0](https://doi.org/10.1016/S0013-9351(03)00086-0)
- Li KC, Xu F, Eriksson KEL (1999) Comparison of fungal laccases and redox mediators in oxidation of a nonphenolic lignin model compound. *Appl Environ Microbiol* 65:2654–2660
- Lloret L, Eibes G, Lú-Chau TA, Moreira MT, Feijoo G, Lema JM (2010) Laccase-catalyzed degradation of anti-inflammatories and estrogens. *Biochem Eng J* 51:124–131
- Loos R, Gawlik BM, Locoro G, Rimaviciute E, Contini S, Bidoglio G (2009) EU-wide survey of polar organic persistent pollutants in European river waters. *Environ Pollut* 157(2):561–568. <https://doi.org/10.1016/j.envpol.2008.09.020>
- Loos R, Locoro G, Comero S, Contini S, Schwesig D, Werres F, Balsa P, Gans O, Weiss S, Blaha L, Bolchi M, Gawlik BM (2010) Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water. *Water Res* 44(14):4115–4126. <https://doi.org/10.1016/j.watres.2010.05.032>
- Mai C, Schormann W, Hüttermann A (2001) Chemo-enzymatically induced copolymerization of phenolics with acrylate compounds. *Appl Microbiol Biotechnol* 55:177–186
- Manda K, Gordes D, Mikolasch A, Hammer E, Schmidt E, Thurow K, Schauer F (2007) Carbon-oxygen bond formation by fungal laccases: cross-coupling of 2,5-dihydroxy-N-(2-hydroxyethyl)-benzamide with the solvents water, methanol, and other alcohols. *Appl Microbiol Biotechnol* 76(2):407–416. <https://doi.org/10.1007/s00253-007-1024-7>
- Marco-Urrea E, Perez-Trujillo M, Cruz-Morato C, Caminal G, Vicent T (2010) Degradation of the drug sodium diclofenac by *Trametes versicolor* pellets and identification of some intermediates by NMR. *J Hazard Mater* 176(1–3):836–842. <https://doi.org/10.1016/j.jhazmat.2009.11.112>
- Margot J, Maillard J, Rossi L, Barry DA, Holliger C (2013) Influence of treatment conditions on the oxidation of micropollutants by *Trametes versicolor* laccase. *N Biotechnol* 30(6):803–813. <https://doi.org/10.1016/j.nbt.2013.06.004>
- Mehinto AC, Hill EM, Tyler CR (2010) Uptake and biological effects of environmentally relevant concentrations of the nonsteroidal anti-inflammatory pharmaceutical diclofenac in rainbow trout (*Oncorhynchus mykiss*). *Environ Sci Technol* 44(6):2176–2182. <https://doi.org/10.1021/es903702m>
- Mikolasch A, Schauer F (2009) Fungal laccases as tools for the synthesis of new hybrid molecules and biomaterials. *Appl Microbiol Biotechnol* 82(4):605–624. <https://doi.org/10.1007/s00253-009-1869-z>
- Minard RD, Liu SY, Bollag JM (1981) Oligomers and quinones from 2,4-dichlorophenol. *J Agr Food Chem* 29(2):250–253
- Mogharabi M, Faramarzi MA (2014) Laccase and laccase-mediated systems in the synthesis of organic compounds. *Adv Synth Catal* 356:897–927. <https://doi.org/10.1002/adsc.201300960>

- Munk L, Andersen ML, Meyer AS (2017) Direct rate assessment of laccase catalysed radical formation in lignin by electron paramagnetic resonance spectroscopy. *Enzyme Microb Technol* 106:88–96
- Nadanaciva S, Aleo MD, Strock CJ, Stedman DB, Wang H, Will Y (2013) Toxicity assessments of nonsteroidal anti-inflammatory drugs in isolated mitochondria, rat hepatocytes, and zebrafish show good concordance across chemical classes. *Toxicol Appl Pharmacol* 272(2):272–280. <https://doi.org/10.1016/j.taap.2013.06.019>
- Nagai M, Sato T, Watanabe H, Saito K, Kawata M, Enei H (2002) Purification and characterization of an extracellular laccase from the edible mushroom *Lentinula edodes*, and decolorization of chemically different dyes. *Appl Microbiol Biotechnol* 60(3):327–335. <https://doi.org/10.1007/s00253-002-1109-2>
- Nair RR, Demarche P, Agathos SN (2013) Formulation and characterization of an immobilized laccase biocatalyst and its application to eliminate organic micropollutants in wastewater. *N Biotechnol* 30(6):814–823. <https://doi.org/10.1016/j.nbt.2012.12.004>
- Nakamura T (1960) On the process of enzymatic oxidation of hydroquinone. *Biochem Biophys Res Commun* 2:111–113
- Nguyen LN, Hai FI, Price WE, Leusch FDL, Roddick F, McAdam EJ, Magram SF, Nghiem LD (2014) Continuous biotransformation of bisphenol A and diclofenac by laccase in an enzymatic membrane reactor. *Int Biodeterior Biodegrad* 95:25–32
- Quednow K, Püttmann W (2008) Endocrine disruptors in freshwater streams of Hesse, Germany: changes in concentration levels in the time span from 2003 to 2005. *Environ Pollut* 152:476–483
- Ramírez-Cavazos LI, Junghanns C, Ornelas-Soto N, Cárdenas-Chávez DL, Hernández-Luna C, Demarche P, Enaud E, García-Morales R, Agathos SN, Parra R (2014) Purification and characterization of two thermostable laccases from *Pycnoporus sanguineus* and potential role in degradation of endocrine disrupting chemicals. *J Mol Catal B Enzym* 108:32–42
- Reinhammar BRM (1972) Oxidation-reduction potentials of the electron acceptors in laccases and stellacyanin. *Biochim Biophys Acta* 275:242–245
- Samaras VG, Stasinakis AS, Mamais D, Thomaidis NS, Lekkas TD (2013) Fate of selected pharmaceuticals and synthetic endocrine disrupting compounds during wastewater treatment and sludge anaerobic digestion. *J Hazard Mater* 244–245:259–267
- Shen S, Marchick MR, Davis MR, Doss GA, Pohl LR (1999) Metabolic activation of diclofenac by human cytochrome P450 3A4: role of 5-hydroxydiclofenac. *Chem Res Toxicol* 12(2):214–222
- Solomon EI, Chen P, Metz M, Lee SK, Palmer AE (2001) Oxygen binding, activation, and reduction to water by copper proteins. *Angew Chem Int Ed* 40(24):4570–4590
- Solomon EI, Augustine AJ, Yoon J (2008) O₂ reduction to H₂O by the multicopper oxidases. *Dalton Trans* 30:3921–3932. <https://doi.org/10.1039/b800799c>
- Tadesse MA, D'Annibale A, Galli C, Gentili P, Sergi F (2008) An assessment of the relative contributions of redox and steric issues to laccase specificity towards putative substrates. *Org Biomol Chem* 6(5):868–878. <https://doi.org/10.1039/b716002j>
- Ternes TA (1998) Occurrence of drugs in German sewage treatment plants and rivers. *Water Res* 32(11):3245–3260
- Thurston CF (1994) The structure and function of fungal laccases. *Microbiology* 140:19–26
- Tran NH, Uruse T, Kusakabe O (2010) Biodegradation characteristics of pharmaceutical substances by whole fungal culture *Trametes versicolor* and its laccase. *J Water Environ Technol* 8(2):125–140
- Uchida H, Fukuda T, Miyamoto H, Kawabata T, Suzuki M, Uwajima T (2001) Polymerization of bisphenol A by purified laccase from *Trametes villosa*. *Biochem Biophys Res Commun* 287(2):355–358. <https://doi.org/10.1006/bbrc.2001.5593>
- Webster R, Pacey M, Winchester T, Johnson P, Jezequel S (1998) Microbial oxidative metabolism of diclofenac: production of 4'-hydroxydiclofenac using *Epicoccum nigrum* IMI354292. *Appl Microbiol Biotechnol* 49(4):371–376
- Wesenberg D, Kyriakides I, Agathos SN (2003) White-rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnol Adv* 22(1–2):161–187
- Xu F (1996) Oxidation of phenols, anilines, and benzenethiols by fungal laccases: correlation between activity and redox potentials as well as halide inhibition. *Biochemistry* 35(23):7608–7614. <https://doi.org/10.1021/bi952971a>
- Xu F, Berka RM, Wahleithner JA, Nelson BA, Shuster JR, Brown SH, Palmer AE, Solomon EI (1998) Site-directed mutations in fungal laccase: effect on redox potential, activity and pH profile. *Biochem J* 334:63–70
- Xu F, Kulys JJ, Duke K, Li KC, Krikstopaitis K, Deussen HJW, Abbate E, Galinyte V, Schneider P (2000) Redox chemistry in laccase-catalyzed oxidation of N-hydroxy compounds. *Appl Environ Microbiol* 66:2052–2056
- Zwiener C, Glauner T, Frimmel FH (2000) Biodegradation of pharmaceutical residues investigated by SPE-GC/ITD-MS and on-line derivatization. *J High Resolut Chrom* 23(7/8):474–478

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com