

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Leibniz-Institut im Forschungsverbund Berlin e. V.

Preprint

ISSN 0946 – 8633

**Analysis and optimal boundary control of a nonstandard
system of phase field equations**

Pierluigi Colli¹, Gianni Gilardi¹, Jürgen Sprekels²

submitted: January 30, 2012

¹ Dipartimento di Matematica “F. Casorati”
Università di Pavia
Via Ferrata, 1
27100 Pavia
Italy
E-Mail: pierluigi.colli@unipv.it
gianni.gilardi@unipv.it

² Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: juergen.sprekels@wias-berlin.de

No. 1681
Berlin 2012



2010 *Mathematics Subject Classification.* 74A15, 35K55, 49K20.

Key words and phrases. Nonlinear phase field systems, Cahn–Hilliard systems, parabolic systems, optimal boundary control, first-order necessary optimality conditions.

P. Colli and G. Gilardi gratefully acknowledge the financial support of the MIUR-PRIN Grant 2008ZKHAWN “*Phase transitions, hysteresis and multiscaling*” and of the IMATI of CNR in Pavia. The work of J. Sprekels was supported by the DFG Research Center MATHEON in Berlin.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 2044975
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Abstract

We investigate a nonstandard phase field model of Cahn-Hilliard type. The model, which was introduced in [16], describes two-species phase segregation and consists of a system of two highly nonlinearly coupled PDEs. It has been studied recently in [5], [6] for the case of homogeneous Neumann boundary conditions. In this paper, we investigate the case that the boundary condition for one of the unknowns of the system is of third kind and nonhomogeneous. For the resulting system, we show well-posedness, and we study optimal boundary control problems. Existence of optimal controls is shown, and the first-order necessary optimality conditions are derived. Owing to the strong nonlinear couplings in the PDE system, standard arguments of optimal control theory do not apply directly, although the control constraints and the cost functional will be of standard type.

1 Introduction

Let $\Omega \subset \mathbb{R}^3$ denote an open and bounded domain whose smooth boundary Γ has outward unit normal \mathbf{n} , let $T > 0$ be a given final time, and let $Q := \Omega \times (0, T)$, $\Sigma := \Gamma \times (0, T)$. In this paper, we study the following initial-boundary value problem:

$$(\varepsilon + 2\rho)\mu_t + \mu\rho_t - \Delta\mu = 0 \quad \text{a. e. in } Q, \quad (1.1)$$

$$\delta\rho_t - \Delta\rho + f'(\rho) = \mu \quad \text{a. e. in } Q, \quad (1.2)$$

$$\frac{\partial\rho}{\partial\mathbf{n}} = 0, \quad \frac{\partial\mu}{\partial\mathbf{n}} = \alpha(u - \mu) \quad \text{a. e. on } \Sigma, \quad (1.3)$$

$$\rho(x, 0) = \rho_0(x), \quad \mu(x, 0) = \mu_0(x), \quad \text{for a. e. } x \in \Omega. \quad (1.4)$$

The PDE system (1.1)–(1.2) constitutes a phase field model of Cahn-Hilliard type that describes phase segregation of two species (atoms and vacancies, say) on a lattice in the presence of diffusion. It has been introduced recently in [16] and [5]; for the general physical background, we refer the reader to [16]. The unknown variables are the *order parameter* ρ , interpreted as a volumetric density, and the *chemical potential* μ . For physical reasons, we must have $0 \leq \rho \leq 1$ and $\mu > 0$ almost everywhere in Q . The boundary (control) function u on the right-hand side of (1.3)₂ plays the role of a *microenergy source*. Moreover, ε and δ are positive constants, and the nonlinearity f is a double-well potential defined in $(0, 1)$, whose derivative f' is singular at the endpoints $\rho = 0$ and $\rho = 1$; a typical example is $f = f_1 + f_2$, with f_2 smooth and $f_1(\rho) = c(\rho \log(\rho) + (1 - \rho) \log(1 - \rho))$, where c is a positive constant.

The PDE system (1.1)–(1.4) is singular, with highly nonlinear and nonstandard coupling. In particular, unpleasant nonlinear couplings involving time derivatives occur in (1.1), and the expression $f'(\rho)$ in (1.2) may become singular. In the recent papers [5], [6], well-posedness and asymptotic behavior for $t \rightarrow \infty$ and $\varepsilon \searrow 0$ of the system (1.1)–(1.4) were established for the

case when the second boundary condition in (1.3) is replaced by the homogeneous Neumann boundary condition $\partial\mu/\partial\mathbf{n} = 0$; a distributed optimal control problem for this situation was analyzed in [7]. We also refer to the papers [3] and [4], where the corresponding Allen-Cahn model was discussed.

The paper is organized as follows: in Section 2, we state the general assumptions and prove the existence of a strong solution to the problem. Section 3 is concerned with the issues of uniqueness and stability. Section 4 then brings the study of a boundary control problem for the system (1.1)–(1.4). We show existence of a solution to the optimal control problem and derive the first-order necessary optimality conditions, as usual given in terms of the adjoint system and a variational inequality.

Throughout the paper, we make repeated use of Hölder's inequality, of the elementary Young inequality

$$ab \leq \gamma a^2 + \frac{1}{4\gamma} b^2, \quad \text{for every } a, b \geq 0 \text{ and } \gamma > 0, \quad (1.5)$$

of the interpolation inequality

$$\|v\|_{L^r(\Omega)} \leq \|v\|_{L^p(\Omega)}^\theta \|v\|_{L^q(\Omega)}^{1-\theta} \quad \forall v \in L^p(\Omega) \cap L^q(\Omega),$$

where $p, q, r \in [1, +\infty]$, $\theta \in [0, 1]$, and $\frac{1}{r} = \frac{\theta}{p} + \frac{1-\theta}{q}$, (1.6)

and, since $\dim \Omega \leq 3$, of the continuity of the embeddings $H^1(\Omega) \subset L^q(\Omega)$ for $1 \leq q \leq 6$, where, with constants $\hat{C}_q > 0$ depending only on Ω ,

$$\|v\|_{L^q(\Omega)} \leq \hat{C}_q \|v\|_{H^1(\Omega)} \quad \forall v \in H^1(\Omega), \quad 1 \leq q \leq 6, \quad (1.7)$$

and where the embeddings are compact for $1 \leq q < 6$. We also use the Sobolev spaces $H^s(\Omega)$ of real order $s > 0$ and recall the compact embeddings $H^s(\Omega) \subset H^1(\Omega)$ and $H^s(\Omega) \subset C(\bar{\Omega})$ for $s > 1$ and $s > 3/2$, respectively, and, e. g., the estimate, with a constant $\hat{C}_\infty > 0$ depending only on Ω ,

$$\|v\|_{C(\bar{\Omega})} \leq \hat{C}_\infty \|v\|_{H^2(\Omega)} \quad \forall v \in H^2(\Omega). \quad (1.8)$$

2 Problem statement and existence

Consider the initial-boundary value problem (1.1)–(1.4). For convenience, we introduce the abbreviated notation

$$H = L^2(\Omega), \quad V = H^1(\Omega), \quad W = \{w \in H^2(\Omega) : \partial w / \partial \mathbf{n} = 0 \text{ on } \Gamma\}.$$

We endow these spaces with their standard norms, for which we use self-explaining notation like $\|\cdot\|_V$; for simplicity, we also write $\|\cdot\|_H$ for the norm in the space $H \times H \times H$. Recall that the embeddings $W \subset V \subset H$ are compact. Moreover, since V is dense in H , we can identify H with a subspace of V^* in the usual way, i. e., by setting $\langle u, v \rangle_{V^*, V} = (u, v)_H$ for all $u \in H$ and $v \in V$, where $\langle \cdot, \cdot \rangle_{V^*, V}$ denotes the duality pairing between V^* and V . Then also the embedding $H \subset V^*$ is compact.

We make the following assumptions on the data:

(A1) $f = f_1 + f_2$, where $f_1 \in C^2(0, 1)$ is convex, $f_2 \in C^2[0, 1]$, and

$$\lim_{r \searrow 0} f_1'(r) = -\infty, \quad \lim_{r \nearrow 1} f_1'(r) = +\infty. \quad (2.1)$$

(A2) $\rho_0 \in W$, $f'(\rho_0) \in H$, $\mu_0 \in V$, and

$$0 < \rho_0(x) < 1 \quad \forall x \in \Omega, \quad \mu_0 \geq 0 \quad \text{a. e. in } \Omega. \quad (2.2)$$

(A3) $u \in H^1(0, T; L^2(\Gamma))$, and $u \geq 0$ a. e. on Σ .

(A4) $\alpha \in L^\infty(\Gamma)$, and $\alpha(x) \geq \alpha_0 > 0$ for almost every $x \in \Gamma$.

Notice that (A2) implies that $\rho_0 \in C(\bar{\Omega})$ and, thanks to the convexity of f_1 , also that $f(\rho_0) \in H$.

The following existence result resembles that of Theorem 2.1 in [5].

Theorem 2.1 *Suppose that the hypotheses (A1)–(A4) are satisfied. Then the system (1.1)–(1.4) has a solution (ρ, μ) such that*

$$\rho \in W^{1,\infty}(0, T; H) \cap H^1(0, T; V) \cap L^\infty(0, T; W), \quad (2.3)$$

$$\mu \in H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; H^{3/2}(\Omega)), \quad (2.4)$$

$$f'(\rho) \in L^\infty(0, T; H), \quad (2.5)$$

$$0 < \rho < 1 \quad \text{a. e. in } Q, \quad \mu \geq 0 \quad \text{a. e. in } Q. \quad (2.6)$$

Remark 2.2 The $H^{3/2}$ space regularity for μ is optimal due to the L^2 space regularity of u given by (A3). Nevertheless, both equation (1.1) and the boundary condition for μ contained in (1.3) can be understood a.e. in Q and a.e. on Σ , respectively, and the standard integration by parts is correct, as we briefly explain (so that we can both refer to that formulation and use integration by parts). In principle, one can replace the equation and the boundary condition by the usual variational formulation, namely

$$\int_{\Omega} [(\varepsilon + 2\rho)\mu_t + \mu\rho_t] v \, dx + \int_{\Omega} \nabla\mu \cdot \nabla v \, dx + \int_{\Gamma} \alpha(\mu - u)v \, d\sigma = 0$$

(where $d\sigma$ stands for the surface measure) for every $v \in V$, a.e. in $(0, T)$, or an integrated-in-time version of it. This implies that (1.1) is satisfied in the sense of distributions, whence $\Delta\mu$ belongs to $L^2(Q)$ by comparison, and the equation can be understood a.e. in Q , a posteriori. The last regularity (2.4) of μ and the condition $\Delta\mu \in L^2(0, T; L^2(\Omega))$ just observed also ensure that the trace $\frac{\partial\mu}{\partial\mathbf{n}}|_{\Sigma}$ has a meaning in the space $L^2(0, T; L^2(\Gamma))$ due to the trace theorem [15, Thm. 7.3] (we just observe that the space $\Xi^{-1/2}(\Omega)$ that enters such a result is larger than $L^2(\Omega)$), so that the boundary condition can be read a.e. on Σ .

Proof of Theorem 2.1. The proof follows closely the lines of the proof of Theorem 2.1 in [5], where a homogeneous Neumann boundary condition for μ was investigated.

Step 1: Approximation. We employ an approximation scheme based on a time delay in the right-hand side of (1.2). To this end, we introduce for $\tau > 0$ the translation operator $\mathcal{T}_\tau : L^1(0, T; H) \rightarrow L^1(0, T; H)$, which for $v \in L^1(0, T; H)$ and almost every $t \in (0, T)$ is defined by

$$(\mathcal{T}_\tau)(t) := v(t - \tau) \quad \text{if } t > \tau, \quad \text{and } (\mathcal{T}_\tau)(t) := \mu_0 \quad \text{if } t \leq \tau. \quad (2.7)$$

Now, let $N \in \mathbb{N}$ be arbitrary, and $\tau := T/N$. We seek functions (ρ^τ, μ^τ) satisfying (2.3)–(2.6) (with (ρ, μ) replaced by (ρ^τ, μ^τ)), which solve the system

$$(\varepsilon + 2\rho^\tau)\mu_t^\tau + \mu^\tau \rho_t^\tau - \Delta\mu^\tau = 0 \quad \text{a. e. in } Q, \quad (2.8)$$

$$\delta\rho_t^\tau - \Delta\rho^\tau + f'(\rho^\tau) = \mathcal{T}_\tau\mu^\tau \quad \text{a. e. in } Q, \quad (2.9)$$

$$\frac{\partial\rho^\tau}{\partial\mathbf{n}} = 0, \quad \frac{\partial\mu^\tau}{\partial\mathbf{n}} = \alpha(u - \mu^\tau) \quad \text{a. e. on } \Sigma, \quad (2.10)$$

$$\rho^\tau(x, 0) = \rho_0(x), \quad \mu^\tau(x, 0) = \mu_0(x), \quad \text{for a. e. } x \in \Omega. \quad (2.11)$$

We note that Remark 2.2 also applies to the approximating problem. To prove the existence of a solution, we put $t_n := n\tau$, $I_n := [0, t_n]$, $1 \leq n \leq N$, and consider for $1 \leq n \leq N$ the problem

$$(\varepsilon + 2\rho^n)\mu_t^n + \mu^n \rho_t^n - \Delta\mu^n = 0 \quad \text{a. e. in } \Omega \times I_n, \quad (2.12)$$

$$\mu^n(0) = \mu_0 \quad \text{a. e. in } \Omega, \quad \frac{\partial\mu^n}{\partial\mathbf{n}} = \alpha(u - \mu^n) \quad \text{a. e. on } \Gamma \times I_n, \quad (2.13)$$

$$\delta\rho_t^n - \Delta\rho^n + f'(\rho^n) = \mathcal{T}_\tau\mu^{n-1} \quad \text{a. e. in } \Omega \times I_n, \quad (2.14)$$

$$\rho^n(0) = \rho_0 \quad \text{a. e. in } \Omega, \quad \frac{\partial\rho^n}{\partial\mathbf{n}} = 0, \quad \text{a. e. on } \Gamma \times I_n. \quad (2.15)$$

Notice that the operator \mathcal{T}_τ acts on functions that are not defined on the entire interval $(0, T)$. However, its meaning is still given by (2.7) if $n > 1$, and for $n = 1$ we simply put $\mathcal{T}_\tau\mu^{n-1} = \mu_0$.

Clearly, we have $(\rho^\tau, \mu^\tau) = (\rho^N, \mu^N)$ if (ρ^N, μ^N) exists. We claim that the systems (2.12)–(2.15) can be uniquely solved by induction for $n = 1, \dots, N$, where, for $1 \leq n \leq N$,

$$\rho^n \in W^{1,\infty}(I_n; H) \cap H^1(I_n; V) \cap L^\infty(I_n; W), \quad (2.16)$$

$$\mu^n \in H^1(I_n; H) \cap C^0(I_n; V) \cap L^2(I_n; H^{3/2}(\Omega)), \quad (2.17)$$

$$0 < \rho^n < 1 \quad \text{a. e. in } \Omega \times I_n, \quad \mu^n \geq 0 \quad \text{a. e. in } \Omega \times I_n. \quad (2.18)$$

To prove the claim, suppose that for some $n \in \{1, \dots, N\}$ the problem (2.12)–(2.15) has a unique solution satisfying (2.16)–(2.18), where the index n is replaced by $n - 1$. Then it follows with exactly the same argument as in the proof of Theorem 2.1 in [5] that the initial-boundary value problem (2.14), (2.15) has a unique solution ρ^n that satisfies (2.16) and the first inequality in (2.18). Substituting ρ^n in (2.12), we infer that the linear initial-boundary value problem (2.12), (2.13) has a unique solution μ^n satisfying (2.17). Notice here that the regularity of μ^n follows from the fact that $u \in H^1(0, T; L^2(\Gamma))$.

It remains to show that μ^n is nonnegative almost everywhere. To this end, we test (2.12) by $-(\mu^n)^-$, where $(\mu^n)^-$ denotes the negative part of μ^n . Using integration by parts and the boundary condition in (2.13), we obtain the identity

$$\begin{aligned} & \frac{1}{2} \int_0^t \int_{\Omega} \frac{d}{dt} \left((\varepsilon + 2\rho^n) |(\mu^n)^-|^2 \right) dx ds + \int_0^t \int_{\Omega} |\nabla(\mu^n)^-|^2 dx ds \\ & + \int_0^t \int_{\Gamma} \alpha |(\mu^n)^-|^2 d\sigma ds + \int_0^t \int_{\Gamma} \alpha u (\mu^n)^- d\sigma ds = 0. \end{aligned}$$

From the fact that $\rho^n, \rho_0, \mu_0, \alpha, u$ are all nonnegative, we infer that

$$\begin{aligned} \varepsilon \int_{\Omega} |(\mu^n)^-(t)|^2 dx & \leq \int_{\Omega} (\varepsilon + 2\rho^n(t)) |(\mu^n)^-(t)|^2 dx \\ & \leq \int_{\Omega} (\varepsilon + 2\rho_0) |\mu_0^-|^2 dx = 0. \end{aligned}$$

Hence, $(\mu^n)^- = 0$, i. e., $\mu^n \geq 0$ a. e. in $\Omega \times I_n$, and the claim is proved.

Step 2: A priori estimates. Now that the well-posedness of the problem (2.8)–(2.11) is established, we perform a number of a priori estimates for its solution. For the sake of a better readability, we will omit the index τ in the calculations. In what follows, we denote by $C > 0$ positive constants that may depend on the data of the system but not on τ . The meaning of C may change from line to line and even in the same chain of inequalities.

First estimate. Since $\partial_t((\varepsilon/2)\mu^2 + \rho\mu^2) = ((\varepsilon + 2\rho)\mu_t + \mu\rho_t)\mu$, testing of (2.8) by μ yields, for every $t \in [0, T]$,

$$\begin{aligned} & \int_{\Omega} \left(\frac{\varepsilon}{2} \mu^2 + \rho\mu^2 \right) (t) dx + \int_0^t \int_{\Omega} |\nabla\mu|^2 dx ds + \int_0^t \int_{\Gamma} \alpha \mu^2 d\sigma ds \\ & = \int_{\Omega} \left(\frac{\varepsilon}{2} \mu_0^2 + \rho_0\mu_0^2 \right) (t) dx + \int_0^t \int_{\Gamma} \alpha u \mu d\sigma ds, \end{aligned}$$

whence, using Young's inequality and (A2)–(A4), we can conclude that

$$\|\mu\|_{L^\infty(0,T;H) \cap L^2(0,T;V)} \leq C. \quad (2.19)$$

Second estimate. Next, we test (2.9) by ρ_t . Applying (2.19), recalling the fact that $f(\rho_0) \in H$, and invoking Young's inequality, we easily see that

$$\|\rho\|_{H^1(0,T;H) \cap L^\infty(0,T;V)} + \|f(\rho)\|_{L^\infty(0,T;L^1(\Omega))} \leq C. \quad (2.20)$$

Third estimate. We rewrite Eq. (2.9) in the form

$$-\Delta\rho + f_1'(\rho) = -\delta\rho_t - f_2'(\rho) + \mathcal{I}_\tau\mu$$

and observe that the right-hand side is bounded in $L^2(Q)$. Hence, applying a standard procedure (e. g., testing by $f_1'(\rho)$), and invoking elliptic regularity, we find that

$$\|\rho\|_{L^2(0,T;W)} + \|f_1'(\rho)\|_{L^2(Q)} \leq C. \quad (2.21)$$

Fourth estimate. We differentiate Eq. (2.9) formally with respect to t and test the resulting equation with ρ_t (this argument can be made rigorous, see [5]). Since, owing to the convexity of f_1 , $f_1''(\rho)$ is nonnegative almost everywhere, we find the estimate

$$\begin{aligned} & \frac{\delta}{2} \|\rho_t(t)\|^2 + \int_0^t \int_{\Omega} |\nabla \rho_t|^2 dx ds \leq \frac{\delta}{2} \|\Delta \rho_0 - f_1'(\rho_0) + \mu_0\|_H^2 \\ & + \max_{0 \leq \rho \leq 1} |f_2''(\rho)| \int_0^t \int_{\Omega} |\rho_t|^2 dx ds + \int_0^t \int_{\Omega} (\partial_t \mathcal{T}_\tau \mu) \rho_t dx ds \\ & \leq C + \int_0^{t-\tau} \int_{\Omega} \mu_t(s) \rho_t(s + \tau) dx ds. \end{aligned} \quad (2.22)$$

In order to estimate the last integral, we substitute for μ_t , using Eq. (2.8). It follows, using integration by parts:

$$\begin{aligned} & \int_0^{t-\tau} \int_{\Omega} \mu_t \rho_t(\cdot + \tau) dx ds = \int_0^{t-\tau} \int_{\Omega} \frac{1}{\varepsilon + 2\rho} (\Delta \mu - \mu \rho_t) \rho_t(\cdot + \tau) dx ds \\ & = \int_0^{t-\tau} \int_{\Omega} \left[-\frac{\nabla \mu}{\varepsilon + 2\rho} \cdot \nabla \rho_t(\cdot + \tau) + \frac{2\rho_t(\cdot + \tau)}{(\varepsilon + 2\rho)^2} \nabla \mu \cdot \nabla \rho \right. \\ & \quad \left. - \frac{1}{\varepsilon + 2\rho} \rho_t \mu \rho_t(\cdot + \tau) \right] dx ds \\ & - \int_0^{t-\tau} \int_{\Gamma} \frac{\alpha}{\varepsilon + 2\rho} (u - \mu) \rho_t(\cdot + \tau) d\sigma ds. \end{aligned} \quad (2.23)$$

Exactly as in the proof of Theorem 2.1 in [5], the domain integral in the second and third lines of (2.23) can be estimated from above by an expression of the form

$$\frac{1}{2} \int_0^t \int_{\Omega} |\nabla \rho_t|^2 dx ds + C \left(1 + \int_0^t \|\mu(s)\|_V^2 \|\rho_t(s)\|_H^2 dx ds \right). \quad (2.24)$$

Observe that, owing to the inequality (2.19), the mapping $s \mapsto \|\mu(s)\|_V^2$ belongs to $L^1(0, T)$. Finally, we estimate the boundary term in the last line of Eq. (2.23). To this end, recall that by the trace theorem there is a constant $c_{\Omega} > 0$, independent of τ , such that $\|v\|_{L^2(\Gamma)} \leq c_{\Omega} \|v\|_V$ for all $v \in V$. Moreover, we have $\rho \geq 0$ and $\alpha \in L^\infty(\Gamma)$. Therefore, we obtain that

$$\begin{aligned}
& \left| \int_0^{t-\tau} \int_{\Gamma} \frac{\alpha}{\varepsilon + 2\rho} (u - \mu) \rho_t(\cdot + \tau) d\sigma ds \right| \\
& \leq C \int_0^{t-\tau} \|\rho_t(s + \tau)\|_{L^2(\Gamma)} (\|u(s)\|_{L^2(\Gamma)} + \|\mu(s)\|_{L^2(\Gamma)}) ds \\
& \leq C \int_0^{t-\tau} \|\rho_t(s + \tau)\|_V (\|u(s)\|_{L^2(\Gamma)} + \|\mu(s)\|_V) ds \\
& \leq \frac{1}{4} \int_0^t \|\rho_t(s)\|_V^2 ds + C. \tag{2.25}
\end{aligned}$$

Now we may combine the estimates (2.22)–(2.25) and employ Gronwall's inequality to conclude that

$$\|\rho_t\|_{L^\infty(0,T;H) \cap L^2(0,T;V)} \leq C. \tag{2.26}$$

The same argument as in the derivation of (2.22) then shows that also

$$\|\rho_t\|_{L^\infty(0,T;W)} + \|f_1'(\rho)\|_{L^\infty(0,T;H)} \leq C. \tag{2.27}$$

Fifth estimate. We test equation (2.8) by μ_t . Formal integration by parts (this can be made rigorous), using (A3), the trace theorem and Young's inequality, yields:

$$\begin{aligned}
& \varepsilon \int_0^t \int_{\Omega} |\mu_t|^2 dx ds + \frac{1}{2} \|\nabla \mu(t)\|_H^2 + \int_{\Gamma} \frac{\alpha}{2} |\mu(t)|^2 d\sigma \\
& \leq C + \int_0^t \int_{\Gamma} \alpha u \mu_t d\sigma + \int_0^t \int_{\Omega} |\mu \rho_t \mu_t| dx ds \\
& \leq C + \int_{\Gamma} \alpha u(t) \mu(t) d\sigma - \int_0^t \int_{\Gamma} \alpha u_t \mu d\sigma ds + \int_0^t \int_{\Omega} |\mu \rho_t \mu_t| dx ds \\
& \leq \frac{C}{\gamma} + \gamma \|\mu(t)\|_V^2 + \int_0^t \|\mu(s)\|_V^2 ds + \int_0^t \int_{\Omega} |\mu| |\rho_t| |\mu_t| dx ds \\
& \leq \frac{C}{\gamma} + \gamma \|\mu(t)\|_V^2 + \int_0^t \|\mu(s)\|_V^2 ds + \frac{\varepsilon}{2} \int_0^t \|\mu_t(s)\|_H^2 ds \\
& \quad + C \int_0^t \|\rho_t(s)\|_{L^4(\Omega)}^2 \|\mu(s)\|_{L^4(\Omega)}^2 ds \\
& \leq \frac{C}{\gamma} + \gamma \|\mu(t)\|_V^2 + \frac{\varepsilon}{2} \int_0^t \|\mu_t(s)\|_H^2 ds \\
& \quad + C \int_0^t (1 + \|\rho_t(s)\|_V^2) \|\mu(s)\|_V^2 ds. \tag{2.28}
\end{aligned}$$

Hence, using (2.26), choosing $\gamma > 0$ sufficiently small, and invoking Gronwall's lemma, we can conclude that

$$\|\mu\|_{H^1(0,T;H) \cap L^\infty(0,T;V)} \leq C. \quad (2.29)$$

Sixth estimate. Since $0 < \rho < 1$ a. e. in Q , and using (2.26), (2.29) and the continuity of the embedding $V \subset L^4(\Omega)$, we can estimate as follows:

$$\begin{aligned} \|(\varepsilon + 2\rho)\mu_t + \mu\rho_t\|_{L^2(Q)} &\leq C \|\mu_t\|_{L^2(Q)} + \|\mu\|_{L^\infty(0,T;L^4(\Omega))} \|\rho_t\|_{L^2(0,T;L^4(\Omega))} \\ &\leq C (\|\mu_t\|_{L^2(Q)} + \|\mu\|_{L^\infty(0,T;V)} \|\rho_t\|_{L^2(0,T;V)}) \leq C. \end{aligned} \quad (2.30)$$

Comparison in (2.8) then shows the boundedness of $\Delta\mu$ in $L^2(Q)$, and it follows from (2.8), (A3) and standard elliptic estimates that also

$$\|\mu\|_{L^2(0,T;H^{3/2}(\Omega))} \leq C. \quad (2.31)$$

Step 3: Conclusion of the proof. Collecting all the above estimates, it turns out that there is some sequence $\tau_k \searrow 0$ such that

$$\begin{aligned} \mu^{\tau_k} &\rightharpoonup \mu && \text{weakly star in} \\ &&& H^1(0,T;H) \cap L^\infty(0,T;V) \cap L^2(0,T;H^{3/2}(\Omega)), \\ \rho^{\tau_k} &\rightharpoonup \rho && \text{weakly star in } W^{1,\infty}(0,T;H) \cap H^1(0,T;V) \cap L^\infty(0,T;W), \\ f_1'(\rho^{\tau_k}) &\rightharpoonup \xi && \text{weakly star in } L^\infty(0,T;H). \end{aligned}$$

Thanks to the Aubin-Lions lemma (cf., [14, Thm. 5.1, p. 58]) and similar results to be found in [17, Sect. 8, Cor. 4], we also deduce (recall that even $H^{3/2}(\Omega)$ is compactly embedded into V) the strong convergences

$$\begin{aligned} \mu^{\tau_k} &\rightarrow \mu && \text{strongly in } C^0([0,T];H) \cap L^2(0,T;V), \\ \rho^{\tau_k} &\rightarrow \rho && \text{strongly in } C^0([0,T];V) \end{aligned}$$

and the Cauchy conditions (1.4) as a consequence. In particular, employing a standard monotonicity argument (cf., e.g., [1, Lemma 1.3, p. 42]), we conclude that $0 < \rho < 1$ and $\xi = f_1'(\rho)$ a. e. in Q . The strong convergence shown above also entails that $f_2'(\rho^{\tau_k}) \rightarrow f_2'(\rho)$ strongly in $C^0([0,T];H)$ (because f_2' is Lipschitz continuous), and that $\mathcal{T}_{\tau_k} \mu^{\tau_k} \rightarrow \mu$ strongly in $L^2(Q)$.

Now notice that the above convergences imply, in particular, that

$$\begin{aligned} \rho^{\tau_k} &\rightarrow \rho && \text{strongly in } C^0([0,T];L^6(\Omega)), \\ \rho_t^{\tau_k} &\rightarrow \rho_t && \text{weakly in } L^2(0,T;L^4(\Omega)), \\ \mu^{\tau_k} &\rightarrow \mu && \text{strongly in } L^2(0,T;L^4(\Omega)), \\ \mu_t^{\tau_k} &\rightarrow \mu_t && \text{weakly in } L^2(Q). \end{aligned}$$

From this, it is easily verified that

$$\begin{aligned}\mu^{\tau_k} \rho_t^{\tau_k} &\rightharpoonup \mu \rho_t && \text{weakly in } L^1(0, T; H), \\ \rho^{\tau_k} \mu_t^{\tau_k} &\rightharpoonup \rho \mu_t && \text{weakly in } L^2(0, T; L^{3/2}(\Omega)).\end{aligned}$$

Now, we are ready to take the limit as $k \rightarrow \infty$ in (2.8)–(2.10) (written for $\tau = \tau_k$). Precisely, we can do that as far as ρ is concerned, while it is easier to take the limit in the variational formulation of (2.8) that accounts for the boundary condition (the same as mentioned in Remark 2.2), or in the following integrated-in-time version of it

$$\begin{aligned}&\int_0^T \int_{\Omega} [(\varepsilon + 2\rho^\tau)\mu_t^\tau + \mu^\tau \rho_t^\tau] v \, dx \, dt + \int_0^T \int_{\Omega} \nabla \mu^\tau \cdot \nabla v \, dx \, dt \\ &+ \int_0^T \int_{\Gamma} \alpha(\mu^\tau - u)v \, d\sigma \, dt = 0 \quad \text{for every } v \in L^\infty(0, T; V).\end{aligned}$$

Then, we obtain the analogue for μ , which implies (1.1) and (1.3)₂. ■

3 Boundedness, uniqueness, and stability

In this section, we derive results concerning boundedness, uniqueness and stability of the solutions to system (1.1)–(1.4). With respect to boundedness, we have the following result, which resembles Theorem 2.3 in [5].

Theorem 3.1 *Suppose that (A1)–(A4) are fulfilled, and suppose that the following conditions are satisfied:*

$$(A5) \quad \mu_0 \in L^\infty(\Omega), \quad \inf_{x \in \Omega} \rho_0(x) > 0, \quad \sup_{x \in \Omega} \rho_0(x) < 1.$$

$$(A6) \quad u \in L^\infty(\Sigma).$$

Then any solution (ρ, μ) of (1.1)–(1.4) fulfilling (2.3)–(2.6) also satisfies

$$\mu \leq \mu^*, \quad \rho \geq \rho_*, \quad \text{and} \quad \rho \leq \rho^* \quad \text{a.e. in } Q \tag{3.1}$$

for some constants $\mu^ > 0$ and $\rho_*, \rho^* \in (0, 1)$ that depend on the structure of the system and T , on the initial data, and on an upper bound for the L^∞ norm of u , only.*

Proof. Let us just show the boundedness of μ and the first estimate (3.1); the results for ρ then follow in exactly the same manner as in the proof of Theorem 2.3 in [5]. Also the result for μ follows – up to some changes that are necessary due to the different boundary condition for μ – by the same chain of arguments as in the proof of Theorem 2.3 in [5]; but since this proof does not seem to be standard, we provide it for the reader’s convenience. So let (ρ, μ) be any solution to the system (1.1)–(1.4), (2.3)–(2.6). We set

$$\Phi_0 := \max \{1, \|\mu_0\|_{L^\infty(\Omega)}, \|u\|_{L^\infty(\Sigma)}\},$$

choose any $k \in \mathbb{R}$ such that $k \geq \Phi_0$, and introduce the auxiliary function $\chi_k \in L^\infty(Q)$ by putting, for almost every $(x, t) \in Q$,

$$\chi_k(x, t) = 1 \quad \text{if } \mu(x, t) > k, \quad \text{and } \chi_k(x, t) = 0 \quad \text{otherwise.}$$

Then, we test (1.1) by $(\mu - k)^+$. We obtain, for any $t \in [0, T]$,

$$\begin{aligned} & \int_{\Omega} \left(\frac{\varepsilon}{2} + \rho(t) \right) |(\mu(t) - k)^+|^2 + \int_0^t \int_{\Omega} |\nabla(\mu - k)^+|^2 dx ds \\ & + \int_0^t \int_{\Gamma} \alpha (\mu - u) (\mu - k)^+ d\sigma ds \\ & = \int_0^t \int_{\Omega} \rho_t |(\mu - k)^+|^2 dx ds - \int_0^t \int_{\Omega} \rho_t \mu (\mu - k)^+ dx ds \\ & = -k \int_0^t \int_{\Omega} \rho_t (\mu - k)^+ dx ds. \end{aligned}$$

Now observe that α and ρ are nonnegative and that, by definition of k ,

$$\alpha (\mu - u) (\mu - k)^+ = \alpha (|(\mu - k)^+|^2 + (k - u) (\mu - k)^+) \geq 0 \quad \text{a. e. in } Q.$$

Hence, using Hölder's inequality, we obtain from the above equality the estimate

$$\begin{aligned} & \frac{\varepsilon}{2} \|(\mu(t) - k)^+\|_H^2 + \int_0^t \int_{\Omega} |\nabla(\mu - k)^+|^2 dx ds \\ & \leq k \int_0^t \|\chi_k(s)\|_{L^{7/2}(\Omega)} \|\rho_t(s)\|_{L^{14/3}(\Omega)} \|(\mu - k)^+(s)\|_{L^2(\Omega)} ds, \end{aligned}$$

whence, using the Gronwall-Bellman lemma as in [2, Lemma A.4, p. 156],

$$\begin{aligned} & \left(\varepsilon \|(\mu - k)^+\|_{C^0([0, T]; H)}^2 + \int_0^T \int_{\Omega} |\nabla(\mu - k)^+|^2 dx dt \right)^{1/2} \\ & \leq \frac{k}{\sqrt{\varepsilon}} \int_0^T \|\chi_k(t)\|_{L^{7/2}(\Omega)} \|\rho_t(t)\|_{L^{14/3}(\Omega)} dt \\ & \leq \frac{k}{\sqrt{\varepsilon}} \|\rho_t\|_{L^{7/3}(0, T; L^{14/3}(\Omega))} \|\chi_k\|_{L^{7/4}(0, T; L^{7/2}(\Omega))}. \end{aligned} \tag{3.2}$$

Next, we apply the continuity of the embedding $V \subset L^6(\Omega)$ and the interpolation inequality (1.6) with $p = 2$, $q = 6$, $r = 14/3$, and $\theta = 1/7$. It follows that

$$\begin{aligned} \|\rho_t\|_{L^{7/3}(0, T; L^{14/3}(\Omega))} & \leq \left(\int_0^T \|\rho_t(t)\|_{L^2(\Omega)}^{1/3} \|\rho_t(t)\|_{L^6(\Omega)}^2 dt \right)^{3/7} \\ & \leq \|\rho_t\|_{L^\infty(0, T; H)}^{1/7} \left(\int_0^T \|\rho_t(t)\|_{L^6(\Omega)}^2 dt \right)^{3/7} \leq C \|\rho_t\|_{L^2(0, T; V)}^{6/7} \leq D_0, \end{aligned}$$

where D_0 is a positive constant depending only on the data of the problem. Moreover, we have

$$\begin{aligned}\|\chi_k\|_{L^{7/4}(0,T;L^{7/2}(\Omega))} &= \left[\int_0^T \left(\int_{\Omega} |\chi_k(x,t)|^{7/2} dx \right)^{1/2} dt \right]^{4/7} \\ &= \left[\int_0^T \left(\int_{\Omega} |\chi_k(x,t)|^4 dx \right)^{1/2} dt \right]^{\frac{1}{2} \cdot \frac{8}{7}} = \|\chi_k\|_{L^2(0,T;L^4(\Omega))}^{8/7}.\end{aligned}$$

Hence, we can infer from (3.2) that for every $k \geq \Phi_0$ it holds the inequality

$$\|(\mu - k)^+\| \leq k D_1 \|\chi_k\|_{L^2(0,T;L^4(\Omega))}^{8/7}, \quad (3.3)$$

where $D_1 = D_0 / \text{Min} \{\varepsilon, 1\}$, and where the norm $\|\cdot\|$ is defined by

$$\|v\|^2 := \max_{t \in [0,T]} \|v(t)\|_H^2 + \int_0^T \int_{\Omega} |\nabla v|^2 dx dt \quad \forall v \in C^0([0,T]; H) \cap L^2(0,T; V).$$

Moreover, owing to the continuity of the embedding $V \subset L^4(\Omega)$, there is some constant $D_2 > 0$, which only depends on Ω and on T , such that

$$\|v\|_{L^2(0,T;L^4(\Omega))} \leq D_2 \|v\| \quad \forall v \in C^0([0,T]; H) \cap L^2(0,T; V). \quad (3.4)$$

At this point, we select a strictly increasing sequence $\{k_j\}$ depending on a real parameter $m > 1$ as follows:

$$k_j := M(2 - 2^{-j}) \quad \text{for } j = 0, 1, \dots, \quad \text{with } M := m \Phi_0. \quad (3.5)$$

Note that $k_0 = M > \Phi_0$ and $\lim_{j \rightarrow \infty} k_j = 2M$. Then, owing to (3.3) and (3.4), it is not difficult to check that

$$\begin{aligned}(k_{j+1} - k_j) \|\chi_{k_{j+1}}\|_{L^2(0,T;L^4(\Omega))} &\leq \|(\mu - k_j)^+\|_{L^2(0,T;L^4(\Omega))} \\ &\leq D_2 \|(\mu - k_j)^+\| \leq k_j D_1 D_2 \|\chi_{k_j}\|_{L^2(0,T;L^4(\Omega))}^{8/7}.\end{aligned} \quad (3.6)$$

Therefore, if we set

$$S_j := \|\chi_{k_j}\|_{L^2(0,T;L^4(\Omega))} \quad \text{for } j = 0, 1, \dots,$$

then we have

$$S_{j+1} \leq \frac{k_j}{k_{j+1} - k_j} D_1 D_2 S_j^{8/7} \leq 4 D_1 D_2 2^j S_j^{8/7} \quad \text{for } j = 0, 1, \dots$$

Using [12, Lemma 5.6, p. 95], we can conclude that $S_j \rightarrow 0$ as $j \rightarrow \infty$, provided that

$$S_0 = \|\chi_{k_0}\|_{L^2(0,T;L^4(\Omega))} \leq (4 D_1 D_2)^{-7} 2^{-49}. \quad (3.7)$$

Now recall that $\chi_{k_0} = \chi_M$ and, owing to (3.5), $M > \Phi_0$ and $m = M/\Phi_0$. Also,

$$\chi_M = 1 < \frac{\mu - \Phi_0}{M - \Phi_0} \quad \text{if } \mu > M, \quad \text{and } \chi_M = 0 \quad \text{otherwise.}$$

Therefore, using (3.3) and (3.4) with $k = k_0 = M$, we find that

$$\begin{aligned} S_0 &\leq \frac{1}{M - \Phi_0} \|(\mu - \Phi_0)^+\|_{L^2(0,T;L^4(\Omega))} \leq \frac{D_2}{M - \Phi_0} \|(\mu - \Phi_0)^+\| \\ &\leq \frac{D_1 D_2}{m - 1} \|\chi_{\Phi_0}\|_{L^2(0,T;L^4(\Omega))}^{8/7} \leq \frac{D_1 D_2}{m - 1} |\Omega|^{\frac{1}{4} \cdot \frac{8}{7}} T^{\frac{1}{2} \cdot \frac{8}{7}}. \end{aligned}$$

We are now in a position to choose $m := 1 + D_1 D_2 |\Omega|^{2/7} T^{4/7} (4 D_1 D_2)^7 2^{49}$. Then, $m > 1$ and (3.7) is satisfied. Consequently,

$$\|\chi_{2M}\|_{L^2(0,T;L^4(\Omega))} = \lim_{j \rightarrow \infty} S_j = 0,$$

due to Beppo Levi's Monotone Convergence Theorem. This implies that $\mu \leq 2M$ a.e. in Q , and the boundedness of μ is proved. \blacksquare

Now that the boundedness condition (3.1) is shown, we can prove the following uniqueness and stability result, which corresponds to Theorem 2.2 in [5].

Theorem 3.2

(i) *Suppose that (A1)–(A6) are fulfilled. Then the system (1.1)–(1.4) has a unique solution (ρ, μ) satisfying (2.3)–(2.6).*

(ii) *Suppose that (A1), (A2), (A4) and (A5) are fulfilled and that the functions u_1, u_2 satisfy the conditions (A3) and (A6). Moreover, let (ρ_i, μ_i) be the solutions to (1.1)–(1.4) corresponding to u_i , $i = 1, 2$, and $u := u_1 - u_2$, $\rho := \rho_1 - \rho_2$ and $\mu := \mu_1 - \mu_2$. Then we have, for every $t \in [0, T]$,*

$$\begin{aligned} \max_{0 \leq s \leq t} (\|\mu(s)\|_H^2 + \|\rho(s)\|_V^2) + \int_0^t \int_{\Omega} (\|\mu(s)\|_V^2 + \|\rho_t(s)\|_H^2 + \|\rho(s)\|_W^2) ds \\ \leq K_1^* \int_0^t \|u(s)\|_{L^2(\Gamma)}^2 ds, \end{aligned} \quad (3.8)$$

with a constant $K_1^* > 0$ that only depends on the data of the system.

Proof. Obviously, the assertion (i) follows directly from (ii). So we only need to show (ii). To this end, observe that by Theorem 3.1 there are constants $M > 0$ and $0 < r_* < r^* < 1$ such that $0 \leq \mu_i \leq M$ and $r_* \leq \rho_i \leq r^*$ a.e. in Q , for $i = 1, 2$. Moreover, the function $r \mapsto r - f'(r)$, $r_* \leq r \leq r^*$, has a Lipschitz constant $L > 0$. Next, we observe that the pair (ρ, μ) is a solution to the system

$$(\varepsilon + 2\rho_1)\mu_t + 2\rho\mu_{2,t} + \mu\rho_{1,t} + \mu_2\rho_t - \Delta\mu = 0 \quad \text{a.e. in } Q, \quad (3.9)$$

$$\delta\rho_t - \Delta\rho = \mu - (f'(\rho_1) - f'(\rho_2)) \quad \text{a.e. in } Q, \quad (3.10)$$

$$\frac{\partial\rho}{\partial\mathbf{n}} = 0, \quad \frac{\partial\mu}{\partial\mathbf{n}} = \alpha(u - \mu) \quad \text{a.e. on } \Sigma, \quad (3.11)$$

$$\rho(x, 0) = \mu(x, 0) = 0, \quad \text{for a.e. } x \in \Omega. \quad (3.12)$$

Now observe that $2\rho_1\mu\mu_t = (\rho_1\mu^2)_t - \mu^2\rho_{1,t}$. Hence, if we test (3.9) by μ then we obtain, using Young's inequality, that for every $t \in [0, T]$ it holds

$$\begin{aligned} & \int_{\Omega} \left(\frac{\varepsilon}{2} + \rho_1(t) \right) \mu^2(t) dx + \int_0^t \int_{\Omega} |\nabla \mu|^2 dx ds + \int_0^t \int_{\Gamma} \alpha |\mu|^2 d\sigma ds \\ & \leq C \int_0^t \int_{\Gamma} |u|^2 d\sigma ds + \int_0^t \int_{\Gamma} \frac{\alpha}{2} |\mu|^2 d\sigma ds \\ & \quad + \int_0^t \int_{\Omega} |\mu| (2|\rho| |\mu_{2,t}| + |\mu_2| |\rho_t|) dx ds. \end{aligned} \quad (3.13)$$

We have, owing to the continuity of the embedding $H^1(\Omega) \subset L^4(\Omega)$ and to Young's inequality,

$$\begin{aligned} & \int_0^t \int_{\Omega} 2|\mu| |\rho| |\mu_{2,t}| dx ds \leq C \int_0^t \|\mu_{2,t}(s)\|_H^2 \|\mu(s)\|_{L^4(\Omega)} \|\rho(s)\|_{L^4(\Omega)} ds \\ & \leq \gamma \int_0^t \|\mu(s)\|_V^2 ds + \frac{C}{\gamma} \int_0^t \|\mu_{2,t}(s)\|_H^2 \|\rho(s)\|_V^2 ds, \end{aligned} \quad (3.14)$$

where, owing to (2.4), the mapping $s \mapsto \|\mu_{2,t}(s)\|_H^2$ belongs to $L^1(0, T)$. Moreover, we also have $\mu_2 \in L^\infty(Q)$, and thus

$$\begin{aligned} & \int_0^t \int_{\Omega} |\mu| |\mu_2| |\rho_t| dx ds \leq C \int_0^t \|\rho_t(s)\|_H \|\mu(s)\|_H ds \\ & \leq \gamma \int_0^t \|\rho_t(s)\|_H^2 ds + \frac{C}{\gamma} \int_0^t \|\mu(s)\|_H^2 ds. \end{aligned} \quad (3.15)$$

Next, we add ρ on both sides of Eq. (3.10) and test the resulting equation by ρ_t . Invoking Young's inequality, it is easily seen that, for every $t \in [0, T]$,

$$\begin{aligned} & \delta \int_0^t \|\rho_t(s)\|_H^2 ds + \|\rho(t)\|_V^2 \\ & \leq \gamma \int_0^t \|\rho_t(s)\|_H^2 ds + \frac{C}{\gamma} \int_0^t (\|\mu(s)\|_H^2 + L^2 \|\rho(s)\|_H^2) ds. \end{aligned} \quad (3.16)$$

Now we can combine (3.13)–(3.16). Choosing $\gamma > 0$ sufficiently small, and applying Gronwall's lemma, we see that (3.8) is satisfied. \blacksquare

The stability estimate (3.8) can be improved if further regularity is assumed for f . The following result is a counterpart of Lemma 3.1 in [7]. We remark at this place that (2.3) implies, in particular, that ρ is weakly continuous as a mapping from $[0, T]$ into W , which justifies the formulation of the estimate (3.17) below.

Theorem 3.3 *Suppose that the assumptions of Theorem 3.2,(ii) are satisfied, and assume that*

$$(A7) \quad f \in C^3(0, 1).$$

Then we have, for every $t \in [0, T]$,

$$\begin{aligned}
& \max_{0 \leq s \leq t} (\|\mu(s)\|_V^2 + \|\rho_t(s)\|_V^2 + \|\rho(s)\|_W^2) \\
& + \int_0^t (\|\mu_t(s)\|_H^2 + \|\rho_t(s)\|_W^2) ds \\
& \leq K_2^* \left\{ \|u(0)\|_{L^2(\Gamma)}^2 + \int_0^t (\|u(s)\|_{L^2(\Gamma)}^2 + \|u_t(s)\|_{L^2(\Gamma)}^2) ds \right\} \quad (3.17)
\end{aligned}$$

with a constant $K_2^* > 0$ that only depends on the data of the system.

Remark 3.4 We note that $\|u(0)\|_{L^2(\Gamma)}^2 \leq I(t)/\max\{1, t\}$ where $I(t)$ denotes the last integral of (3.17). It follows that $\|u(0)\|_{L^2(\Gamma)}^2$ can be dropped if one pretends (3.17) just for $t = T$.

Proof of Theorem 3.3. We closely follow the lines of the proof of Lemma 3.1 in [7]. Since the proof given there carries over to our situation with minor changes, we can afford to be brief. First, observe that by Theorem 3.1 there are constants $M > 0$ and $0 < r_* < r^* < 1$ such that $0 \leq \mu_i \leq M$ and $r_* \leq \rho_i \leq r^*$ a. e. in Q , for $i = 1, 2$. Next, we recall that the pair (ρ, μ) is a solution to the system (3.9)–(3.12). We test Eq. (3.9) formally by μ_t . It then follows, with the use of Young's inequality, that

$$\begin{aligned}
& \varepsilon \int_0^t \|\mu_t(s)\|_H^2 ds + \frac{1}{2} \|\nabla \mu(t)\|_H^2 + \frac{1}{2} \int_\Gamma \alpha |\mu(t)|^2 d\sigma \\
& \leq \int_0^t \int_\Gamma \alpha u \mu_t d\sigma ds \\
& + \int_0^t \int_\Omega (2|\rho| |\mu_{2,t}| + |\mu| |\rho_{1,t}| + |\mu_2| |\rho_t|) |\mu_t| dx ds. \quad (3.18)
\end{aligned}$$

Now, by virtue of integration by parts with respect to t , and invoking (3.8), Young's inequality and the trace theorem,

$$\begin{aligned}
& \left| \int_0^t \int_\Gamma \alpha u \mu_t d\sigma ds \right| \\
& \leq \gamma \int_\Gamma \alpha |\mu(t)|^2 d\sigma + \frac{C}{\gamma} \int_\Gamma |u(t)|^2 d\sigma + \frac{C}{\gamma} \int_0^t \int_\Gamma |u_t| |\mu| d\sigma ds \\
& \leq C \gamma \|\mu(t)\|_V^2 + \frac{C}{\gamma} \int_\Gamma |u(0)|^2 d\sigma + \frac{C}{\gamma} \int_0^t \int_\Gamma (|u|^2 + |u_t|^2) d\sigma ds. \quad (3.19)
\end{aligned}$$

Employing almost exactly the same arguments as in the proof of Lemma 3.1 in [7] (the minor necessary changes are left as an easy exercise to the reader), and taking advantage of (3.8),

we conclude the estimate (where $\gamma > 0$ is arbitrary)

$$\begin{aligned}
& \int_0^t \int_{\Omega} (2|\rho| |\mu_{2,t}| + |\mu| |\rho_{1,t}| + |\mu_2| |\rho_t|) |\mu_t| dx ds \\
& \leq 3\gamma \int_0^t \|\mu_t(s)\|_H^2 ds + \frac{C}{\gamma} \int_0^t \|\rho_{1,t}(s)\|_V^2 \|\mu(s)\|_V^2 ds \\
& \quad + \frac{C}{\gamma} \int_0^t \|\mu_{2,t}(s)\|_H^2 \|\Delta\rho(s)\|_H^2 ds + \frac{C}{\gamma} \int_0^t \int_{\Gamma} |u|^2 d\sigma ds. \tag{3.20}
\end{aligned}$$

Next, we test (3.10) formally by $-\Delta\rho_t$. By the same token as in the proof of Lemma 3.1 in [7], we deduce for arbitrary $\gamma > 0$ the estimate

$$\begin{aligned}
& \delta \int_0^t \|\nabla\rho_t(s)\|_H^2 ds + \frac{1}{4} \|\Delta\rho(t)\|_H^2 \leq \gamma \int_0^t \|\mu_t(s)\|_H^2 ds \\
& \quad + \frac{C}{\gamma} \int_0^t (1 + \|\rho_{2,t}(s)\|_V^2) \|\Delta\rho(s)\|_H^2 ds + C \int_0^t \int_{\Gamma} |u|^2 d\sigma ds. \tag{3.21}
\end{aligned}$$

Now observe that, owing to Theorem 2.1, the mappings $s \mapsto \|\rho_{i,t}(s)\|_V^2$, $i = 1, 2$, and $s \mapsto \|\mu_{2,t}(s)\|_H^2$ all belong to $L^1(0, T)$. Hence, combining the estimates (3.18)–(3.21), adjusting $\gamma > 0$ sufficiently small, and invoking Gronwall's lemma, we can conclude that for every $t \in [0, T]$ it holds

$$\begin{aligned}
& \int_0^t (\|\nabla\rho_t(s)\|_H^2 + \|\mu_t(s)\|_H^2) ds + \max_{0 \leq s \leq t} (\|\mu(s)\|_V^2 + \|\rho(s)\|_W^2) \\
& \leq C \left\{ \|u(0)\|_{L^2(\Gamma)}^2 + \int_0^t \int_{\Gamma} (|u|^2 + |u_t|^2) d\sigma ds \right\}. \tag{3.22}
\end{aligned}$$

Next, we formally differentiate (3.10) with respect to t , and obtain

$$\delta\rho_{tt} - \Delta\rho_t = \mu_t - f''(\rho_1)\rho_t - (f''(\rho_1) - f''(\rho_2))\rho_{2,t}, \tag{3.23}$$

with zero initial and Neumann boundary conditions for ρ_t . Hence, testing (3.23) by ρ_t , invoking Young's inequality, and recalling (3.8) and (3.22), we find that

$$\begin{aligned}
& \frac{\delta}{2} \|\rho_t(t)\|_H^2 + \int_0^t \|\nabla\rho_t(s)\|_H^2 ds \\
& \leq C \left\{ \|u(0)\|_{L^2(\Gamma)}^2 + \int_0^t \int_{\Gamma} (|u|^2 + |u_t|^2) d\sigma ds \right\} \\
& \quad + \int_0^t \int_{\Omega} |\rho_{2,t}| |f''(\rho_1) - f''(\rho_2)| |\rho_t| dx ds. \tag{3.24}
\end{aligned}$$

Moreover, using Hölder's and Young's inequalities, (A7) and (3.8), we see that

$$\begin{aligned}
& \int_0^t \int_{\Omega} |\rho_{2,t}| |f''(\rho_1) - f''(\rho_2)| |\rho_t| dx ds \\
& \leq C \int_0^t \|\rho_{2,t}(s)\|_{L^4(\Omega)} \|\rho(s)\|_{L^4(\Omega)} \|\rho_t(s)\|_H ds \\
& \leq C \left(\int_0^t \|\rho_t(s)\|_H^2 ds + \max_{0 \leq s \leq t} \|\rho(s)\|_V^2 \int_0^t \|\rho_{2,t}(s)\|_V^2 ds \right) \\
& \leq C \int_0^t \int_{\Gamma} (|u|^2 + |u_t|^2) d\sigma ds. \tag{3.25}
\end{aligned}$$

Finally, we test (3.23) by $-\Delta\rho_t$. Using Young's inequality and (3.22), we find that

$$\begin{aligned}
& \frac{\delta}{2} \|\nabla\rho_t(t)\|_H^2 + \int_0^t \|\Delta\rho_t(s)\|_H^2 ds \\
& \leq \gamma \int_0^t \|\Delta\rho_t(s)\|_H^2 ds + \frac{C}{\gamma} \left\{ \|u(0)\|_{L^2(\Gamma)}^2 + \int_0^t \int_{\Gamma} (|u|^2 + |u_t|^2) d\sigma ds \right\} \\
& \quad + \int_0^t \int_{\Omega} |\rho_{2,t}| |f''(\rho_1) - f''(\rho_2)| |\Delta\rho_t| dx ds \\
& \leq 2\gamma \int_0^t \|\Delta\rho_t(s)\|_H^2 ds + \frac{C}{\gamma} \left\{ \|u(0)\|_{L^2(\Gamma)}^2 + \int_0^t \int_{\Gamma} (|u|^2 + |u_t|^2) d\sigma ds \right\} \\
& \quad + \frac{C}{\gamma} \max_{0 \leq s \leq t} \|\rho(s)\|_V^2 \int_0^t \|\rho_{2,t}(s)\|_V^2 ds \\
& \leq 2\gamma \int_0^t \|\Delta\rho_t(s)\|_H^2 ds \\
& \quad + \frac{C}{\gamma} \left\{ \|u(0)\|_{L^2(\Gamma)}^2 + \int_0^t \int_{\Gamma} (|u|^2 + |u_t|^2) d\sigma ds \right\}. \tag{3.26}
\end{aligned}$$

Choosing $\gamma > 0$ appropriately small, we can infer that the estimate (3.17) is in fact true. This concludes the proof. \blacksquare

4 An optimal boundary control problem

In this section, we consider the following optimal boundary control problem:

(CP) Minimize the cost functional

$$\begin{aligned}
J(u, \rho, \mu) &= \frac{1}{2} \int_{\Omega} |\rho(x, T) - \rho_T(x)|^2 dx + \frac{\beta_1}{2} \int_0^T \int_{\Gamma} |u(x, t)|^2 d\sigma dt \\
& \quad + \frac{\beta_2}{2} \int_0^T \int_{\Omega} |\mu(x, t) - \mu_T(x, t)|^2 dx dt \tag{4.1}
\end{aligned}$$

subject to the state system (1.1)–(1.4) and to the control constraints

$$u \in \mathcal{U}_{\text{ad}} := \left\{ v \in H^1(0, T; L^2(\Gamma)) \cap L^\infty(\Sigma) : U_1 \leq v \leq U_2 \text{ a. e. on } \Sigma, \right. \\ \left. \|v_t\|_{L^2(0, T; L^2(\Gamma))} \leq R \right\}. \quad (4.2)$$

In this connection, we require that the hypotheses (A1)–(A7) be satisfied. In addition, we postulate:

$$(A8) \quad R > 0, \beta_i \geq 0, i = 1, 2, \rho_T \in L^2(\Omega), \mu_T \in L^2(Q), U_1, U_2 \in L^\infty(\Sigma), \text{ and there are constants } 0 < u_* < u^* < +\infty \text{ such that}$$

$$u_* \leq U_1 \leq U_2 \leq u^* \text{ a. e. on } \Sigma. \quad (4.3)$$

In what follows, we denote

$$\mathcal{X} := H^1(0, T; L^2(\Gamma)) \cap L^\infty(\Sigma), \quad \|u\|_{\mathcal{X}} := \|u\|_{H^1(0, T; L^2(\Gamma))} + \|u\|_{L^\infty(\Sigma)},$$

where $\|\cdot\|_{H^1(0, T; L^2(\Gamma))}$ denotes the standard norm in $H^1(0, T; L^2(\Gamma))$. Obviously, \mathcal{U}_{ad} is a nonempty, bounded, closed and convex subset of \mathcal{X} , and \mathcal{U}_{ad} is contained in the open set $\mathcal{U} \subset \mathcal{X}$ given by

$$\mathcal{U} := \left\{ v \in \mathcal{X} : \frac{1}{2} u_* < \text{ess inf } v, \text{ess sup } v < \frac{3}{2} u^*, \|v_t\|_{L^2(0, T; L^2(\Gamma))} < R+1 \right\}.$$

By Theorem 3.3, the control-to-state mapping $u \mapsto S(u) := (\rho, \mu)$ is Lipschitz continuous as a mapping from the set $\mathcal{U} \subset \mathcal{X}$ into the space

$$(H^1(0, T; W) \cap C^1([0, T]; V)) \times (H^1(0, T; H) \cap C^0([0, T]; V)).$$

We may without loss of generality assume (by possibly taking a larger K_2^*) that (3.17) is valid on the whole set \mathcal{U} with the same constant $K_2^* > 0$. It also follows from Theorem 3.1 that there exist constants $\mu^* > 0$ and $0 < r_* < r^* < 1$ such that for every $u \in \mathcal{U}$ it holds

$$0 \leq \mu \leq \mu^* \quad \text{and} \quad 0 < r_* \leq \rho \leq r^* < 1 \quad \text{a. e. in } Q, \quad (4.4)$$

where $(\rho, \mu) = S(u)$. Moreover, a closer inspection of the proof of Theorem 2.1 reveals that there is a constant $K_3^* > 0$ such that we have, for any $u \in \mathcal{U}$,

$$\|\rho\|_{W^{1, \infty}(0, T; H) \cap H^1(0, T; V) \cap L^\infty(0, T; W)} \\ + \|\mu\|_{H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; H^{3/2}(\Omega)) \cap L^\infty(Q)} \leq K_3^*. \quad (4.5)$$

Remark 4.1 Thanks to (4.4) and to $f \in C^3(0, 1)$, it holds $f'(\rho) \in L^\infty(Q)$. Also, by the embedding $V \subset L^6(\Omega)$, we have $\mu \in C^0([0, T]; L^6(\Omega))$. Notice also that (2.3) implies, in particular, that ρ is continuous from $[0, T]$ to $H^s(\Omega)$ for all $s < 2$; thus, since $H^s(\Omega) \subset$

$C(\overline{\Omega})$ for $s > 3/2$, we also have $\rho \in C(\overline{Q})$. Therefore, possibly choosing a larger constant K_3^* , we may without loss of generality assume that

$$\|\rho\|_{C(\overline{Q})} + \|\mu\|_{C^0([0,T];L^6(\Omega))} + \|\rho_t\|_{L^2(0,T;L^6(\Omega))} \leq K_3^* \quad \forall u \in \mathcal{U}. \quad (4.6)$$

Remark 4.2 The mathematical literature on control problems for phase field systems is scarce and usually restricted to the so-called *Caginalp model* of phase transitions (see, e. g., [11], [9], [10], [18], and the references given there). More general, thermodynamically consistent phase field models were the subject of [13]. In [7], the present authors investigated a control problem for the system (1.1)–(1.4) with distributed controls. Since many of the arguments employed in [7] carry over to the boundary control considered here, we can afford to be sketchy in some of the proofs in the following exposition.

4.1 Existence

We begin our discussion of the control problem **(CP)** with the following existence result:

Theorem 4.3 *Suppose that the conditions (A1)–(A8) are satisfied. Then the optimal control problem **(CP)** has a solution $\bar{u} \in \mathcal{U}_{\text{ad}}$.*

Proof. Let $\{u_n\} \subset \mathcal{U}_{\text{ad}}$ be a minimizing sequence for **(CP)**, and let $\{(\rho_n, \mu_n)\}$ be the sequence of the associated solutions to (1.1)–(1.4). We then can infer from (4.5) the existence of a triple $(\bar{u}, \bar{\rho}, \bar{\mu})$ such that, for a suitable subsequence again indexed by n , we have

$$\begin{aligned} u_n &\rightharpoonup \bar{u} \quad \text{weakly star in } H^1(0, T; L^2(\Gamma)) \cap L^\infty(\Sigma), \\ \rho_n &\rightharpoonup \bar{\rho} \quad \text{weakly star in } W^{1,\infty}(0, T; H) \cap H^1(0, T; V) \cap L^\infty(0, T; W), \\ \mu_n &\rightharpoonup \bar{\mu} \quad \text{weakly star in } H^1(0, T; H) \cap L^\infty([0, T]; V) \cap L^2(0, T; H^{3/2}(\Omega)). \end{aligned}$$

Clearly, we have that $\bar{u} \in \mathcal{U}_{\text{ad}}$. Moreover, by virtue of the Aubin-Lions lemma (cf. [14, Thm. 5.1, p. 58]) and similar compactness results (cf. [17, Sect. 8, Cor. 4]), we also have the strong convergences

$$\begin{aligned} \rho_n &\rightarrow \bar{\rho} \quad \text{strongly in } C^0([0, T]; H^s(\Omega)) \quad \text{for all } s < 2, \\ \mu_n &\rightarrow \bar{\mu} \quad \text{strongly in } C^0([0, T]; H) \cap L^2(0, T; V). \end{aligned}$$

From this we infer, possibly selecting another subsequence again indexed by n , that $\rho_n \rightarrow \bar{\rho}$ pointwise a. e. (actually, uniformly) in Q . In particular, $r_* \leq \bar{\rho} \leq r^*$ a. e. in Q and, since $f \in C^2(0, 1)$, also $f'(\rho_n) \rightarrow f'(\bar{\rho})$ strongly in $L^2(Q)$. Now notice that the above convergences imply, in particular, that

$$\begin{aligned} \rho_n &\rightarrow \bar{\rho} \quad \text{strongly in } C^0([0, T]; L^6(\Omega)), \\ \partial_t \rho_n &\rightharpoonup \partial_t \bar{\rho} \quad \text{weakly in } L^2(0, T; L^4(\Omega)), \\ \mu_n &\rightarrow \bar{\mu} \quad \text{strongly in } L^2(0, T; L^4(\Omega)), \\ \partial_t \mu_n &\rightharpoonup \partial_t \bar{\mu} \quad \text{weakly in } L^2(Q). \end{aligned}$$

From this, it is easily verified that

$$\begin{aligned}\mu_n \partial_t \rho_n &\rightarrow \bar{\mu} \partial_t \bar{\rho} \quad \text{weakly in } L^1(0, T; H), \\ \rho_n \partial_t \mu_n &\rightarrow \bar{\rho} \partial_t \bar{\mu} \quad \text{weakly in } L^2(0, T; L^{3/2}(\Omega)).\end{aligned}$$

In summary, if we pass to the limit as $n \rightarrow \infty$ in the state equations (1.1)–(1.4) written for the triple (u_n, ρ_n, μ_n) , we find that $(\bar{\rho}, \bar{\mu}) = S(\bar{u})$, that is, the triple $(\bar{u}, \bar{\rho}, \bar{\mu})$ is admissible for the control problem **(CP)**. From the weak sequential lower semicontinuity of the cost functional J it finally follows that \bar{u} , together with $(\bar{\rho}, \bar{\mu}) = S(\bar{u})$, is a solution to **(CP)**. This concludes the proof. \blacksquare

Remark 4.4 It can be shown that this existence result holds for much more general cost functionals. All we need is that J enjoy appropriate weak sequential lower semicontinuity properties that match the above weak convergences.

Remark 4.5 Since the state component ρ is continuous on \bar{Q} , the existence result remains valid if suitable pointwise state constraints for ρ are added (provided the admissible set is not empty).

4.2 Necessary optimality conditions

In this section, we derive the first-order necessary conditions of optimality for problem **(CP)**. To this end, we first show that the control-to-state operator $S : u \mapsto (\rho, \mu)$ is Fréchet differentiable as a mapping from $\mathcal{U} \subset \mathcal{X}$ into the Banach space $(\mathcal{Y}, \|\cdot\|_{\mathcal{Y}})$, where

$$\begin{aligned}\mathcal{Y} &:= (H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; W)) \\ &\quad \times (C^0([0, T]; H) \cap L^2(0, T; V)).\end{aligned}$$

4.2.1 The linearized system

Let $\bar{u} \in \mathcal{U}$ and $h \in \mathcal{X}$ be given and $(\bar{\rho}, \bar{\mu}) = S(\bar{u})$. As a preparatory step, we consider the following system, which is obtained by linearizing the system (1.1)–(1.4) at $(\bar{\rho}, \bar{\mu})$:

$$(\varepsilon + 2\bar{\rho}) \eta_t - \Delta \eta + 2\bar{\mu}_t \xi + \bar{\mu} \xi_t + \bar{\rho}_t \eta = 0 \quad \text{a. e. in } Q, \quad (4.7)$$

$$\delta \xi_t - \Delta \xi = -f''(\bar{\rho}) \xi + \eta \quad \text{a. e. in } Q, \quad (4.8)$$

$$\frac{\partial \xi}{\partial \mathbf{n}} = 0, \quad \frac{\partial \eta}{\partial \mathbf{n}} = \alpha(h - \eta) \quad \text{a. e. on } \Sigma, \quad (4.9)$$

$$\xi(x, 0) = \eta(x, 0) = 0 \quad \text{for a. e. } x \in \Omega. \quad (4.10)$$

We expect for the Fréchet derivative $DS(\bar{u})$ at \bar{u} (if it exists) that $(\xi, \eta) = DS(\bar{u})h$, provided that (4.7)–(4.10) admits a unique solution (ξ, η) . In view of (2.3), (2.4), and (3.1), we can guess the regularity of ξ and η :

$$\xi \in H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; W) \cap L^\infty(Q), \quad (4.11)$$

$$\eta \in H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; H^{3/2}(\Omega)). \quad (4.12)$$

Notice that also in this case we cannot expect that $\eta(t) \in H^2(\Omega)$ a.e. in $(0, T)$ due to the low space regularity of h , and we could repeat Remark 2.2 here. Nevertheless, if (4.11) and (4.12) hold, then the collection of source terms in (4.7), i. e., the part $-2\bar{\mu}_t \xi - \bar{\mu} \xi_t - \bar{\rho}_t \eta$, belongs to $L^2(Q)$, whereas the regularity (4.12) for η allows us to conclude from (4.8) that also $\xi \in C(\bar{Q})$ (by applying maximal parabolic regularity theory, see, e. g., [8, Thm. 6.8] or [18, Lemma 7.12]).

In fact, ξ is even more regular: indeed, we may differentiate (4.8) with respect to t to find that

$$\delta \xi_{tt} - \Delta \xi_t = -f'''(\bar{\rho}) \bar{\rho}_t \xi - f''(\bar{\rho}) \xi_t + \eta_t, \quad (4.13)$$

with zero initial and Neumann boundary conditions for ξ_t . Since the right-hand side of (4.13) belongs to $L^2(Q)$, we may test by any of the functions ξ_t , ξ_{tt} , and $-\Delta \xi_t$, to obtain that even

$$\xi \in H^2(0, T; H) \cap C^1([0, T]; V) \cap H^1(0, T; W). \quad (4.14)$$

Notice, however, that this fact has no bearing on the regularity of η , since the coefficient $\bar{\mu}_t$ in (4.7) only belongs to $L^2(Q)$.

The following well-posedness result resembles Proposition 3.2 in [7].

Proposition 4.6 *Suppose that (A1)–(A8) are fulfilled. Then the system (4.7)–(4.10) has a unique solution (ξ, η) satisfying (4.12), (4.14), and*

$$\begin{aligned} & \|\xi\|_{H^2(0, T; H) \cap C^1([0, T]; V) \cap H^1(0, T; W)} + \|\eta\|_{H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; H^{3/2}(\Omega))} \\ & \leq K_4^* \|h\|_{H^1(0, T; L^2(\Gamma))}, \end{aligned} \quad (4.15)$$

with a constant $K_4^* > 0$ that is independent of the choice of $\bar{u} \in \mathcal{U}$ and $h \in \mathcal{X}$.

Remark 4.7 It follows from Proposition 4.6, in particular, that the linear mapping $h \mapsto (\xi, \eta)$ is continuous from \mathcal{X} into \mathcal{Y} .

Proof. We follow the lines of the proof of our previous existence results and proceed in a series of steps.

Step 1: Approximation. As in the proof of Theorem 2.1, we use an approximation technique based on a delay in the right-hand side of (4.8). To this end, for $\tau > 0$ we resume the definition of the translation operator $\mathcal{T}_\tau : L^1(0, T; H) \rightarrow L^1(0, T; H)$ by putting, for every $v \in L^1(0, T; H)$ and almost every $t \in (0, T)$,

$$(\mathcal{T}_\tau v)(t) = v(t - \tau) \text{ if } t \geq \tau, \quad \text{and} \quad (\mathcal{T}_\tau v)(t) = 0 \text{ if } t < \tau. \quad (4.16)$$

Notice that, for any $v \in L^2(Q)$ and any $\tau > 0$, we obviously have $\|\mathcal{T}_\tau v\|_{L^2(Q)} \leq \|v\|_{L^2(Q)}$. Then, for any fixed $\tau > 0$, we look for functions (ξ^τ, η^τ) , which satisfy (4.11) and (4.12) and the system:

$$(\varepsilon + 2\bar{\rho}) \eta_t^\tau - \Delta \eta^\tau + 2\bar{\mu}_t \xi^\tau + \bar{\mu} \xi_t^\tau + \bar{\rho}_t \eta^\tau = 0 \quad \text{a. e. in } Q, \quad (4.17)$$

$$\delta \xi_t^\tau - \Delta \xi^\tau + f''(\bar{\rho}) \xi^\tau = \mathcal{T}_\tau \eta^\tau \quad \text{a. e. in } Q, \quad (4.18)$$

$$\frac{\partial \xi^\tau}{\partial \mathbf{n}} = 0, \quad \frac{\partial \eta^\tau}{\partial \mathbf{n}} = \alpha(h - \eta^\tau) \quad \text{a. e. on } \Sigma, \quad (4.19)$$

$$\xi^\tau(x, 0) = \eta^\tau(x, 0) = 0 \quad \text{for a. e. } x \in \Omega. \quad (4.20)$$

Precisely, we choose for $\tau > 0$ the discrete values $\tau = T/N$, where $N \in \mathbb{N}$ is arbitrary, and put $t_n = n\tau$, $0 \leq n \leq N$, and $I_n = (0, t_n)$. For $1 \leq n \leq N$, we solve the problem

$$(\varepsilon + 2\bar{\rho})\eta_{n,t} - \Delta\eta_n + 2\bar{\mu}_t\xi_n + \bar{\mu}\xi_{n,t} + \bar{\rho}_t\eta_n = 0 \quad \text{a. e. in } \Omega \times I_n, \quad (4.21)$$

$$\frac{\partial\eta_n}{\partial\mathbf{n}} = \alpha(h - \eta_n) \quad \text{a. e. on } \Gamma \times I_n, \quad \eta_n(x, 0) = 0 \quad \text{for a. e. } x \in \Omega, \quad (4.22)$$

$$\delta\xi_{n,t} - \Delta\xi_n + f''(\bar{\rho})\xi_n = \mathcal{T}_\tau\eta_n \quad \text{a. e. in } \Omega \times I_n, \quad (4.23)$$

$$\frac{\partial\xi_n}{\partial\mathbf{n}} = 0 \quad \text{a. e. on } \Gamma \times I_n, \quad \xi_n(x, 0) = 0 \quad \text{for a. e. } x \in \Omega, \quad (4.24)$$

where the variables η_n and ξ_n , defined on I_n , have obvious meaning. Here, \mathcal{T}_τ acts on functions that are not defined on the entire interval $(0, T)$; however, for $n > 1$ it is still defined by (4.16), while for $n = 1$ we simply put $\mathcal{T}_\tau\eta_n = 0$. Notice that whenever the pairs (ξ_k, η_k) with

$$\xi_k \in H^1(I_k; H) \cap C^0(\bar{I}_k; V) \cap L^2(I_k; W) \cap C(\overline{\Omega \times I_k}), \quad (4.25)$$

$$\eta_k \in H^1(I_k; H) \cap C^0(\bar{I}_k; V) \cap L^2(I_k; H^{3/2}(\Omega)), \quad (4.26)$$

have been constructed for $1 \leq k \leq n < N$, then we look for the pair (ξ_{n+1}, η_{n+1}) that coincides with (ξ_n, η_n) in I_n , and note that the linear parabolic problem (4.23), (4.24) has a unique solution ξ_{n+1} on $\Omega \times I_{n+1}$ that satisfies (4.25) for $k = n + 1$. Inserting ξ_{n+1} in (4.21) (where n is replaced by $n + 1$), we then find that the linear parabolic problem (4.21), (4.22) admits a unique solution η_{n+1} that fulfills (4.26) for $k = n + 1$. Hence, we conclude that $(\xi^\tau, \eta^\tau) = (\xi_N, \eta_N)$ satisfies (4.17)–(4.20), and (4.11), (4.12).

Step 2: A priori estimates. We now prove a series of a priori estimates for the functions (ξ^τ, η^τ) . In the following, we denote by C_i ($i \in \mathbb{N}$) some generic positive constants, which may depend on $\varepsilon, \delta, \rho_*, \rho^*, \mu^*, T, K_1^*, K_2^*, K_3^*$, but not on τ (i. e., not on N). For the sake of simplicity, we omit the superscript τ and simply write (ξ, η) .

First a priori estimate. Observe that $2\bar{\rho}\eta\eta_t = (\bar{\rho}\eta^2)_t - \bar{\rho}_t\eta^2$. Hence, testing (4.17) by η , and invoking (4.19) and Young's inequality, we have, for $0 \leq t \leq T$,

$$\begin{aligned} & \int_\Omega \left(\frac{\varepsilon}{2} + \bar{\rho}(t) \right) \eta(t)^2 dx + \int_0^t \|\nabla\eta(s)\|_H^2 ds + \int_0^t \int_\Gamma \alpha \eta^2 d\sigma ds \\ & \leq \int_0^t \int_\Gamma \frac{\alpha}{2} \eta^2 d\sigma ds + C_1 \int_0^t \int_\Gamma h^2 d\sigma ds \\ & \quad + 2 \int_0^t \int_\Omega |\bar{\mu}_t| |\xi| |\eta| dx ds + \int_0^t \int_\Omega |\bar{\mu}| |\xi_t| |\eta| dx ds. \end{aligned} \quad (4.27)$$

For any $\gamma > 0$, we have, by Young's inequality and (4.4), that

$$\begin{aligned} & \int_0^t \int_\Omega |\bar{\mu}| |\xi_t| |\eta| dx ds \leq \|\bar{\mu}\|_{L^\infty(Q)} \int_0^t \|\eta(s)\|_H \|\xi_t(s)\|_H ds \\ & \leq \gamma \int_0^t \|\xi_t(s)\|_H^2 ds + \frac{C_2}{\gamma} \int_0^t \|\eta(s)\|_H^2 ds. \end{aligned} \quad (4.28)$$

Moreover,

$$\begin{aligned} \int_0^t \int_{\Omega} |\bar{\mu}_t| |\xi| |\eta| dx ds &\leq \int_0^t \|\bar{\mu}_t(s)\|_H \|\xi(s)\|_{L^4(\Omega)} \|\eta(s)\|_{L^4(\Omega)} ds \\ &\leq \gamma \int_0^t \|\eta(s)\|_V^2 ds + \frac{C_3}{\gamma} \int_0^t \|\bar{\mu}_t(s)\|_H^2 \|\xi(s)\|_V^2 ds. \end{aligned} \quad (4.29)$$

Notice that, by virtue of (4.5), the mapping $s \mapsto \|\bar{\mu}_t(s)\|_H^2$ is bounded by a function in $L^1(0, T)$.

Next, we add ξ on both sides of Eq. (4.18) and test the resulting equation by ξ_t . On using Young's inequality again, we obtain:

$$\begin{aligned} \frac{\delta}{4} \int_0^t \|\xi_t(s)\|_H^2 ds + \frac{1}{2} (\|\xi(t)\|_H^2 + \|\nabla \xi(t)\|_H^2) \\ \leq C_4 \left(\int_0^t \|\eta(s)\|_H^2 ds + \int_0^t \|\xi(s)\|_H^2 ds \right). \end{aligned} \quad (4.30)$$

Combining the inequalities (4.27)–(4.30), and choosing $\gamma > 0$ sufficiently small, we conclude from Gronwall's lemma that

$$\begin{aligned} \int_0^T (\|\xi_t(t)\|_H^2 + \|\eta(t)\|_V^2) dt + \max_{0 \leq t \leq T} (\|\xi(t)\|_V^2 + \|\eta(t)\|_H^2) \\ \leq C_5 \int_0^T \int_{\Gamma} |h|^2 d\sigma dt. \end{aligned} \quad (4.31)$$

Thanks to (4.19), we may also infer (possibly by choosing a larger C_5) that

$$\|\xi(t)\|_W^2 \leq C_5 \left(\|\Delta \xi(t)\|_H^2 + \int_0^T \int_{\Gamma} |h|^2 d\sigma dt \right) \quad \text{for all } t \in [0, T]. \quad (4.32)$$

Second a priori estimate. We test (4.17) by η_t and apply Young's inequality in order to obtain

$$\begin{aligned} \varepsilon \int_0^t \|\eta_t(s)\|_H^2 ds + \frac{1}{2} \|\nabla \eta(t)\|_H^2 + \int_{\Gamma} \frac{\alpha}{2} |\eta(t)|^2 d\sigma \leq \int_0^t \int_{\Gamma} \alpha h \eta_t d\sigma ds \\ + \int_0^t \int_{\Omega} (2|\bar{\mu}_t| |\xi| + |\bar{\mu}| |\xi_t| + |\bar{\rho}_t| |\eta|) |\eta_t| dx ds. \end{aligned} \quad (4.33)$$

By (4.4), we can infer from Young's inequality that

$$\int_0^t \int_{\Omega} |\bar{\mu}| |\xi_t| |\eta_t| dx ds \leq \gamma \int_0^t \|\eta_t(s)\|_H^2 ds + \frac{C_6}{\gamma} \int_0^t \|\xi_t(s)\|_H^2 ds. \quad (4.34)$$

Moreover, by virtue of Hölder's and Young's inequalities,

$$\begin{aligned} \int_0^t \int_{\Omega} |\bar{\rho}_t| |\eta| |\eta_t| dx ds \\ \leq \gamma \int_0^t \|\eta_t(s)\|_H^2 ds + \frac{C_7}{\gamma} \int_0^t \|\bar{\rho}_t(s)\|_{L^4(\Omega)}^2 \|\eta(s)\|_{L^4(\Omega)}^2 ds \\ \leq \gamma \int_0^t \|\eta_t(s)\|_H^2 ds + \frac{C_8}{\gamma} \int_0^t \|\bar{\rho}_t(s)\|_V^2 \|\eta(s)\|_V^2 ds. \end{aligned} \quad (4.35)$$

Observe that by (4.5) the mapping $s \mapsto \|\bar{\rho}_t(s)\|_V^2$ is bounded by a function in $L^1(0, T)$. Also, we have, owing to the continuity of the embedding $W \subset L^\infty(\Omega)$ and (4.32),

$$\begin{aligned}
& \int_0^t \int_\Omega 2 |\bar{\mu}_t| |\xi| |\eta_t| dx ds \\
& \leq \gamma \int_0^t \|\eta_t(s)\|_H^2 ds + \frac{C_9}{\gamma} \int_0^t \|\bar{\mu}_t(s)\|_H^2 \|\xi(s)\|_{L^\infty(\Omega)}^2 ds \\
& \leq \gamma \int_0^t \|\eta_t(s)\|_H^2 ds + \frac{C_{10}}{\gamma} \left(\int_0^T \int_\Gamma |h|^2 d\sigma dt \right. \\
& \qquad \qquad \qquad \left. + \int_0^t \|\bar{\mu}_t(s)\|_H^2 \|\Delta\xi(s)\|_H^2 ds \right), \tag{4.36}
\end{aligned}$$

where, owing to (4.5), the mapping $s \mapsto \|\bar{\mu}_t(s)\|_H^2$ is bounded by a function in $L^1(0, T)$.

Finally, we employ integration by parts, Young's inequality, (4.31), and the trace theorem to obtain

$$\begin{aligned}
& \left| \int_0^t \int_\Gamma \alpha h \eta_t d\sigma ds \right| \leq \int_\Gamma \alpha |h(t)| |\eta(t)| d\sigma + \int_0^t \int_\Gamma \alpha |h_t| |\eta| d\sigma ds \\
& \leq \int_\Gamma \frac{\alpha}{4} |\eta(t)|^2 d\sigma + C_{11} \|h\|_{H^1(0, T; L^2(\Gamma))}^2. \tag{4.37}
\end{aligned}$$

Next, we formally test (4.18) by $-\Delta\xi_t$ to obtain, for every $t \in [0, T]$,

$$\delta \int_0^t \|\nabla\xi_t(s)\|_H^2 ds + \frac{1}{2} \|\Delta\xi(t)\|_H^2 = \int_0^t \int_\Omega (-\mathcal{T}_\tau\eta) - f''(\bar{\rho}) \xi \Delta\xi_t dx ds. \tag{4.38}$$

Now, by virtue of (4.31) and invoking Young's inequality, we have

$$\begin{aligned}
& \left| \int_0^t \int_\Omega (\mathcal{T}_\tau\eta) \Delta\xi_t dx ds \right| \\
& \leq \int_\Omega |(\mathcal{T}_\tau\eta)(t)| |\Delta\xi(t)| dx + \int_0^t \int_\Omega |\partial_t(\mathcal{T}_\tau\eta)| |\Delta\xi| dx ds \\
& \leq \frac{1}{8} \|\Delta\xi(t)\|_H^2 + C_{12} \int_0^T \int_\Gamma |h|^2 d\sigma dt \\
& \quad + \gamma \int_0^t \|\eta_t(s)\|_H^2 ds + \frac{1}{4\gamma} \int_0^t \|\Delta\xi(s)\|_H^2 ds. \tag{4.39}
\end{aligned}$$

Moreover, it turns out that

$$\begin{aligned}
& \left| \int_0^t \int_\Omega f''(\bar{\rho}) \xi \Delta\xi_t dx ds \right| \leq \int_\Omega |f''(\bar{\rho}(t))| |\xi(t)| |\Delta\xi(t)| dx \\
& \quad + \int_0^t \int_\Omega |f'''(\bar{\rho}) \bar{\rho}_t \xi + f''(\bar{\rho}) \xi_t| |\Delta\xi| dx ds. \tag{4.40}
\end{aligned}$$

We have, owing to (4.4) and (4.31),

$$\int_{\Omega} |f''(\bar{\rho}(t))| |\xi(t)| |\Delta\xi(t)| dx \leq \frac{1}{8} \|\Delta\xi(t)\|_H^2 + C_{13} \int_0^T \int_{\Gamma} |h|^2 d\sigma dt. \quad (4.41)$$

Also the second integral on the right-hand side of (4.40) is bounded, since (4.4), (4.5), and (4.31) imply that

$$\begin{aligned} & \int_0^t \int_{\Omega} |f'''(\bar{\rho}) \bar{\rho}_t \xi + f''(\bar{\rho}) \xi_t| |\Delta\xi| dx ds \\ & \leq C_{14} \int_0^t (\|\bar{\rho}_t(s)\|_{L^4(\Omega)}^2 \|\xi(s)\|_{L^4(\Omega)}^2 + \|\xi_t(s)\|_H^2) ds + \int_0^t \|\Delta\xi(s)\|_H^2 ds \\ & \leq C_{15} \left(\max_{0 \leq t \leq T} \|\xi(t)\|_V^2 \int_0^t \|\bar{\rho}_t(s)\|_V^2 ds + \int_0^t \|\xi_t(s)\|_H^2 ds \right) + \int_0^t \|\Delta\xi(s)\|_H^2 ds \\ & \leq C_{16} \int_0^T \int_{\Gamma} |h|^2 d\sigma dt + \int_0^t \|\Delta\xi(s)\|_H^2 ds, \end{aligned} \quad (4.42)$$

thanks to the continuity of the embedding $V \subset L^4(\Omega)$. Thus, combining the estimates (4.33)–(4.42), choosing $\gamma > 0$ sufficiently small, and invoking Gronwall's inequality, we can infer that

$$\begin{aligned} & \int_0^T \left(\|\eta_t(t)\|_H^2 + \|\xi_t(t)\|_V^2 \right) dt + \max_{0 \leq t \leq T} (\|\eta(t)\|_V^2 + \|\xi(t)\|_W^2) \\ & \leq C_{17} \|h\|_{H^1(0,T;L^2(\Gamma))}^2. \end{aligned} \quad (4.43)$$

Next, we compare terms in (4.17) and, arguing as in the derivation of (4.33)–(4.37), we readily find that

$$\int_0^T \|\Delta\eta(t)\|_H^2 dt \leq C_{18} \|h\|_{H^1(0,T;L^2(\Gamma))}^2.$$

Thus, by owing to elliptic regularity (cf. (4.19) and Remark 2.2), we conclude that

$$\int_0^T \|\eta(t)\|_{H^{3/2}(\Omega)}^2 dt \leq C_{19} \|h\|_{H^1(0,T;L^2(\Gamma))}^2. \quad (4.44)$$

Finally, we differentiate Eq. (4.18) with respect to t . We obtain:

$$\delta \xi_{tt} - \Delta \xi_t = \partial_t(\mathcal{T}_\tau \eta) - f'''(\bar{\rho}) \bar{\rho}_t \xi - f''(\bar{\rho}) \xi_t \quad \text{a. e. in } Q. \quad (4.45)$$

From (4.4)–(4.6), (4.43) and (4.44), we can infer that we may test (4.45) by any of the functions ξ_t , $-\Delta\xi_t$, and ξ_{tt} , in order to find that

$$\int_0^T \left(\|\xi_{tt}(t)\|_H^2 + \|\Delta\xi_t(t)\|_H^2 \right) dt + \max_{0 \leq t \leq T} \|\xi_t(t)\|_V^2 \leq C_{20} \|h\|_{H^1(0,T;L^2(\Gamma))}^2. \quad (4.46)$$

Step 3: Passage to the limit. Let (ξ^τ, η^τ) denote the solution to the system (4.17)–(4.20) associated with $\tau = T/N$, for $N \in \mathbb{N}$. In Step 2, we have shown that there is some $C > 0$, which does not depend on τ , such that

$$\begin{aligned} & \|\xi^\tau\|_{H^2(0,T;H) \cap C^1([0,T];V) \cap H^1(0,T;W) \cap C(\bar{Q})} \\ & + \|\eta^\tau\|_{H^1(0,T;H) \cap C^0([0,T];V) \cap L^2(0,T;H^{3/2}(\Omega))} \leq C \|h\|_{H^1(0,T;L^2(\Gamma))}. \end{aligned} \quad (4.47)$$

Hence, there is a subsequence $\tau_k \searrow 0$ such that

$$\begin{aligned}\xi^{\tau_k} &\rightharpoonup \xi \text{ weakly star in } H^2(0, T; H) \cap W^{1, \infty}(0, T; V) \cap H^1(0, T; W), \\ \eta^{\tau_k} &\rightharpoonup \eta \text{ weakly star in } H^1(0, T; H) \cap L^\infty(0, T; V) \cap L^2(0, T; H^{3/2}(\Omega)).\end{aligned}\tag{4.48}$$

From the trace theorem we can infer that ξ satisfies the boundary condition given in (4.9), while the boundary condition for η will be satisfied (either in the variational sense or in the sense of the appropriate trace theorem, see Remark 2.2) once we prove that we can pass to the limit in the products of (4.7), as shown below. Moreover, it is easily seen that also (4.10) is fulfilled. By compact embedding, we also have, in particular,

$$\xi^{\tau_k} \rightarrow \xi \text{ strongly in } C(\overline{Q}), \quad \eta^{\tau_k} \rightarrow \eta \text{ strongly in } L^2(Q),\tag{4.49}$$

so that $\bar{\rho} \eta_t^{\tau_k} \rightarrow \bar{\rho} \eta_t$ and $\bar{\mu} \xi_t^{\tau_k} \rightarrow \bar{\mu} \xi_t$, both weakly in $L^2(Q)$, $f''(\bar{\rho}) \xi^{\tau_k} \rightarrow f''(\bar{\rho}) \xi$ strongly in $L^2(Q)$, as well as $\bar{\mu}_t \xi_t^{\tau_k} \rightarrow \bar{\mu}_t \xi_t$ and $\bar{\rho}_t \eta^{\tau_k} \rightarrow \bar{\rho}_t \eta$, both strongly in $L^1(Q)$. Finally, it is easily verified that $\{\mathcal{T}_{\tau_k} \eta^{\tau_k}\}$ converges strongly in $L^2(Q)$ to η . In conclusion, we may pass to the limit as $k \rightarrow \infty$ in the system (4.17)–(4.20) (written for τ_k) to find that the pair (ξ, η) is in fact a solution to the linearized system (4.7)–(4.10).

We now show the uniqueness. If (ξ_1, η_1) , (ξ_2, η_2) are two solutions having the above properties, then the pair (ξ, η) , where $\xi = \xi_1 - \xi_2$ and $\eta = \eta_1 - \eta_2$, satisfies (4.7)–(4.10) with $h = 0$. We thus may repeat the first a priori estimate in Step 2 to conclude that $\xi = \eta = 0$.

Finally, taking the limit as $\tau \searrow 0$ in (4.47) and invoking the lower semicontinuity of norms, we obtain the inequality (4.15). This concludes the proof. \blacksquare

4.2.2 Fréchet differentiability of the control-to-state mapping

In this section, we prove the following result.

Proposition 4.8 *Suppose that the assumptions (A1)–(A8) are satisfied. Then the solution operator S , viewed as a mapping from \mathcal{X} to \mathcal{Y} , is Fréchet differentiable on \mathcal{U} . For any $\bar{u} \in \mathcal{U}$ the Fréchet derivative $DS(\bar{u})$ is for $h \in \mathcal{X}$ given by $DS(\bar{u})h = (\xi, \eta)$, where (ξ, η) is the unique solution to the linearized system (4.7)–(4.10).*

Proof. Let $\bar{u} \in \mathcal{U}$ be given and $(\bar{\rho}, \bar{\mu}) = S(\bar{u})$. Since \mathcal{U} is an open subset of \mathcal{X} , there is some $\lambda > 0$ such that $\bar{u} + h \in \mathcal{U}$ whenever $h \in \mathcal{X}$ satisfies $\|h\|_{\mathcal{X}} \leq \lambda$. In the following, we consider such perturbations $h \in \mathcal{X}$, and we define $(\rho^h, \mu^h) := S(\bar{u} + h)$ and put

$$z^h := \mu^h - \bar{\mu} - \eta^h, \quad y^h := \rho^h - \bar{\rho} - \xi^h,\tag{4.50}$$

where (ξ^h, η^h) denotes the unique solution to the linearized system (4.7)–(4.10) associated with h . Since the linear mapping $h \mapsto (\xi^h, \eta^h)$ is by Proposition 4.6 continuous from \mathcal{X} into \mathcal{Y} , it obviously suffices to show that there is an increasing function $g : [0, \lambda] \rightarrow [0, +\infty)$ which satisfies $\lim_{r \searrow 0} g(r)/r^2 = 0$ and

$$\begin{aligned}&\|y^h\|_{H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; W)}^2 + \|z^h\|_{C^0([0, T]; H) \cap L^2(0, T; V)}^2 \\ &\leq g(\|h\|_{H^1(0, T; L^2(\Gamma))}).\end{aligned}\tag{4.51}$$

Using the state system (1.2)–(1.4) and the linearized system (4.7)–(4.10), we easily verify that for $h \in \mathcal{X}$ with $\|h\|_{\mathcal{X}} \leq \lambda$ the pair (y^h, z^h) is a strong solution to the system

$$\begin{aligned} & (\varepsilon + 2\bar{\rho}) z_t^h + \bar{\rho}_t z^h + \bar{\mu} y_t^h + 2\bar{\mu}_t y^h - \Delta z^h \\ & = -2(\mu_t^h - \bar{\mu}_t)(\rho^h - \bar{\rho}) - (\rho_t^h - \bar{\rho}_t)(\mu^h - \bar{\mu}) \quad \text{a. e. in } Q, \end{aligned} \quad (4.52)$$

$$\delta y_t^h - \Delta y^h + f'(\rho^h) - f'(\bar{\rho}) - f''(\bar{\rho}) \xi^h = z^h, \quad \text{a. e. in } Q, \quad (4.53)$$

$$\frac{\partial y^h}{\partial \mathbf{n}} = 0, \quad \frac{\partial z^h}{\partial \mathbf{n}} = -\alpha z^h, \quad \text{a. e. on } \Sigma, \quad (4.54)$$

$$y^h(x, 0) = z^h(x, 0) = 0 \quad \text{for a. e. } x \in \Omega. \quad (4.55)$$

Notice that

$$\begin{aligned} y^h & \in H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; W) \cap C(\bar{Q}), \\ z^h & \in H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; H^{3/2}(\Omega)). \end{aligned}$$

For the sake of a better readability, in the following estimates we omit the superscript h of y^h and z^h . Also, we denote by C_i ($i \in \mathbb{N}$) certain positive constants that only depend on $\varepsilon, \delta, \rho_*, \rho^*, \mu^*, T, K_1^*, K_2^*, K_3^*, K_4^*$, but not on h .

We now add y on both sides of Eq. (4.53) and test the resulting equation by y_t . Using Young's inequality, we find that for all $t \in [0, T]$ it holds

$$\begin{aligned} & \frac{\delta}{2} \int_0^t \|y_t(s)\|_H^2 ds + \frac{1}{2} (\|\nabla y(t)\|_H^2 + \|y(t)\|_H^2) \leq \frac{2}{\delta} \int_0^t \|z(s)\|_H^2 ds \\ & + C_1 \int_0^t \|y(s)\|_H^2 ds + C_2 \int_0^t \|(f'(\rho^h) - f'(\bar{\rho}) - f''(\bar{\rho}) \xi^h)(s)\|_H^2 ds. \end{aligned} \quad (4.56)$$

In order to handle the third term on the right-hand side of (4.56), we note that the stability estimate (3.17) implies, in particular, that

$$\|\rho^h - \bar{\rho}\|_{L^\infty(Q)}^2 \leq K_2^* \|h\|_{H^1(0, T; L^2(\Gamma))}^2, \quad (4.57)$$

that is, $\rho^h \rightarrow \bar{\rho}$ uniformly on \bar{Q} as $\|h\|_{H^1(0, T; L^2(\Gamma))} \rightarrow 0$. Since $f \in C^3(0, 1)$, we can infer from Taylor's theorem and (4.4) that

$$|f'(\rho^h) - f'(\bar{\rho}) - f''(\bar{\rho}) \xi^h| \leq \max_{r_* \leq \sigma \leq r^*} \frac{|f'''(\sigma)|}{2} |\rho^h - \bar{\rho}|^2 + |f''(\bar{\rho})| |y| \quad \text{on } \bar{Q}. \quad (4.58)$$

It then follows from the estimates (3.17) and (4.56)–(4.57) that

$$\begin{aligned} & \frac{\delta}{2} \int_0^t \|y_t(s)\|_H^2 ds + \frac{1}{2} \|y(t)\|_V^2 \leq \frac{2}{\delta} \int_0^t \|z(s)\|_H^2 ds + C_3 \int_0^t \|y(s)\|_H^2 ds \\ & + C_4 \|h\|_{H^1(0, T; L^2(\Gamma))}^4. \end{aligned} \quad (4.59)$$

Next, observe that $2\bar{\rho} z z_t = (\bar{\rho} z^2)_t - \bar{\rho}_t z^2$. Therefore, testing (4.52) by z yields for every $t \in [0, T]$ that

$$\begin{aligned} & \int_{\Omega} \left(\frac{\varepsilon}{2} + \bar{\rho}(t) \right) z^2(t) dx + \int_0^t \|\nabla z(s)\|_H^2 ds + \int_0^t \int_{\Gamma} \alpha |z|^2 d\sigma dt \\ &= - \int_0^t \int_{\Omega} (\bar{\mu} y_t + 2\bar{\mu}_t y) z dx ds - 2 \int_0^t \int_{\Omega} (\mu_t^h - \bar{\mu}_t) (\rho^h - \bar{\rho}) z dx ds \\ & \quad - \int_0^t \int_{\Omega} (\rho_t^h - \bar{\rho}_t) (\mu^h - \bar{\mu}) z dx ds. \end{aligned} \quad (4.60)$$

We estimate the terms on the right-hand side of (4.60) individually. At first, using (4.4) and Young's inequality, we find that

$$\int_0^t \int_{\Omega} |\bar{\mu}| |y_t| |z| dx ds \leq \gamma \int_0^t \|y_t(s)\|_H^2 ds + \frac{C_5}{\gamma} \int_0^t \|z(s)\|_H^2 ds. \quad (4.61)$$

Moreover, using the continuity of the embedding $H^1(\Omega) \subset L^4(\Omega)$, as well as Hölder's and Young's inequalities, we have

$$\begin{aligned} 2 \int_0^t \int_{\Omega} |\bar{\mu}_t| |y| |z| dx ds &\leq 2 \int_0^t \|\bar{\mu}_t(s)\|_H \|z(s)\|_{L^4(\Omega)} \|y(s)\|_{L^4(\Omega)} ds \\ &\leq \gamma \int_0^t \|z(s)\|_V^2 ds + \frac{C_6}{\gamma} \int_0^t \|\bar{\mu}_t(s)\|_H^2 \|y(s)\|_V^2 ds. \end{aligned} \quad (4.62)$$

Observe that by (2.4) the mapping $s \mapsto \|\bar{\mu}_t(s)\|_H^2$ belongs to $L^1(0, T)$.

At this point, we can conclude from (3.17) and (4.57), invoking Young's inequality, that

$$\begin{aligned} & \int_0^t \int_{\Omega} 2 |\mu_t^h - \bar{\mu}_t| |\rho^h - \bar{\rho}| |z| dx ds \\ &\leq 2 \int_0^t \|(\mu_t^h - \bar{\mu}_t)(s)\|_H \|(\rho^h - \bar{\rho})(s)\|_{L^\infty(\Omega)} \|z(s)\|_H ds \\ &\leq C_7 \|\rho^h - \bar{\rho}\|_{L^\infty(Q)}^2 \int_0^t \|(\mu_t^h - \bar{\mu}_t)(s)\|_H^2 ds + \int_0^t \|z(s)\|_H^2 ds \\ &\leq \int_0^t \|z(s)\|_H^2 ds + C_8 \|h\|_{H^1(0,T;L^2(\Gamma))}^4. \end{aligned} \quad (4.63)$$

Finally, we invoke (3.17) and Hölder's and Young's inequalities, as well as the continuity of the embedding $H^1(\Omega) \subset L^4(\Omega)$, to obtain that

$$\begin{aligned} & \int_0^t \int_{\Omega} |\rho_t^h - \bar{\rho}_t| |\mu^h - \bar{\mu}| |z| dx ds \\ &\leq \max_{0 \leq s \leq t} \|z(s)\|_H \int_0^t \|(\rho_t^h - \bar{\rho}_t)(s)\|_{L^4(\Omega)} \|(\mu^h - \bar{\mu})(s)\|_{L^4(\Omega)} ds \\ &\leq \gamma \max_{0 \leq s \leq t} \|z(s)\|_H^2 + \frac{C_9}{\gamma} \int_0^t \|(\rho_t^h - \bar{\rho}_t)(s)\|_V^2 ds \int_0^t \|(\mu^h - \bar{\mu})(s)\|_V^2 ds \\ &\leq \gamma \max_{0 \leq s \leq t} \|z(s)\|_H^2 + C_{10} \|h\|_{H^1(0,T;L^2(\Gamma))}^4. \end{aligned} \quad (4.64)$$

Combining the estimates (4.59)–(4.64), taking the maximum with respect to $t \in [0, T]$, adjusting $\gamma > 0$ appropriately small, and invoking Gronwall's lemma, we arrive at the conclusion that $(y^h, z^h) = (y, z)$ satisfies the inequality

$$\begin{aligned} \|y^h\|_{H^1(0,T;H) \cap C^0([0,T];V)}^2 + \|z^h\|_{C^0([0,T];H) \cap L^2([0,T];V)}^2 \\ \leq C_{11} \|h\|_{H^1(0,T;L^2(\Gamma))}^4. \end{aligned} \quad (4.65)$$

Finally, testing (4.53) by $-\Delta y^h$, and using (4.58), we find that also

$$\|y^h\|_{L^2(0,T;W)}^2 \leq C_{12} \|h\|_{H^1(0,T;L^2(\Gamma))}^4. \quad (4.66)$$

Therefore, the function $g(r) := (C_{11} + C_{12}) r^4$ has the requested properties. This concludes the proof of the assertion. \blacksquare

Corollary 4.9 *Let the assumptions (A1)–(A8) be fulfilled, and let $\bar{u} \in \mathcal{U}_{\text{ad}}$ be an optimal control for the problem (CP) with associated state $(\bar{\rho}, \bar{\mu}) = S(\bar{u})$. Then, for every $v \in \mathcal{U}_{\text{ad}}$,*

$$\int_0^T \int_{\Gamma} \beta_1 \bar{u}(v - \bar{u}) \, d\sigma \, dt + \int_{\Omega} (\bar{\rho}(T) - \rho_T) \xi(T) \, dx + \int_0^T \int_{\Omega} \beta_2 (\bar{\mu} - \mu_T) \eta \, dx \, dt \geq 0, \quad (4.67)$$

where (ξ, η) is the unique solution to the linearized system (4.7)–(4.10) associated with $h = v - \bar{u}$.

Proof. Let $v \in \mathcal{U}_{\text{ad}}$ be arbitrary and $h = v - \bar{u}$. Then $\bar{u} + \lambda h \in \mathcal{U}_{\text{ad}}$ for $0 < \lambda \leq 1$. For any such λ , we have

$$\begin{aligned} 0 &\leq \frac{J(\bar{u} + \lambda h, S(\bar{u} + \lambda h)) - J(\bar{u}, S(\bar{u}))}{\lambda} \\ &\leq \frac{J(\bar{u} + \lambda h, S(\bar{u} + \lambda h)) - J(\bar{u}, S(\bar{u} + \lambda h))}{\lambda} \\ &\quad + \frac{J(\bar{u}, S(\bar{u} + \lambda h)) - J(\bar{u}, S(\bar{u}))}{\lambda}. \end{aligned}$$

It follows immediately from the definition of the cost functional J that the first summand on the right-hand side of this inequality converges to $\int_0^T \int_{\Gamma} \beta_1 \bar{u}(v - \bar{u}) \, d\sigma \, dt$ as $\lambda \searrow 0$. For the second summand, we obtain from Proposition 4.8 that

$$\begin{aligned} \lim_{\lambda \searrow 0} \frac{J(\bar{u}, S(\bar{u} + \lambda h)) - J(\bar{u}, S(\bar{u}))}{\lambda} \\ = \int_{\Omega} (\bar{\rho}(x, T) - \rho_T(x)) \xi(x, T) \, dx + \int_0^T \int_{\Omega} \beta_2 (\bar{\mu} - \mu_T) \eta \, dx \, dt, \end{aligned}$$

whence the assertion follows. \blacksquare

4.2.3 The optimality system

Let $\bar{u} \in \mathcal{U}_{\text{ad}}$ be an optimal control for **(CP)** with associated state $(\bar{\rho}, \bar{\mu}) = S(\bar{u})$. Then, for every $v \in \mathcal{U}_{\text{ad}}$, (4.67) holds. We now aim to eliminate (ξ, η) by introducing the adjoint state variables. To this end, we consider the *adjoint system*:

$$-(\varepsilon + 2\bar{\rho}) q_t - \bar{\rho}_t q - \Delta q = p + \beta_2 (\bar{\mu} - \mu_T) \quad \text{a. e. in } Q, \quad (4.68)$$

$$\frac{\partial q}{\partial \mathbf{n}} = -\alpha q \quad \text{a. e. in } \Sigma, \quad q(x, T) = 0 \quad \text{for a. e. } x \in \Omega, \quad (4.69)$$

$$-\delta p_t - \Delta p + f''(\bar{\rho}) p = \bar{\mu} q_t - \bar{\mu}_t q \quad \text{in } Q, \quad (4.70)$$

$$\frac{\partial p}{\partial \mathbf{n}} = 0 \quad \text{on } \Sigma, \quad \delta p(T) = \bar{\rho}(T) - \rho_T \quad \text{in } \Omega, \quad (4.71)$$

which is a linear backward-in-time parabolic system for the adjoint state variables p and q .

It must be expected that the adjoint state variables (p, q) be less regular than the state variables $(\bar{\rho}, \bar{\mu})$. Indeed, we only have $p(T) \in L^2(\Omega)$, and thus (4.70) and (4.71) should be interpreted in the usual weak sense. That is, we look for a vector-valued function $p \in H^1(0, T; V^*) \cap C^0([0, T]; H) \cap L^2(0, T; V)$ that, in addition to the final time condition (4.71), satisfies

$$\begin{aligned} & \langle -\delta p_t(t), v \rangle_{V^*, V} + \int_{\Omega} \nabla p(t) \cdot \nabla v \, dx + \int_{\Omega} f''(\bar{\rho}(t)) p(t) v \, dx \\ &= \int_{\Omega} (\bar{\mu}(t) q_t(t) - \bar{\mu}_t(t) q(t)) v \, dx, \end{aligned} \quad (4.72)$$

for every $v \in V$ and almost every $t \in (0, T)$. Notice that if $q \in H^1(0, T; H) \cap C^0([0, T]; V)$, then it is easily seen that $\bar{\mu} q_t - \bar{\mu}_t q \in L^{3/2}(Q)$, so that the integral on the right-hand side of (4.72) makes sense. On the other hand, if p has the expected regularity then the solution to (4.68), (4.69) should belong to $H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; H^2(\Omega))$.

The following result is an analogue of Theorem 3.7 in [7].

Theorem 4.10 *Suppose that $\bar{u} \in \mathcal{U}_{\text{ad}}$ is an optimal control for **(CP)** with associated state $(\bar{\rho}, \bar{\mu}) = S(\bar{u})$. Then the adjoint system (4.68)–(4.71) has a unique weak solution (p, q) with $p \in H^1(0, T; V^*) \cap C^0([0, T]; H) \cap L^2(0, T; V)$, $q \in H^1(0, T; H) \cap C^0([0, T]; V) \cap L^2(0, T; H^2(\Omega))$; moreover, for any $v \in \mathcal{U}_{\text{ad}}$, we have the inequality*

$$\int_0^T \int_{\Gamma} \beta_1 \bar{u} (v - \bar{u}) \, d\sigma \, dt + \int_0^T \int_{\Gamma} \alpha q (v - \bar{u}) \, d\sigma \, dt \geq 0. \quad (4.73)$$

Proof. The existence and uniqueness result for the adjoint state variables p and q follows using the same line of arguments as in the proof of Proposition 3.6 in [7], with only minor and straightforward changes that are due to the different boundary condition for q . Now let $v \in \mathcal{U}_{\text{ad}}$ be given. A standard calculation (which can be left as an easy exercise to the reader), using

the linearized system (4.7)–(4.10) with $h = v - \bar{u}$, repeated integration by parts, and the well-known integration by parts formula

$$\int_0^T (\langle v_t(t), w(t) \rangle_{V^*, V} + \langle w_t(t), v(t) \rangle_{V^*, V}) dt = \int_{\Omega} (v(T)w(T) - v(0)w(0)) dx$$

(which holds for all functions $v, w \in H^1(0, T; V^*) \cap L^2(0, T; V)$), yields the identity

$$\begin{aligned} & \int_{\Omega} (\bar{\rho}(x, T) - \rho_T(x)) \xi(x, T) dx + \int_0^T \int_{\Omega} \beta_2(\bar{\mu} - \mu_T) \eta dx dt \\ &= \int_0^T \int_{\Gamma} \alpha q (v - \bar{u}) d\sigma dt. \end{aligned} \tag{4.74}$$

The variational inequality (4.73) is thus a direct consequence of Corollary 4.9. ■

References

- [1] V. BARBU, “*Nonlinear semigroups and differential equations in Banach spaces*”, Noordhoff, Leyden, 1976.
- [2] H. BREZIS, “*Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*”, North-Holland Math. Stud. **5**, North-Holland, Amsterdam, 1973.
- [3] P. COLLI, G. GILARDI, P. PODIO-GUIDUGLI, AND J. SPREKELS, *Existence and uniqueness of a global-in-time solution to a phase segregation problem of the Allen-Cahn type*, Math. Models Methods Appl. Sci. **20** (2010), pp. 519–541.
- [4] P. COLLI, G. GILARDI, P. PODIO-GUIDUGLI, AND J. SPREKELS, *A temperature-dependent phase segregation problem of the Allen-Cahn type*, Adv. Math. Sci. Appl. **20** (2010), pp. 219–234.
- [5] P. COLLI, G. GILARDI, P. PODIO-GUIDUGLI, AND J. SPREKELS, *Well-posedness and long-time behavior for a nonstandard viscous Cahn-Hilliard system*, SIAM J. Appl. Math. **71** (2011), pp. 1849–1870.
- [6] P. COLLI, G. GILARDI, P. PODIO-GUIDUGLI, AND J. SPREKELS, *An asymptotic analysis for a nonstandard viscous Cahn-Hilliard system*. WIAS Preprint No. 1652, Berlin 2011. Accepted for publication in Discrete Contin. Dyn. Syst. Ser. S.
- [7] P. COLLI, G. GILARDI, P. PODIO-GUIDUGLI, AND J. SPREKELS, *Distributed optimal control of a nonstandard system of phase field equations*, WIAS Preprint No. 1630, Berlin 2011. In print in Contin. Mech. Thermodyn.
- [8] J. A. GRIEPENTROG, *Maximal regularity for nonsmooth parabolic problems in Sobolev-Morrey spaces*, Adv. Differential Equations **12** (2007), pp. 1031–1078.

- [9] M. HEINKENSCHLOSS, *The numerical solution of a control problem governed by a phase field model*, Optim. Methods Softw. **7** (1997), pp. 211–263.
- [10] M. HEINKENSCHLOSS, AND F. TRÖLTZSCH, *Analysis of the Lagrange-SQP-Newton method for the control of a phase field equation*, Control Cybernet. **28** (1999), pp. 178-211.
- [11] K.-H. HOFFMANN, AND L. JIANG, *Optimal control problem of a phase field model for solidification*, Numer. Funct. Anal. Optim. **40** (1992), pp. 11–27.
- [12] O. A. LADYŽENSKAYA, V. A. SOLONNIKOV, AND N. N. URAL'CEVA, “*Linear and quasilinear equations of parabolic type*”, Trans. Amer. Math. Soc. Vol. **23**, Providence, Rhode Island, 1968.
- [13] C. LEFTER, AND J. SPREKELS, *Control of a phase field system modeling non-isothermal phase transitions*, Adv. Math. Sci. Appl. **17** (2007), pp. 181–194.
- [14] J.-L. LIONS, “*Quelques méthodes de résolution des problèmes aux limites non linéaires*”, Dunod Gauthier-Villars, Paris, 1969.
- [15] J.-L. LIONS, E. MAGENES, “*Problèmes aux limites non homogènes et applications, Vol. I*”, Dunod, Paris, 1968.
- [16] P. PODIO-GUIDUGLI, *Models of phase segregation and diffusion of atomic species on a lattice*, Ric. Mat. **55** (2006), pp. 105–118.
- [17] J. SIMON, *Compact sets in the space $L^p(0, T; B)$* , Ann. Mat. Pura. Appl. **146** (1987), pp. 65–96.
- [18] F. TRÖLTZSCH, “*Optimal Control of Partial Differential Equations: Theory, Methods and Applications*”, Graduate Studies in Mathematics Vol. **112**, American Mathematical Society, Providence, Rhode Island, 2010.