

**Weierstraß-Institut**  
**für Angewandte Analysis und Stochastik**  
**Leibniz-Institut im Forschungsverbund Berlin e. V.**

Preprint

ISSN 2198-5855

**Convergence to equilibrium in  
energy-reaction-diffusion systems using  
vector-valued functional inequalities**

*Dedicated to Peter Markowich on the occasion of his sixtieth birthday.*

Alexander Mielke<sup>1,2</sup>, Markus Mittnenzweig<sup>1</sup>

submitted: December 9, 2016

<sup>1</sup> Weierstrass Institute  
Mohrenstraße 39  
10117 Berlin  
Germany

E-Mail: alexander.mielke@wias-berlin.de  
markus.mittnenzweig@wias-berlin.de

<sup>2</sup> Institut für Mathematik  
Humboldt-Universität zu Berlin  
Rudower Chaussee 25  
12489 Berlin-Adlershof  
Germany

No. 2349  
Berlin 2016



---

2010 *Mathematics Subject Classification.* 35K57, 35B40, 35Q79, 92E20.

*Key words and phrases.* Energy-reaction-diffusion systems, vector-valued inequalities, cross diffusion, log-Sobolev inequality, entropy functional, exponential decay of relative entropy, convexity method.

*Acknowledgments.* The research of M.M. was supported by ERC via AdG 267802 AnaMultiScale, and A.M. was partially supported by the Einstein Stiftung Berlin via ECMath project SE2.

Edited by  
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)  
Leibniz-Institut im Forschungsverbund Berlin e. V.  
Mohrenstraße 39  
10117 Berlin  
Germany

Fax: +49 30 20372-303  
E-Mail: [preprint@wias-berlin.de](mailto:preprint@wias-berlin.de)  
World Wide Web: <http://www.wias-berlin.de/>

## Abstract

We discuss how the recently developed energy-dissipation methods for reaction-diffusion systems can be generalized to the non-isothermal case. For this we use concave entropies in terms of the densities of the species and the internal energy, with the important feature, that the equilibrium densities may depend on the internal energy. Using the log-Sobolev estimate and variants for lower-order entropies as well as estimates for the entropy production of the nonlinear reactions we give two methods to estimate the relative entropy by the total entropy production, namely a somewhat restrictive convexity method, which provides explicit decay rates, and a very general, but weaker compactness method.

## 1 Introduction

The aim of this paper is to generalize the recently developed methods of uniform exponential decay for the isothermal case (see [FeT16, DFT16, MHM15] as well as the references therein) and the older and somewhat forgotten, but very elegant compactness method developed for semiconductor problems in [GIH97]. A first step towards the non-isothermal case was done in [HH\*16], where a more explicit approach close to [DeF06, FeT16] was developed. Here we want to show that also the convexity method from [MHM15] and the compactness method from [GIH97] can be adapted to energy-reaction diffusion systems (ERDS).

Following [AGH02, Mie11b, Mie13] we use the internal energy density  $u(t, \cdot) : \Omega \rightarrow \mathbb{R}$  as the main thermodynamic variable, while the absolute temperature is a dependent variable obtained from the constitutive entropy function  $s = S(x, \mathbf{c}, u)$  via

$$\theta(t, x) = \frac{1}{\partial_u S(x, \mathbf{c}(t, x), u(t, x))},$$

where  $\mathbf{c} = (c_1, \dots, c_I) \in [0, \infty[^I$  is the vector of concentrations. The major advantage is that thermodynamics imposes that  $S(x, \cdot, \cdot) : [0, \infty[^{I+1} \rightarrow \mathbb{R}$  is a concave function that allows us to apply entropy production (EEP) estimates. Moreover,  $S(x, \mathbf{c}, u)$  is a strictly increasing function in  $u$  to make the absolute temperature non-negative.

Our typical model for relevant entropy relations is given by

$$S(x, \mathbf{c}, u) = s(u) - \mathfrak{B}(\mathbf{c} | \mathbf{w}(x, u)) = \widehat{s}(x, u) + \sum_{i=1}^I \left( c_i \log(w_i(x, u)) - \lambda_B(c_i) \right), \quad (1.1)$$

where the Boltzmann function  $\lambda_B$  and the relative Boltzmann entropy are defined via

$$\lambda_B(z) := z \log z - z + 1 \quad \text{and} \quad \mathfrak{B}(\mathbf{c} | \mathbf{b}) := \sum_{i=1}^I b_i \lambda_B(c_i/b_i). \quad (1.2)$$

Here each  $w_i(x, \cdot) : [0, \infty[ \rightarrow [0, \infty[$  is a non-decreasing and concave function as well as  $\widehat{s}(x, \cdot) : u \mapsto s(x, u) + I - \sum_{i=1}^I w_i(x, u)$ , see the thermodynamic discussions in Section 2 where Proposition 2.1 provides the necessary concavity result. From  $\partial_{\mathbf{c}} S(x, \mathbf{c}, u) = (\log(c_i/w_i(x, u)))_{i=1, \dots, I}$  we see that the vector  $\mathbf{w}(x, u) = (w_i(x, u))_{i=1, \dots, I}$  gives the thermodynamic equilibrium concentrations for a given internal energy  $u$ . Further on, we assume that the constitutive function  $S(x, \mathbf{c}, u)$ , and hence  $w_i$  do not depend on  $x$ , i.e. the material

is homogeneous. However, following the ideas in [GHI97] and [MHM15, Sec. 5.1], it should be possible to generalize the methods developed here also to the inhomogeneous case.

One way of generating thermodynamically consistent energy-reaction-diffusion systems (ERDS) on a physical domain  $\Omega \subset \mathbb{R}^d$  is that of using the theory of gradient systems. For this we use the entropy functional

$$\mathcal{S}(\mathbf{c}, u) := \int_{\Omega} S(\mathbf{c}(x), u(x)) \, dx$$

and an Onsager operator  $\mathbb{K}(\mathbf{c}, u) = \mathbb{K}_{\text{diff}}(\mathbf{c}, u) + \mathbb{K}_{\text{R}}(\mathbf{c}, u)$  modeling the entropy production by diffusion or heat transfer and by reaction, respectively. We always assume that they are of the form

$$\mathbb{K}_{\text{diff}}(\mathbf{c}, u) \begin{pmatrix} \boldsymbol{\zeta} \\ \mu \end{pmatrix} = -\operatorname{div} \left( \mathbb{M}(\mathbf{c}, u) \nabla \begin{pmatrix} \boldsymbol{\zeta} \\ \mu \end{pmatrix} \right) \quad \text{and} \quad \mathbb{K}_{\text{R}}(\mathbf{c}, u) = \begin{pmatrix} \mathbb{L}(\mathbf{c}, u) & 0 \\ 0 & 0 \end{pmatrix}.$$

True Onsager operators  $\mathbb{K}$  satisfy the symmetry (reciprocal) relation (cf. [Ons31])  $\mathbb{K} = \mathbb{K}^*$ . Moreover, the second law of thermodynamics enforces  $\mathbb{K}$ . Obviously, this is satisfied if the matrices  $\mathbb{M}$  and  $\mathbb{L}$  are symmetric and positive semi-definite. For our EEP estimates the symmetry will not be important, since we will only rely on the positive semi-definiteness

$$\mathbf{A} : \mathbb{M}(\mathbf{c}, u) \mathbf{A} \geq 0 \quad \text{and} \quad \boldsymbol{\zeta} \cdot \mathbb{L}(\mathbf{c}, u) \boldsymbol{\zeta} \geq 0.$$

Of course, we will need quantitative lower bounds for  $\mathbb{M}$  and  $\mathbb{L}$ . Indeed, starting from the entropy functional  $\mathcal{S}$  we consider the ERDS for  $\mathbf{y} = (\mathbf{c}, u)$ , which is derived formally from  $\dot{\mathbf{y}} = \mathbb{K}(\mathbf{y}) \text{D}\mathcal{S}(\mathbf{y})$  giving

$$\begin{pmatrix} \dot{\mathbf{c}} \\ \dot{u} \end{pmatrix} = -\operatorname{div} \left( \mathbb{M}(\mathbf{c}, u) \begin{pmatrix} \nabla \text{D}_{\mathbf{c}} \mathcal{S}(\mathbf{c}, u) \\ \nabla \partial_u \mathcal{S}(\mathbf{c}, u) \end{pmatrix} \right) + \begin{pmatrix} \mathbb{L}(\mathbf{c}, u) \text{D}_{\mathbf{c}} \mathcal{S}(\mathbf{c}, u) \\ 0 \end{pmatrix} \quad \text{in } \Omega, \\ \boldsymbol{\nu} \cdot \nabla \text{D}_{\mathbf{c}} \mathcal{S}(\mathbf{c}, u) = \mathbf{0} \quad \text{and} \quad \boldsymbol{\nu} \cdot \nabla \partial_u \mathcal{S}(\mathbf{c}, u) = 0 \quad \text{on } \partial\Omega.$$

Obviously the divergence-form of the equation for  $u$  together with the no-flux boundary condition imply that the average internal energy

$$\overline{u(t)} := \frac{1}{|\Omega|} \int_{\Omega} u(t, x) \, dx$$

is constant along solutions of the ERDS. Throughout this work we assume that  $\Omega \subset \mathbb{R}^d$  is a bounded domain with Lipschitz boundary and, without loss of generality, with unit mass, i.e.  $|\Omega| = 1$ . Thus, averages on  $\Omega$  are simply equal to the integral over  $\Omega$ .

Typically in RDS there are further conserved quantities (like the total atomic masses or the charges in semiconductor models), which we assume to be given by a matrix  $\mathbf{Q} : \mathbb{R}^I \rightarrow \mathbb{R}^m$  such that  $\mathbf{Q} \mathbb{L}(\mathbf{c}, u) \boldsymbol{\zeta} = \mathbf{0}$ . Then, again using the non-flux boundary conditions, we see that

$$\mathbf{Q} \overline{\mathbf{c}(t)} = \int_{\Omega} \mathbf{Q} \mathbf{c}(t, x) \, dx \in \mathbb{R}^m$$

is constant along solutions. In particular, we define the associated flow-invariant subsets

$$\mathfrak{S}(\mathbf{q}, \mathbf{U}) := \left\{ (\mathbf{c}, u) \in L^1(\Omega)^{I+1} \mid c_i, u \geq 0, \overline{u} = \mathbf{U}, \mathbf{Q} \overline{\mathbf{c}} = \mathbf{q} \right\}.$$

The strictly concave functional  $\mathcal{S}$  has a unique maximizer, which is spatially constant and which we denote by  $\mathbf{w}(\mathbf{q}, \mathbf{U}) \in ]0, \infty[^I$ . With this we define the relative entropy

$$\begin{aligned} \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) &:= \int_{\Omega} H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}(x), u(x)) \, dx \text{ with} \\ H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) &= S(\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U}) + \text{DS}(\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U}) \cdot \begin{pmatrix} \mathbf{c} - \mathbf{w}(\mathbf{q}, \mathbf{U}) \\ u - \mathbf{U} \end{pmatrix} - S(\mathbf{c}, u). \end{aligned}$$

Hence, we have  $\mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) \geq 0$  with equality if and only if  $(\mathbf{c}, u) = (\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$ .

The entropy production  $\mathcal{P}$  is now defined as the increase of  $\mathcal{S}$  along solutions  $\mathbf{y}(t) = (\mathbf{c}(t), u(t))$ , namely

$$\mathcal{P}(\mathbf{y}) := \frac{d}{dt} \mathcal{S}(\mathbf{y}) = \langle \text{DS}(\mathbf{y}), \mathbb{K}(\mathbf{y}) \text{DS}(\mathbf{y}) \rangle,$$

which has a diffusive contribution  $\mathcal{P}_D$  and a reactive contribution  $\mathcal{P}_R$  given by

$$\begin{aligned} \mathcal{P}_D(\mathbf{y}) &:= \int_{\Omega} \nabla \mathbf{y}(x) : \mathbb{W}(\mathbf{y}(x)) \nabla \mathbf{y}(x) \, dx \quad \text{and} \quad \mathcal{P}_R(\mathbf{y}) := \int_{\Omega} P_R(\mathbf{y}(x)) \, dx \\ \text{with } \mathbb{W}(\mathbf{y}) &= \text{D}^2 S(\mathbf{y}) \mathbb{M}(\mathbf{y}) \text{D}^2 S(\mathbf{y}) \quad \text{and} \quad P_R(\mathbf{y}) = \text{DS}(\mathbf{y}) \cdot \mathbb{L}(\mathbf{y}) \text{DS}(\mathbf{y}). \end{aligned} \quad (1.3)$$

The aim of this work is to provide methods to derive entropy entropy-production (EEP) estimates in the form

$$\forall (\mathbf{q}, \mathbf{U}) \in \mathfrak{Q} \exists K(\mathbf{q}, \mathbf{U}) > 0 \forall (\mathbf{c}, u) \in \mathfrak{S}(\mathbf{q}, \mathbf{U}) : \mathcal{P}(\mathbf{c}, u) \geq K(\mathbf{q}, u) \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u), \quad (1.4)$$

which can be seen as a vector-valued generalization of the usually scalar functional inequalities of log-Sobolev type.

Indeed, we emphasize that we do neither address the question of existence of solutions for equations of the type (1.3) (see [Fis15, Jün15, DD\*16] and the references therein for recent advances) nor the question in which norms the exponential decay of of the solutions can be established. We are content with the construction of  $K(\mathbf{q}, \mathbf{U}) > 0$ , which implies that along sufficiently good notions of weak solutions we have the uniform exponential decay of the relative entropy, namely

$$(\mathbf{c}(0), u(0)) \in \mathfrak{S}(\mathbf{q}, \mathbf{U}) \implies \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}(t), u(t)) \leq e^{-K(\mathbf{q}, \mathbf{U})t} \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}(0), u(0)) \text{ for all } t \geq 0.$$

Clearly, this implies convergence  $(\mathbf{c}(t), u(t)) \rightarrow (\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$  in  $L^1(\Omega)$  when exploiting a suitable variant of the Csiscár-Kullback-Pinsker estimate.

We discuss two possible strategies to establish the fundamental EEP estimate (1.4). Both rely on estimating  $\mathbb{W}$  and  $P_R$  by suitable simplified functions, namely

$$\mathbb{W}(\mathbf{c}, u) \geq \delta_* \text{diag}\left(\frac{1}{c_1}, \dots, \frac{1}{c_I}, \frac{1}{u^{2-\gamma}}\right) \text{ and} \quad (1.5a)$$

$$P_R(\mathbf{c}, u) \geq \kappa_* \widehat{P}_R(\mathbf{c}, u) \text{ with } \widehat{P}_R(\mathbf{c}, u) := \sum_{r=1}^I G\left(\frac{\mathbf{c}^{\alpha^r}}{\mathbf{w}(u)^{\alpha^r}}, \frac{\mathbf{c}^{\beta^r}}{\mathbf{w}(u)^{\beta^r}}\right) \quad (1.5b)$$

with  $G(a, b) = (a-b)(\log a - \log b) \geq 0$ . We show that these estimates can be realized for ERDS of the form

$$\begin{pmatrix} \dot{\mathbf{c}} \\ \dot{u} \end{pmatrix} = \begin{pmatrix} (\delta_i \Delta c_i)_i \\ \delta_u \Delta u \end{pmatrix} + \begin{pmatrix} \mathbf{R}(\mathbf{c}, u) \\ 0 \end{pmatrix} \text{ with } \mathbf{R}(\mathbf{c}, u) = \sum_{r=1}^R \kappa_r \left( \frac{\mathbf{c}^{\alpha^r}}{\mathbf{w}(u)^{\alpha^r}} - \frac{\mathbf{c}^{\beta^r}}{\mathbf{w}(u)^{\beta^r}} \right) (\beta^r - \alpha^r), \quad (1.6)$$

if the reaction-rate coefficients satisfy  $\kappa_r \geq \kappa_*$  and if for all  $i = 1, \dots, I$  there exist  $\rho_i > 0$  such that the diffusion coefficients  $\delta_i$  and the equilibrium functions  $w_i$  satisfy the relations

$$\rho_i \leq \frac{4\delta_u(\delta_i - \delta_*)}{(\delta_i + \delta_u)^2} \quad \text{and} \quad \frac{1}{\rho_i} (w'_i(u))^2 \leq (w'_i(u))^2 - w_i(u)w''_i(u) \quad \text{for } u > 0,$$

see Proposition 2.2, which establishes (1.5a). We interpret these relations as a bound on cross-diffusion, since the heat equation without cross-diffusion should have the form

$$\dot{u} = -\operatorname{div} \left( M_{u,u}(\mathbf{c}, u) \nabla \frac{1}{\theta} \right) \quad \text{with} \quad \frac{1}{\theta} = \partial_u S(\mathbf{c}, u) = s'(u) + \sum_1^I c_i w'_i(u) / w_i(u),$$

see the discussion in Section 2.2. The assumption (1.5a) provides the lower bound

$$\mathcal{P}_D(\mathbf{c}, u) \geq \delta_* p_\Omega \int_\Omega \left( \bar{u} \lambda_B \left( \frac{u}{\bar{u}} \right) + \sum_{i=1}^I \bar{c}_i \lambda_B \left( \frac{c_i}{\bar{c}_i} \right) \right) dx$$

with  $p_\Omega = \min\{\rho(\Omega, \gamma, 1), \rho(\Omega, 1, 1)\}$ , where the optimal generalized log-Sobolev constants  $\rho(\Omega, \gamma, \alpha)$  are defined in (3.6).

For  $\widehat{P}_R$  of the entropy production we impose the finite-dimensional conditions

$$\forall (\mathbf{q}, \mathbf{U}) \in \mathfrak{Q} \exists K_R(\mathbf{q}, \mathbf{U}) > 0 \forall \mathbf{c} \text{ with } \mathbf{Q}\mathbf{c} = \mathbf{q} : \widehat{P}_R(\mathbf{c}, \mathbf{U}) \geq K_R(\mathbf{q}, \mathbf{U}) \mathfrak{B}(\mathbf{c} | \mathbf{w}(\mathbf{q}, \mathbf{U})), \quad (1.7)$$

which, for our mass-action kinetics, is equivalent to the unique-equilibrium condition that  $\mathbf{c} = \mathbf{w}(\mathbf{q}, \mathbf{U})$  is the only solution of  $\mathbf{R}(\mathbf{c}, \mathbf{U}) = 0$  satisfying  $\mathbf{Q}\mathbf{c} = \mathbf{q}$ .

The counterpart of the convexity method for isothermal RDS developed in [MHM15] is formulated in Theorem 3.2. Assuming that there exists a  $\mu(\mathbf{U}) \geq 0$  such that the function

$$(\mathbf{c}, u) \mapsto \mu(\mathbf{U}) \left( \mathbf{U}^\gamma \lambda_B(u/\mathbf{U}) + \sum_i \lambda_B(c_i) \right) + \widehat{P}_R(\mathbf{c}, u) \quad \text{is convex}$$

we obtain the constructive lower bound

$$K(\mathbf{q}, \mathbf{U}) \geq \frac{1}{C_H(\mathbf{q}, \mathbf{U})} \min \left\{ \kappa_* K_R(\mathbf{q}, \mathbf{U}), \delta_* p_\Omega \frac{K_R(\mathbf{q}, \mathbf{U})}{\mu(\mathbf{U}) + K_R(\mathbf{q}, \mathbf{U})} \right\},$$

where the  $C_H$  is characterized in Lemma 3.1.

In Section 4 we discuss the generalization of the compactness argument developed in [GIH97] for isothermal semiconductor models. There only the case  $d \leq 2$  is treated because of the coupling via the electric potential, but without this there is no dimension restriction, see also [Mie16]. In contrast to the convexity method the theory is much more general, but the result is also significantly weaker, since the constant  $K(\mathbf{q}, \mathbf{U})$  in the EEP estimate is not constructive and moreover, it also depends on an upper bound  $M$  for  $\mathcal{H}_{\mathbf{q}, \mathbf{U}}$ , namely

$$\begin{aligned} \forall (\mathbf{q}, \mathbf{U}) \in \mathfrak{Q} \quad M > 0 \exists K_M(\mathbf{q}, \mathbf{U}) > 0 \forall (\mathbf{c}, u) \in \mathfrak{S}(\mathbf{q}, \mathbf{U}) : \\ \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) \leq M \implies \mathcal{P}(\mathbf{c}, u) \geq K_M(\mathbf{q}, \mathbf{U}) \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u), \end{aligned} \quad (1.8)$$

## 2 Thermodynamical modeling

Here we describe the general approach to obtain thermodynamically consistent models for ERDS. We start from a concave entropy and derive the coupled system of PDEs as a formal

gradient system as introduced in [Mie11a, Mie13]. Using the Boltzmann entropy for the concentrations it is possible to reproduce the exact mass-action kinetics, if it satisfies a detailed-balance condition, which we assume throughout this work. The detailed-balance seems to be necessary for the gradient structure; however, the uniform exponential decay estimates can be obtained under the weaker notion of complex balancing, see [DFT16, Mie16]. Indeed it would be straightforward to generalize the present work on ERDS to the case of complex-balanced kinetics, but we refrain from doing so in the interest of a simpler notation.

## 2.1 The entropy functional

We study general systems that are thermodynamically consistent, in the sense that they can be described as a gradient system in terms of the total entropy. We allow for  $I$  different species  $X_1, \dots, X_I$ , whose concentrations denoted by  $c_i(t, x) \geq 0$  form the density vector  $\mathbf{c} = (c_1, \dots, c_I) : \Omega \rightarrow [0, \infty[^I$ . In addition, we have the scalar internal energy density  $u(t, x) \in \mathbb{R}$ . The total entropy  $\mathcal{S}$  is the integral over the entropy density  $s = S(x, \mathbf{c}, u)$ , where  $S(x, \cdot)$  is a strictly concave function of  $(\mathbf{c}, u) \in [0, \infty[^I \times \mathbb{R}$ , see e.g. [LiY99]. Thus, we set

$$\mathcal{S}(\mathbf{c}, u) = \int_{\Omega} S(x, \mathbf{c}(x), u(x)) \, dx.$$

The absolute temperature is now a dependent variable given by

$$\theta(t, x) = \Theta(x, \mathbf{c}, u) := \frac{1}{\partial_u S(x, \mathbf{c}(t, x), u(t, x))}.$$

At a given point  $x \in \Omega$  and for a fixed internal energy the equilibrium densities are obtained by maximizing the  $\mathbf{c} \mapsto S(x, \mathbf{c}, u)$  providing the vector

$$\mathbf{w}(x, u) = (w_1(x, u), \dots, w_I(x, u)) \in ]0, \infty[^I,$$

i.e.  $D_{\mathbf{c}} S(x, \mathbf{w}(x, u), u) = 0$ . One of the major difficulties in this work is that the vector  $\mathbf{w}$  depends on  $u$ , i.e. the local equilibrium concentrations depend on  $u$ , and hence implicitly on  $\theta$ .

So far, we also allowed for an explicit dependence on the material point  $x \in \Omega$ . Such a dependence is crucial for ERDS posed inside of solids, e.g. in semiconductor physics where  $c_i$  describes the concentrations of charge carriers. The dependence on  $x$  then models the domains of the different materials including doping structures. However, for our work this dependence is not essential, so we will drop it for notational simplicity. Following [GIH97] it should be possible to adapt the subsequent analysis to the general case by using suitable relative densities.

For later usage we give a suitable class of entropy functions based on the Boltzmann entropy using the function  $\lambda_B(z) := z \log z - z + 1$ :

$$S(\mathbf{c}, u) = s(u) - \sum_{i=1}^I w_i(u) \lambda_B\left(\frac{c_i}{w_i(u)}\right) = \widehat{s}(u) + \sum_{i=1}^I \left( c_i \log(w_i(u)) - \lambda_B(c_i) \right), \quad (2.1)$$

$$\text{where } \widehat{s}(u) = s(u) + I - \sum_{i=1}^I w_i(u).$$

The following result gives necessary and sufficient conditions on  $s$  and  $w_i$  for the concavity of this function  $S$ .

**Proposition 2.1 (Concavity and monotonicity of  $S$ )** Consider smooth functions  $s : [0, \infty[ \rightarrow \mathbb{R}$  and  $w_i : [0, \infty[ \rightarrow [0, \infty[$  and define  $S : [0, \infty[^{I+1} \rightarrow \mathbb{R}$  as in (2.1). Then,  
(a)  $S$  is concave if and only if

$$w_1, w_2, \dots, w_I, \text{ and } \widehat{s} \text{ are concave;} \quad (2.2a)$$

(b)  $S$  is non-decreasing in  $u$  if and only if

$$w_1, w_2, \dots, w_I, \text{ and } \widehat{s} \text{ are increasing.} \quad (2.2b)$$

**Proof:** We simply calculate the second derivative, which yields

$$D^2S(\mathbf{c}, u) = \begin{pmatrix} -\frac{1}{c_1} & 0 & \dots & 0 & \frac{w'_1(u)}{w_1(u)} \\ 0 & -\frac{1}{c_2} & 0 & & \frac{w'_2(u)}{w_2(u)} \\ \vdots & & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & -\frac{1}{c_I} & \frac{w'_I(u)}{w_I(u)} \\ \frac{w'_1(u)}{w_1(u)} & \frac{w'_2(u)}{w_2(u)} & \dots & \frac{w'_I(u)}{w_I(u)} & \partial_u^2 S(\mathbf{c}, u) \end{pmatrix} \quad (2.3)$$

with  $\partial_u^2 S(\mathbf{c}, u) = \widehat{s}''(u) + \sum_{i=1}^I c_i \frac{w_i(u)w_i''(u) - (w_i'(u))^2}{(w_i(u))^2} < 0$ . In particular, we find

$$\begin{pmatrix} \zeta \\ \tau \end{pmatrix} \cdot D^2S(\mathbf{c}, u) \begin{pmatrix} \zeta \\ \tau \end{pmatrix} = \left( \widehat{s}''(u) + \sum_{i=1}^I c_i \frac{w_i''(u)}{w_i(u)} \right) \tau^2 - \sum_{i=1}^I c_i \left( \frac{\zeta_i}{c_i} + \frac{w_i'(u)}{w_i(u)} \tau \right)^2,$$

and (2.2a) follows immediately since  $c_i$  and  $u$  may range freely in  $[0, \infty[$ .

For (2.2b)part (b) we simply use  $\partial_u S(\mathbf{c}, u) = \widehat{s}'(u) + \sum_{i=1}^I c_i w_i'(u)/w_i(u)$  and the fact that  $c_i$  and  $u$  may range freely in  $[0, \infty[$ . ■

The simplest example of a suitable entropy functional  $S(\mathbf{c}, u)$  (cf. [Mie13, (3.9)] or [HH\*16, Sec. 4.3]) is given by the choices

$$\widehat{s}(u) = s_0 \log u \quad \text{and} \quad w_i(u) = r_i u^{\rho_i} \quad \text{with} \quad s_0, r_i > 0 \quad \text{and} \quad \rho_i \in [0, 1].$$

With  $\boldsymbol{\rho} = (\rho_1, \dots, \rho_I)$  we then have the explicit relations

$$S(\mathbf{c}, u) = (s_0 + \boldsymbol{\rho} \cdot \mathbf{c}) \log u + \tilde{\sigma}(\mathbf{c}) \quad \text{with} \quad \tilde{\sigma}(\mathbf{c}) := \sum_{i=1}^I \left( c_i \log r_i - \lambda_B(c_i) \right) \quad \text{and}$$

$$\theta = \Theta(\mathbf{c}, u) = u / (s_0 + \boldsymbol{\rho} \cdot \mathbf{c}), \quad u = U(\mathbf{c}, \theta) = (s_0 + \boldsymbol{\rho} \cdot \mathbf{c}) \theta.$$

Moreover, we may express the entropy and the free energy  $\psi = u - \theta s$  in  $\mathbf{c}$  and  $\theta$ , which gives, using the shorthand  $c^\rho := s_0 + \boldsymbol{\rho} \cdot \mathbf{c} > 0$ ,

$$\overline{S}(\mathbf{c}, \theta) = S(\mathbf{c}, U(\mathbf{c}, \theta)) = c^\rho (\log \theta + \log c^\rho) + \tilde{\sigma}(\mathbf{c}),$$

$$\overline{\psi}(\mathbf{c}, \theta) = U(\mathbf{c}, \theta) - \theta \overline{S}(\mathbf{c}, \theta) = c^\rho (1 - \lambda_B(\theta)) - \theta (\tilde{\sigma}(\mathbf{c}) + c^\rho \log c^\rho).$$

It can be shown that  $\overline{S}$  is concave if and only if  $\rho_i \in [0, 1/2]$  for all  $i$ . Similarly,  $\overline{\psi}(\cdot, \theta)$  is convex if and only if  $\rho_i \in [0, 1/2]$  for all  $i$ . This shows that  $\theta$  is not a good variable to describe ERDS.

This family can be generalized substantially by choosing a continuously differentiable function  $V : [0, \infty[ \rightarrow ]0, \infty[$  such that

$$V'(u) \geq \max\{\rho_i \mid i = 1, \dots, I\} \quad \text{for all } u > 0.$$

Then we define  $\phi : [0, \infty[ \rightarrow ]0, \infty[$  via  $\phi'(u) = \phi(u)/V(u)$ ,  $\phi(0) = \phi_0 \geq 0$  and set

$$w_i(u) = r_i \phi(u)^{\rho_i} \quad \text{for } i = 1, \dots, I \quad \text{and } \widehat{s}(u) = s_0 \log(\phi(u)),$$

with  $r_i$  and  $\rho_i$  as above. The monotonicity of  $w_i$  and  $\widehat{s}$  is obvious, and the concavities follow from

$$w_i''(u) \leq 0 \quad \iff \quad (1 - \rho_i)(\phi')^2 \geq \phi \phi'' \quad \iff \quad V' \geq \rho_i.$$

The result for  $\widehat{s}$  is obtained the same way but with  $\rho = 0$ . The special form with the same  $\phi$  for each  $w_i$  leads to the explicit relation  $V(u) = (s_0 + \boldsymbol{\rho} \cdot \mathbf{c})\theta$ , which now may be nonlinear, but is still monotone.

## 2.2 Diffusion and heat transfer

Our ERDS will be of the general form

$$\begin{pmatrix} \dot{\mathbf{c}} \\ \dot{u} \end{pmatrix} = \operatorname{div} \left( M(\mathbf{c}, u) \nabla \begin{pmatrix} \mathbf{c} \\ u \end{pmatrix} \right) + \begin{pmatrix} \mathbf{R}(\mathbf{c}, u) \\ 0 \end{pmatrix} \text{ in } \Omega, \quad \nabla \begin{pmatrix} \mathbf{c} \\ u \end{pmatrix} \cdot \boldsymbol{\nu} \text{ on } \partial\Omega. \quad (2.4)$$

Here  $M(\mathbf{c}, u)$  could be a tensor of fourth order to model anisotropy, however, only consider  $M(\mathbf{c}, u) \in \mathbb{R}^{(I+1) \times (I+1)}$  such that  $M$  contains the mobilities of the species  $X_i$  with densities  $c_i$ , the heat conductivity  $M_{I+1, I+1}$  as well as the cross terms  $M_{i, I+1}$  and  $M_{I+1, i}$ . Of course, we also allow for cross diffusion between the species, i.e.  $M_{ij}$  may be nonzero for  $i \neq j$ . One way of generating suitable matrices  $M$  is by using an diffusion Onsager operator

$$\mathbb{K}_{\text{diff}}(\mathbf{c}, u) \begin{pmatrix} \zeta \\ \mu \end{pmatrix} := -\operatorname{div} \left( \mathbb{M}(\mathbf{c}, u) \nabla \begin{pmatrix} \zeta \\ \mu \end{pmatrix} \right), \quad \text{where } \mathbb{M}(\mathbf{c}, u) = \mathbb{M}(\mathbf{c}, u)^\top \geq 0.$$

Then, we generate the matrix  $M$  by applying  $\mathbb{K}_{\text{diff}}$  to  $\mathcal{S}$ , namely

$$\begin{pmatrix} \dot{\mathbf{c}} \\ \dot{u} \end{pmatrix} = \operatorname{div} \left( M(\mathbf{c}, u) \nabla \begin{pmatrix} \mathbf{c} \\ u \end{pmatrix} \right) = \mathbb{K}(\mathbf{c}, u) \operatorname{D}\mathcal{S}(\mathbf{c}, u). \quad (2.5)$$

By using that  $\begin{pmatrix} \zeta \\ \mu \end{pmatrix} = \operatorname{D}\mathcal{S}(\mathbf{c}, u) = \begin{pmatrix} \operatorname{D}_{\mathbf{c}}\mathcal{S}(\mathbf{c}) \\ \operatorname{D}_u\mathcal{S}(\mathbf{c}, u) \end{pmatrix}$  we find the relation

$$M(\mathbf{c}, u) = \mathbb{M}(\mathbf{c}, u) (-\operatorname{D}^2\mathcal{S}(\mathbf{c}, u)).$$

Hence  $M$  may be non-symmetric, but it is the product of two symmetric matrices, which implies that it is still diagonalizable with real eigenvalues.

Of course, the simplest choice for  $\mathbb{M}(\mathbf{c}, u)$ , which is often used nevertheless, is simply to take  $\mathbb{M}$  as a scalar multiple  $\delta$  of  $(-\operatorname{D}^2\mathcal{S}(\mathbf{c}, u))^{-1}$  like in the case of Wasserstein diffusion, see [JKO98, Ott01, LiM13]. It is still an open problem to construct sufficiently large classes of mobility tensors  $\mathbb{M}$ , such that  $M = \mathbb{M}(-\operatorname{D}^2\mathcal{S})$  is sufficiently simple to allow for existence results for solutions.

In the present work we do not discuss the structure of  $\mathbb{M}$ , since we do not need the pure diffusion equation (2.4) with  $\mathbf{R} \equiv 0$  to be a gradient system with respect to  $\mathcal{S}$ . The Onsager form in (2.5) and the choice  $M = -\mathbb{M}\operatorname{D}^2\mathcal{S}$  is just an easy way to generate a large class of admissible mobilities. Our focus concerns the entropy production generated by  $M$  along solutions of  $\dot{\mathbf{y}} = \operatorname{div}(M(\mathbf{y})\nabla\mathbf{y})$ , viz.

$$\frac{d}{dt}\mathcal{S}(\mathbf{y}) = \int_{\Omega} \operatorname{D}\mathcal{S}(\mathbf{y}) \cdot \dot{\mathbf{y}} \, dx = \int_{\Omega} \nabla\mathbf{y} \cdot (-M(\mathbf{y})\operatorname{D}^2\mathcal{S}(\mathbf{y})) \nabla\mathbf{y} \, dx =: \mathcal{P}_{\text{diff}}(\mathbf{y}). \quad (2.6)$$

Thus, we have to choose  $M$  such that  $\mathcal{P}_{\text{diff}}(\mathbf{y}) \geq 0$  for all  $\mathbf{y}$ . This is clearly implied by the condition that the symmetric part of  $-M(\mathbf{y})D^2S(\mathbf{y})$  is positive semi-definite. However, using suitable integration by parts (which is equivalent to exploit null-Lagrangians) there may be more cases, see e.g. the theory developed in [JüM06].

To obtain our desired uniform exponential decay estimates we need a suitable quantitative lower bound  $\mathcal{P}_{\text{diff}}(\mathbf{y}) \geq \delta_* \widehat{\mathcal{P}}_\gamma(\mathbf{y})$ , for which we choose

$$\widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) := \int_{\Omega} \left( \frac{|\nabla u|^2}{u^{2-\gamma}} + \sum_{i=1}^I \frac{|\nabla c_i|^2}{c_i} \right) dx, \quad \text{where } \gamma \in [0, 1[. \quad (2.7)$$

The parameter  $\gamma$  for the internal energy relates to the fact that typical entropy functions  $s$  are of the form  $s(u) = s_1 u^\gamma$  for  $\gamma \in ]0, 1[$  or  $s(u) = s_0 \log u$  for  $\gamma = 0$ , while the Boltzmann entropy  $\lambda_B(z) = z \log z - z + 1$  for the densities  $c_i$  corresponds to the case  $\gamma = 1$ .

The following result provides conditions on the entropy  $S$  in (2.1) and a constant diagonal matrix  $M$  such that  $\mathcal{P}_{\text{diff}}$  can be bounded from below by  $\widehat{\mathcal{P}}_\gamma$ .

**Proposition 2.2 (Entropy production by linear diffusion)** *Assume that  $S$  is given by (2.1) with  $w_i$  and  $\widehat{s}$  strictly increasing and concave, and that there exists  $\gamma, \rho_i \in ]0, 1[$  for  $i = 1, \dots, I$  and  $\sigma > 0$  such that*

$$\forall i, u > 0: \quad -w_i(u)w_i''(u) \geq \frac{1-\rho_i}{\rho_i} (w_i'(u))^2 \quad \text{and} \quad -\widehat{s}''(u) \geq \frac{\widehat{s}_0}{u^{2-\gamma}}. \quad (2.8)$$

*Then, for the constant diagonal matrix  $M(\mathbf{y}) = \text{diag}(\delta_1, \dots, \delta_I, \delta_u)$  with  $\delta_i, \delta_u > 0$  the entropy production  $\mathcal{P}_{\text{diff}}$  defined in (2.6) satisfies the lower bound*

$$\mathcal{P}_{\text{diff}}(\mathbf{c}, u) \geq \delta_* \widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) \quad \text{with } \delta_* = \min \left\{ \delta_u \widehat{s}_0, \left( \delta_i - \frac{\rho_i(\delta_i + \delta_u)^2}{4\delta_u} \right) \mid i = 1, \dots, I \right\} \quad (2.9)$$

for all  $(\mathbf{c}, u)$  with  $\sqrt{c_i}, u^{\gamma/2} \in H^1(\Omega)$ .

**Proof:** An explicit calculation using  $D^2S$  from (2.3) gives  $\mathcal{P}_{\text{diff}}(\mathbf{y}) = \int_{\Omega} Q(\mathbf{y}; \nabla \mathbf{y}) dx$  with

$$Q(\mathbf{c}, u; \nabla c, \nabla u) = -\delta_u \widehat{s}'' |\nabla u|^2 + \sum_{i=1}^I \left( \delta_i \frac{|\nabla c_i|^2}{c_i} - (\delta_i + \delta_u) \frac{w_i'}{w_i} \nabla u \cdot \nabla c_i + \delta_u c_i \frac{w_i'^2 - w_i w_i''}{w_i^2} |\nabla u|^2 \right),$$

where we dropped the argument  $u$  from  $\widehat{s}$  and  $w_i$  for notational simplicity.

Abbreviating  $a_i = w_i'/w_i$  and using the assumption (2.8) we obtain the lower bound

$$\begin{aligned} Q(\mathbf{c}, u; \nabla c, \nabla u) &\geq \frac{\delta_u \widehat{s}_0}{u^{2-\gamma}} |\nabla u|^2 + \sum_{i=1}^I \left( \delta_i \frac{|\nabla c_i|^2}{c_i} - (\delta_i + \delta_u) a_i \nabla u \cdot \nabla c_i + \frac{\delta_u c_i a_i^2}{\rho_i} |\nabla u|^2 \right) \\ &\geq \frac{\delta_u \widehat{s}_0}{u^{2-\gamma}} |\nabla u|^2 + \sum_{i=1}^I \left( \delta_i - \frac{\rho_i(\delta_i + \delta_u)^2}{4\delta_u} \right) \frac{|\nabla c_i|^2}{c_i}, \end{aligned} \quad (2.10)$$

where we estimated by Young's inequality each term in the sum individually. This proves the assertion.  $\blacksquare$

**Remark 2.3 (Dropped mixed terms)** In estimate (2.10) we dropped a positive term to eliminate the mixed terms  $\nabla u \cdot \nabla c_i$ . Dropping the index  $i$  for simplicity and letting

$$\tilde{\kappa} = \frac{2\delta_u}{\rho_i(\delta_i + \delta_u)}, \quad \tilde{\rho} = \frac{\rho_i(\delta_i + \delta_u)^2}{4\delta_u}, \quad \text{and} \quad A(u) = w_i(u)^{\tilde{\kappa}}$$

this term takes the form

$$\begin{aligned} \tilde{\rho} \left( \frac{|\nabla c|^2}{c} - 2A'(u)\nabla u \cdot \nabla c + cA'(u)^2|\nabla u|^2 \right) &= c\tilde{\rho} |\nabla(\log c) - \nabla(\log A(u))|^2 \\ &= c\tilde{\rho} \left| \nabla \log(c/A(u)) \right|^2 = \frac{\tilde{\rho}}{\alpha^2} c^{1-2\alpha} A(u)^{2\alpha} \left| \nabla \left( \frac{c^\alpha}{A(u)^\alpha} \right) \right|^2, \end{aligned}$$

where  $\alpha > 0$  is arbitrary. For a way that this term is exploited, we refer to [HH\*16, Prop. 6.2], where  $w(u) = u^{1/2}$ ,  $\delta_u = \delta$  and  $\rho = \alpha = 1/2$  giving  $A(u) = u$ .

The estimate (2.9) is of course only useful for  $\delta_* > 0$ . For the case  $M = \delta I_{I+1}$  this is the case with  $\delta_* = \delta \min\{1 - \rho_i \mid i = 1, \dots, I\}$ . If the diffusion constants are different we obtain restrictions on the  $\rho_i$ , namely we need

$$\rho_i < \frac{4\delta_i\delta_u}{(\delta_i + \delta_u)^2} \leq 1. \quad (2.11)$$

This provides an interesting connection between the entropy structure and the diffusion mechanisms. Indeed, this connection may be attributed to cross-diffusion, if we take into account that the temperature is defined via  $\theta = 1/\partial_u S(\mathbf{c}, u)$ . In thermodynamical modeling it is usually assumed that heat flow is not driven by  $\nabla u$  but rather by  $\nabla\theta$  (or more precisely by  $\nabla(1/\theta) = \nabla(\partial_u S(\mathbf{c}, u))$ ), see e.g. [AGH02, Mie13]. Thus, our system written in terms of  $(\mathbf{c}, u)$  must have cross-diffusion when written in terms of  $(\mathbf{c}, \theta)$ . More precisely, for  $S$  in (2.1) we have

$$\nabla DS(\mathbf{c}, u) = \left( \begin{array}{c} -\frac{1}{c_i} \nabla c_i + a_i \nabla u \\ (\tilde{s}'' + \sum_1^I c_i a_i') \nabla u + \sum_1^I a_i \nabla c_i \end{array} \right) \quad \text{with} \quad a_i(u) = \frac{w_i'(u)}{w_i(u)}.$$

To generate a term that is always proportional to  $\nabla u$  we need to take a very specific combination of the components of  $\nabla DS(\mathbf{c}, u)$ , namely

$$\nabla(\partial_u S(\mathbf{c}, u)) + \sum_{i=1}^I a_i c_i \nabla(\partial_{c_i} S(\mathbf{c}, u)) = \left( \tilde{s}''(u) + \sum_{i=1}^I c_i \frac{w_i''(u)}{w_i(u)} \right) \nabla u.$$

In turn, this implies that the linear decoupled diffusion system  $\dot{\mathbf{y}} = M\Delta\mathbf{y}$  with the constant matrix  $M = \text{diag}(\delta_1, \dots, \delta_I, \delta_u)$  can be written in the Onsager form (2.5), i.e.

$$\begin{pmatrix} \dot{\mathbf{c}} \\ \dot{u} \end{pmatrix} = M \begin{pmatrix} \Delta\mathbf{c} \\ \Delta u \end{pmatrix} = \begin{pmatrix} (\delta_i \Delta c_i)_{1, \dots, I} \\ \delta_u \Delta u \end{pmatrix} = \mathbb{K}(\mathbf{c}, u) DS(\mathbf{c}, u) = -\text{div} \left( \mathbb{M}(\mathbf{c}, u) \nabla (DS(\mathbf{c}, u)) \right),$$

if the mobility matrix  $\mathbb{M}(\mathbf{c}, u) \in \mathbb{R}^{(I+1) \times (I+1)}$  satisfies the relation

$$\mathbb{M}(\mathbf{c}, u) (0, \dots, 0, 1)^\top = \tilde{m}(\mathbf{c}, u) (c_1 a_1(u), \dots, c_I a_I(u), 1)^\top$$

for a suitable scalar function  $\tilde{m}$ . Recall that the case  $w_i(u) = r_i u^{\rho_i}$  leads to  $a_i(u) = \rho_i/u$ , i.e. we have

$$\mathbb{M}_{i, I+1}(\mathbf{c}, u) = \tilde{m}(\mathbf{c}, u) \frac{\rho_i c_i}{u} \quad \text{for } i = 1, \dots, I.$$

Thus, we may interpret the condition (2.11) on  $\rho_i$  as a condition on the strength of cross-diffusion.

**Remark 2.4** We may check whether  $M = \text{diag}(\delta_1, \dots, \delta_u)$  can actually be generated by an Onsager operator with  $\mathbb{M}$ . For this we use the block inversion formula

$$\left( \begin{array}{c|c} \mathbf{C}^{-1} & \mathbf{C}^{-1}\mathbf{a} \\ \hline (\mathbf{C}^{-1}\mathbf{a})^\top & \sigma_0 \end{array} \right)^{-1} = \left( \begin{array}{c|c} \mathbf{C} + \mu\mathbf{a}\otimes\mathbf{a} & -\mu\mathbf{a} \\ \hline -\mu\mathbf{a}^\top & \mu \end{array} \right) \text{ with } \frac{1}{\mu} = \sigma_0 - \mathbf{a} \cdot \mathbf{C}^{-1}\mathbf{a}.$$

Thus, using (2.3) we can calculate  $\mathbb{M}(\mathbf{c}, u) = M(-D^2S(\mathbf{c}, u))^{-1}$  explicitly, which leads to

$$\mathbb{M}(\mathbf{c}, u) = \begin{pmatrix} \delta_1 c_1 + \mu \left( \frac{c_1 w'_1}{w_1} \right)^2 & \cdots & \mu \frac{c_1 c_I w'_1 w'_I}{w_1 w_I} & \delta_1 \mu c_1 \frac{w'_1}{w_1} \\ \vdots & \ddots & \vdots & \vdots \\ \mu \frac{c_I c_1 w'_I w'_1}{w_I w_1} & \cdots & \delta_I c_I + \mu \left( \frac{c_I w'_I}{w_I} \right)^2 & \delta_I \mu c_I \frac{w'_I}{w_I} \\ \delta_u \mu c_1 \frac{w'_1}{w_1} & \cdots & \delta_u \mu c_I \frac{w'_I}{w_I} & \delta_u \mu \end{pmatrix}$$

with  $\frac{1}{\mu} = -\widehat{s}''(u) - \sum_{i=1}^I c_i \frac{w''_i(u)}{w_i(u)} > 0$ . Obviously, this matrix is only symmetric if  $\delta_i = \delta_u$  for all  $i$ ; however, one can check that the symmetric part of  $\mathbb{M}$  is positive definite under the conditions for which  $\delta_*$  in (2.9) is positive.

It remains a challenging open question to find suitable symmetric and positive semi-definite mobilities  $\mathbb{M}(\mathbf{c}, u)$  and suitable entropies  $S(\mathbf{c}, u)$  such that the induced diffusion matrix  $M(\mathbf{c}, u) = -\mathbb{M}(\mathbf{c}, u)D^2S(\mathbf{c}, u)$  is nontrivial but still simple enough to allow for an existence theory for solutions. Some results are available for the isothermal case (e.g. [BD\*10, LiM13]), but nothing exists in the non-isothermal case.

## 2.3 Mass-action reaction kinetics


Next we discuss the reaction part  $\dot{\mathbf{c}} = \mathbf{R}(\mathbf{c}, u)$  in the ERDS (2.4), which is a simple ODE, where the internal energy  $u > 0$  acts as a parameter. Thus, the theory for the kinetics is very close to the isothermal one, if we take care properly of the parametric dependence on  $u$ .

First we observe, that for a fixed  $u$  our entropy  $\mathbf{c} \mapsto \mathcal{S}(\mathbf{c}, u)$  in (2.1) is, up to a constant, a classical Boltzmann entropy with the thermal equilibrium  $\mathbf{c} = \mathbf{w}(u)$ . Thus, our basic assumption is that  $\mathcal{S}(\cdot, u)$  is a Liapunov function for  $\dot{\mathbf{c}} = \mathbf{R}(\mathbf{c}, u)$ , i.e.

$$P_{\text{react}}(\mathbf{c}, u) := \mathbf{R}(\mathbf{c}, u) \cdot D_{\mathbf{c}}\mathcal{S}(\mathbf{c}, u) \geq 0.$$

Indeed, as for the diffusion part, we want to quantify this relation to obtain exponential decay.

Second, we specify the reaction terms further and assume mass-action kinetics with  $R$  reaction pairs for the species  $X_1, \dots, X_I$  given by



Here  $\alpha_i^r, \beta_i^r \in \mathbb{N}_0$  are the stoichiometric coefficients and  $k_{\text{fw}}^r$  and  $k_{\text{bw}}^r$  are the forward and backward reaction-rate coefficients of the  $r$ th reaction. Defining the stoichiometric vectors  $\boldsymbol{\alpha}^r := (\alpha_i^r)_i$  and  $\boldsymbol{\beta}^r := (\beta_i^r)_i$  in  $\mathbb{N}_0^I$  the associated reaction-rate equation reads

$$\dot{\mathbf{c}} = \mathbf{R}(\mathbf{c}, u) = \sum_{r=1}^R (k_{\text{fw}}^r(\mathbf{c}, u) \mathbf{c}^{\boldsymbol{\alpha}^r} - k_{\text{bw}}^r(\mathbf{c}, u) \mathbf{c}^{\boldsymbol{\beta}^r}) (\boldsymbol{\beta}^r - \boldsymbol{\alpha}^r).$$

Third, we assume the *detailed-balance condition* for the  $R$  reaction pairs

$$(DBC) \quad \forall r = 1, \dots, R \quad \forall u > 0 : \quad k_{\text{fw}}^r(u) (\mathbf{w}(u))^{\boldsymbol{\alpha}^r} = k_{\text{bw}}^r(u) (\mathbf{w}(u))^{\boldsymbol{\beta}^r} =: \kappa_r(u), \quad (2.12)$$

which means that all the reactions pairs are simultaneously in balance at the thermodynamic equilibrium  $\mathbf{w}(u)$ . We emphasize that this condition is stronger than needed: following [DFT16, Mie16] it should be possible to generalize our arguments to the case of complex-balanced reaction-rate equation, where still a quantitative estimate between the entropy production and the entropy is possible. For simplicity we stay with the DBC (2.12) and follow the approach developed in [Mie11a, Mie13, HH\*16]. Using  $D_c S(\mathbf{c}, u) = -(\log(c_i/w_i(u)))_{i=1, \dots, I}$  we obtain the special form

$$\dot{\mathbf{c}} = \mathbf{R}(\mathbf{c}, u) = \sum_{r=1}^R \kappa_r(u) \left( \frac{\mathbf{c}^{\boldsymbol{\alpha}^r}}{\mathbf{w}(u)^{\boldsymbol{\alpha}^r}} - \frac{\mathbf{c}^{\boldsymbol{\beta}^r}}{\mathbf{w}(u)^{\boldsymbol{\beta}^r}} \right) (\boldsymbol{\beta}^r - \boldsymbol{\alpha}^r) = \mathbb{L}(\mathbf{c}, u) D_c S(\mathbf{c}, u), \quad (2.13)$$

where the monomials  $\mathbf{c}^\gamma$  are defined via  $\prod_{i=1}^I c_i^{\gamma_i}$  and where the Onsager matrix  $\mathbb{L} \in \mathbb{R}^{I \times I}$  is a sum of rank-one contributions for each reaction, viz.

$$\mathbb{L}(\mathbf{c}, u) = \sum_{r=1}^R \kappa_r(u) \Lambda \left( \frac{\mathbf{c}^{\boldsymbol{\alpha}^r}}{\mathbf{w}(u)^{\boldsymbol{\alpha}^r}}, \frac{\mathbf{c}^{\boldsymbol{\beta}^r}}{\mathbf{w}(u)^{\boldsymbol{\beta}^r}} \right) (\boldsymbol{\alpha}^r - \boldsymbol{\beta}^r) \otimes (\boldsymbol{\alpha}^r - \boldsymbol{\beta}^r) \quad \text{with} \quad \Lambda(a, b) = \frac{a - b}{\log(a/b)}.$$

We now define the entropy production through reaction via

$$P_{\text{react}}(\mathbf{c}, u) := -\mathbf{R}(\mathbf{c}, u) \cdot (\log(c_i/w_i(u)))_i = D_c S(\mathbf{c}, u) \cdot \mathbb{L}(\mathbf{c}, u) D_c S(\mathbf{c}, u) \geq 0. \quad (2.14)$$

**Remark 2.5 (Arrhenius-type reaction-rate coefficients)** *It is also possible to consider reaction-rate coefficients  $\kappa_r > 0$  depending on  $\mathbf{c}$  and  $u$ , which leads to an implicit dependence on the temperature, e.g.  $\kappa_r(\mathbf{c}, u) = A_r \exp(-E_r \partial_u S(\mathbf{c}, u)/\mathbb{R}) = A_r \exp(\frac{-E_r}{\mathbb{R}\theta})$ , where  $E_r$  is the activation energy and  $\mathbb{R}$  the gas constant, see [BoD15, DD\*16].*

Fourth, we remark that  $\mathbb{L}$  usually has lower rank, because the stoichiometric subspace

$$\mathbb{S} := \text{span}\{\boldsymbol{\alpha}^r - \boldsymbol{\beta}^r \mid r = 1, \dots, R\} \subset \mathbb{R}^I$$

may have a dimension  $m_{\mathbb{S}} := \dim \mathbb{S}$  less than  $I$ . In that case (2.13) has natural conserved quantities (often called conservation of atomic masses) which can be described by a matrix  $\mathbf{Q} : \mathbb{R}^I \rightarrow \mathbb{R}^m$  where  $m = I - m_{\mathbb{S}}$  such that

$$\text{kernel}(\mathbf{Q}) = \mathbb{S} \quad \text{and} \quad \text{range}(\mathbf{Q}^\top) = \mathbb{S}^\perp := \{\boldsymbol{\xi} \in \mathbb{R}^I \mid \boldsymbol{\xi} \cdot \mathbf{v} = 0 \text{ for } \mathbf{v} \in \mathbb{S}\}.$$

Obviously, solutions of (2.13) satisfy  $\mathbf{Q}\mathbf{c}(t) = \mathbf{Q}\mathbf{c}(0)$  and thus stay inside the convex sets

$$\mathbf{C}_{\mathbf{q}} := \mathbf{c}(0) + \mathbb{S} \cap [0, \infty[^I, \quad \text{where } \mathbf{q} := \mathbf{Q}\mathbf{c}(0).$$

For later usage we also define  $\mathfrak{Q} \subset \mathbb{R}^m \times ]0, \infty[$  as the set of all relevant conservation values, namely

$$\mathfrak{Q} := \{(\mathbf{q}, \mathbf{U}) = (\mathbf{Q}\mathbf{c}, u) \in \mathbb{R}^{m+1} \mid (\mathbf{c}, u) \in [0, \infty[^I \times ]0, \infty[ \}.$$

Note that  $\mathbf{U} = 0$  is the case of zero energy, which implies  $u(t, x) = 0$  for all  $t$  and  $x$ , which reduces to the case treated in [DeF06, DFT16, FeT16, Mie16]. Hence, it is no restriction to assume  $\mathbf{U} > 0$  subsequently.

Finally, we remark that for each fixed  $u$  and  $\mathbf{q} = \mathbf{Q}\mathbf{c}(0)$  there exists a unique maximizer  $\mathbf{w} = \mathbf{w}(\mathbf{q}, u)$  of  $\mathcal{S}(\cdot, u)$  on  $\mathbf{C}_{\mathbf{q}}$  such that

$$\mathbf{w}(\mathbf{q}, u) := \operatorname{argmax}\left\{ S(\mathbf{c}, u) \mid \mathbf{c} \in \mathbf{C}_{\mathbf{q}} \right\} \implies \exists \eta \in \mathbb{R}^m: \left( \log \left( \frac{\mathbf{w}_i(\mathbf{q}, u)}{w_i(u)} \right) \right)_{1, \dots, I} = \mathbf{Q}^\top \eta. \quad (2.15)$$

Here  $\eta \in \mathbb{R}^m$  is the Lagrange multiplier associated with the constraint  $\mathbf{Q}\mathbf{c} = \mathbf{q}$ . Since  $\mathbf{w}(u)$  is a global maximizer of  $S(\cdot, u)$  we obviously have  $\mathbf{w}(\mathbf{q}_u, u) = \mathbf{w}(u)$  if  $\mathbf{q}_u = \mathbf{Q}\mathbf{w}(u)$ .

Since  $\mathbf{Q}(\boldsymbol{\alpha}^r - \boldsymbol{\beta}^r) = 0$  for all  $r$ , it is easy to see that each  $\mathbf{w}(\mathbf{q}, u)$  is a steady state of (2.13) and, moreover, it still satisfies the DBC (2.12), see also [MHM15, Sec. 2.1]. Thus, if necessary, we may assume that for a given initial condition we already have  $\mathbf{Q}\mathbf{c}(0) = \mathbf{Q}\mathbf{w}(u)$ .

We now return to the question of quantitative estimates for the exponential decay. Assuming that all reaction-rate coefficients  $\kappa_r(u)$  are bounded from below by  $\kappa_*$ , the reactive entropy-production  $P_{\text{react}}(\mathbf{c}, u)$  as defined in (2.14) is bounded from below by  $\kappa_*$  times the simplified entropy-production potential

$$\widehat{P}_{\text{R}}(\mathbf{c}, u) := \sum_{r=1}^I G\left(\frac{\mathbf{c}^{\boldsymbol{\alpha}^r}}{\mathbf{w}(u)^{\boldsymbol{\alpha}^r}}, \frac{\mathbf{c}^{\boldsymbol{\beta}^r}}{\mathbf{w}(u)^{\boldsymbol{\beta}^r}}\right) \text{ with } G(a, b) = (a-b)(\log a - \log b) \geq 0. \quad (2.16)$$

Our major assumption for both methods is that  $\widehat{P}_{\text{R}}(\cdot, u)$  vanishes in each stoichiometric subspace  $\mathbf{C}_{\mathbf{q}}$  exactly in one point, namely  $\mathbf{w}(\mathbf{q}, u)$ . Using the special structure of  $\mathbf{R}(\cdot, u)$  induced by the mass-action kinetic and the detailed-balance condition this is equivalent to the condition that  $\mathbf{R}(\mathbf{c}, u) = 0$  has exactly one solution in each  $\mathbf{C}_{\mathbf{q}}$ , namely  $\mathbf{w}(\mathbf{q}, u)$ . We formulate this *Unique-Equilibrium Condition* (UEC) for later reference:

$$\text{(UEC)} \quad \{ \mathbf{c} \in \mathbf{C}_{\mathbf{q}} \mid \widehat{P}_{\text{R}}(\mathbf{c}, u) \} = \{ \mathbf{w}(\mathbf{q}, u) \} = \{ \mathbf{c} \in \mathbf{C}_{\mathbf{q}} \mid \mathbf{R}(\mathbf{c}, u) = 0 \}. \quad (2.17)$$

Since it is well known that any additional solutions of  $\mathbf{R}(\mathbf{c}, u) = 0$  have to lie on the boundary of  $[0, \infty]^I$ , one easy sufficient condition for the UEC is that no reaction contains an autocatalytic species, which means that  $\alpha_i^r$  and  $\beta_i^r$  are both positive. Hence, we have the implication

$$\left( \forall i = 1, \dots, I \text{ and } r = 1, \dots, R : \alpha_i^r \beta_i^r = 0 \right) \implies \text{UEC holds.}$$

We refer to [Fei73, GIH97] for more details. Using the relative Boltzmann entropy  $\mathfrak{B}(\mathbf{c} \mid \mathbf{w})$  defined in (1.2) the UEC allows us to show that

$$\forall (\mathbf{q}, \mathbf{U}) \in \mathfrak{Q} \exists K_{\text{R}}(\mathbf{q}, \mathbf{U}) > 0 \forall \mathbf{c} \in \mathbf{C}_{\mathbf{q}} : \widehat{P}_{\text{R}}(\mathbf{c}, \mathbf{U}) \geq K_{\text{R}}(\mathbf{q}, \mathbf{U}) \mathfrak{B}(\mathbf{c} \mid \mathbf{w}(\mathbf{q}, \mathbf{U})). \quad (2.18)$$

Indeed, this follows from a compactness argument for  $\mathbf{c} \in \mathbf{C}_{\mathbf{Q}\mathbf{w}(u)}$ , which is a special case of the result in Section 4.

## 2.4 Combining entropy production from diffusion and reaction

We return to the full ERDS

$$\begin{pmatrix} \dot{\mathbf{c}} \\ \dot{u} \end{pmatrix} = \operatorname{div} \left( M(\mathbf{c}, u) \nabla \begin{pmatrix} \mathbf{c} \\ u \end{pmatrix} \right) + \begin{pmatrix} \mathbf{R}(\mathbf{c}, u) \\ 0 \end{pmatrix} \text{ in } \Omega, \quad \nu \cdot \nabla \begin{pmatrix} \mathbf{c} \\ u \end{pmatrix} = 0 \text{ on } \partial\Omega, \quad (2.19)$$

and consider sufficiently smooth solutions  $(\mathbf{c}(t, \cdot), u(t, \cdot)) : \Omega \rightarrow [0, \infty]^{I+1}$ . From the no-flux boundary conditions on  $\partial\Omega$  and the fact  $\mathbf{R}(\mathbf{c}, u) \in \mathbb{S}$  we deduce the conservation relations

$$\int_{\Omega} u(t, x) \, dx = \int_{\Omega} u(0, x) \, dx =: \mathbf{U} \quad \text{and} \quad \int_{\Omega} \mathbf{Q}\mathbf{c}(t, x) \, dx = \int_{\Omega} \mathbf{Q}\mathbf{c}(0, x) \, dx =: \mathbf{q}.$$

Thus, the solutions  $(\mathbf{c}, u)$  will stay inside the sets

$$\mathfrak{S}(\mathbf{q}, \mathbf{U}) := \left\{ (\mathbf{c}, u) \in L^1(\Omega)^{I+1} \mid u, c_i \geq 0, \int_{\Omega} u \, dx = \mathbf{U}, \int_{\Omega} \mathbf{Q}\mathbf{c} \, dx = \mathbf{q} \right\}. \quad (2.20)$$

Clearly,  $\mathfrak{S}(\mathbf{q}, \mathbf{U})$  contains exactly one constant state, namely  $(\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$ .

For a fixed  $(\mathbf{q}, \mathbf{U})$  and using the concave entropy density  $S$  we define the nonnegative and convex relative entropy density

$$H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) := S(\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U}) + DS(\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U}) \cdot (\mathbf{c} - \mathbf{w}(\mathbf{q}, \mathbf{U}), u - \mathbf{U}) - S(\mathbf{c}, u).$$

Similarly, for entropies  $s(\cdot)$  only dependent on the internal energy  $u$ , we define

$$h_{\mathbf{U}}(u) = s(\mathbf{U}) + s'(U) \cdot (u - \mathbf{U}) - s(u) \quad \text{and} \quad \widehat{h}_{\mathbf{U}}(u) = h_{\mathbf{U}}(u) + I - \sum_{i=1}^I w_i(u). \quad (2.21)$$

Clearly,  $h_{\mathbf{U}}(u) \geq 0$  and  $H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) \geq 0$  with equality if and only if  $(\mathbf{c}, u) = (\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$ .

We define a relative entropy functional on  $L^1(\Omega)^{I+1}$  via

$$\mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) := \int_{\Omega} H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}(x), u(x)) \, dx$$

which is convex and non-negative. The unique minimizer is the constant function  $(\mathbf{c}, u) \equiv (\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$  which lies in the convex set  $\mathfrak{S}(\mathbf{q}, \mathbf{U})$ . Moreover, by construction we know that

$$\forall (\mathbf{c}, u) \in \mathfrak{S}(\mathbf{q}, \mathbf{U}) : \quad \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) = S(\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U}) - S(\mathbf{c}, u).$$

Thus, for solutions  $(\mathbf{c}(t), u(t))$  in  $\mathfrak{S}(\mathbf{q}, \mathbf{U})$  the entropy production can be calculated via

$$\frac{d}{dt} S(\mathbf{c}, u) = -\frac{d}{dt} \mathcal{H}_{\mathbf{q}}(\mathbf{c}, u) = \mathcal{P}(\mathbf{c}, u) = \mathcal{P}_{\text{diff}}(\mathbf{c}, u) + \mathcal{P}_{\text{react}}(\mathbf{c}, u) \quad (2.22)$$

$$\text{where } \mathcal{P}_{\text{react}}(\mathbf{c}, u) := \int_{\Omega} P_{\text{react}}(\mathbf{c}(x), u(x)) \, dx. \quad (2.23)$$

Under the assumptions of Sections 2.2 and 2.3 we know that  $\mathcal{P}_{\text{diff}}(\mathbf{c}, u) = 0$  if and only if  $(\mathbf{c}, u) = (\bar{\mathbf{c}}, \bar{u})$ , i.e. all functions are constant. In contrast the condition  $\mathcal{P}_{\text{react}}(\mathbf{c}, u) = 0$  implies that  $(\mathbf{c}, u)$  are in local equilibrium only, i.e.  $(\mathbf{c}(x), u(x)) = (\mathbf{w}(\tilde{\mathbf{q}}(x), \tilde{u}(x)), \tilde{u}(x))$  for arbitrary functions  $(\tilde{\mathbf{q}}, \tilde{u}) : \Omega \rightarrow \mathbb{R}^m \times [0, \infty[$ . However, combining these two conditions and the property  $(\mathbf{c}, u) \in \mathfrak{S}(\mathbf{q}, \mathbf{U})$ , we see that the entropy production  $\mathcal{P}(\mathbf{c}, u)$  restricted to  $\mathfrak{S}(\mathbf{q}, \mathbf{U})$  is 0 if and only if  $(\mathbf{c}, u) = (\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$ .

Thus, it is natural to ask whether there is a EEP estimate of the form

$$\forall (\mathbf{q}, \mathbf{U}) \in \mathfrak{Q} \exists K(\mathbf{q}, \mathbf{U}) > 0 \forall (\mathbf{c}, u) \in \mathfrak{S}(\mathbf{q}, \mathbf{U}) : \quad \mathcal{P}(\mathbf{c}, u) \geq K(\mathbf{q}, \mathbf{U}) \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u). \quad (2.24)$$

To establish the above estimate it is convenient to estimate the two terms in  $\mathcal{P}$  by its model contributions, namely

$$\begin{aligned} \mathcal{P}_{\text{diff}}(\mathbf{c}, u) &\geq \delta_* \widehat{\mathcal{P}}_{\gamma}(\mathbf{c}, u) \quad \text{and} \quad \mathcal{P}_{\text{react}}(\mathbf{c}, u) \geq \kappa_* \widehat{\mathcal{P}}_{\text{R}}(\mathbf{c}, u), \\ \text{where } \widehat{\mathcal{P}}_{\text{R}}(\mathbf{c}, u) &:= \int_{\Omega} \widehat{P}_{\text{R}}(\mathbf{c}(x), u(x)) \, dx \end{aligned} \quad (2.25)$$

with  $\widehat{\mathcal{P}}_{\gamma}$  and  $\widehat{P}_{\text{R}}$  from (2.7) and (2.16), respectively. In the sequel we will hence simplify the analysis and the notation by only estimating  $\widehat{\mathcal{P}}_{\gamma}(\mathbf{c}, u) + \widehat{\mathcal{P}}_{\text{R}}(\mathbf{c}, u)$  from below by  $\mathcal{H}_{\mathbf{q}, \mathbf{U}}$ ,

and moreover by always assuming that  $\mathbf{q} = \mathbf{Q}\mathbf{w}(\mathbf{U})$ . As was explained in Section 2.3 this is always possible by adjusting  $\mathbf{w}$  suitably. However, we need to be sure that this does not change the necessary structural properties of the vector-valued function  $u \mapsto \mathbf{w}(u)$ , such as monotonicity and concavity, see Section 2.1. However, when fixing on  $\mathbf{U}$  and  $\mathbf{q}$  we can change  $\mathbf{w}(\cdot)$  by pre-multiplying it with the diagonal matrix  $A = \text{diag}(e^{a_1}, \dots, e^{a_I})$ , where  $\mathbf{a} = \mathbf{Q}^\top \boldsymbol{\eta}$  is the vector given in (2.15). Then, obviously the new vector-valued function  $\tilde{\mathbf{w}}(u) = A\mathbf{w}(u)$  satisfies  $\tilde{\mathbf{w}}(\mathbf{U}) = \mathbf{w}(\mathbf{q}, \mathbf{U})$ . However, such a pre-multiplication does not change any of the properties we assumed so far or we will use later. Of course, this will change all the constants, in particular  $K_R(\mathbf{U}, q)$ .

Thus, with a slight abuse of notation we will use the notation

$$\widehat{\mathfrak{G}}(\mathbf{U}) := \left\{ (\mathbf{c}, u) \in L^1(\Omega)^{I+1} \mid u, c_i \geq 0, \int_{\Omega} u \, dx = \mathbf{U}, \int_{\Omega} \mathbf{Q}\mathbf{c} \, dx = \mathbf{Q}\mathbf{w}(\mathbf{U}) \right\},$$

which leads us to our remaining task, which is to show the following estimate:

$$\forall \mathbf{U} > 0 \exists \widehat{K}(\mathbf{U}) > 0 \forall (\mathbf{c}, u) \in \widehat{\mathfrak{G}}(\mathbf{U}) : \quad (\widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) + \widehat{\mathcal{P}}_R(\mathbf{c}, u)) \geq \widehat{K}(\mathbf{U}) \widehat{\mathcal{H}}_{\mathbf{U}}(\mathbf{c}, u), \quad (2.26)$$

where  $\widehat{H}_{\mathbf{U}} = H_{\mathbf{Q}\mathbf{w}(\mathbf{U}), \mathbf{U}}$  and  $\widehat{\mathcal{H}}_{\mathbf{U}}(\mathbf{c}, u) = \int_{\Omega} \widehat{H}_{\mathbf{U}}(\mathbf{c}, u) \, dx = \mathcal{H}_{\mathbf{Q}\mathbf{w}(\mathbf{U}), \mathbf{U}}(\mathbf{c}, u)$ . Clearly, if (2.26) holds, then we obtain the bound  $K(\mathbf{Q}\mathbf{w}(\mathbf{U}), \mathbf{U}) \geq \min\{\delta_*, \kappa_*\} \widehat{K}(\mathbf{U})$  in (2.24).

### 3 EEP estimates via convexity

In this section we describe the general method on how to obtain estimates of the entropy in terms of the entropy-production by using a convexity argument in the way as derived in [MHM15] for the isothermal case. It is based on the fact that the diffusion part  $\widehat{\mathcal{P}}_{\text{diff}}$  can be estimated by the log-Sobolev inequality and variants of it, see Section 3.1. If this method works it provides stronger results that are not dependent on the entropy of the initial condition as in the case of the compactness method from [GIH97] discussed in Section 4.

#### 3.1 Generalized log-Sobolev inequalities

For  $\alpha \geq 0$  we consider the entropy functions

$$F_\alpha(z) = \begin{cases} \frac{1}{\alpha(\alpha-1)}(z^\alpha - \alpha z + \alpha - 1) & \text{for } \alpha \in \mathbb{R} \setminus \{0, 1\}, \\ \lambda_B(z) = z \log z - z + 1 & \text{for } \alpha = 1, \\ z - \log z - 1 & \text{for } \alpha = 0. \end{cases} \quad (3.1)$$

We see that all  $F_\alpha$  are convex and satisfy  $F_\alpha(1) = F'_\alpha(1) = 0 \leq F_\alpha(z)$ . In particular, we have  $F''_\alpha(z) = z^{\alpha-2} > 0$ . Moreover, for  $\alpha > 0$  we have the monotonicities

$$\partial_\alpha(\alpha F_\alpha(z)) = \frac{z}{(1-\alpha)^2} F_1(z^{\alpha-1}) \geq 0 \quad \text{and} \quad \partial_\alpha((1-\alpha)F_\alpha(z)) = -\frac{1}{\alpha^2} F_1(z^\alpha) \leq 0. \quad (3.2)$$

Thus, for all  $\alpha \in ]0, 1[$  all these entropies are equivalent:

$$0 < \alpha_1 < \alpha_2 < 1 \quad \implies \quad \frac{1-\alpha_2}{1-\alpha_1} F_{\alpha_2}(z) \leq F_{\alpha_1}(z) \leq \frac{\alpha_2}{\alpha_1} F_{\alpha_2}(z).$$

Moreover, we have  $F_1 = \lambda_B$  and the following identities

$$F_1(uv) = vF_1(u) + uF_1(v) + (u-1)(v-1), \quad (3.3)$$

$$a^\alpha F_\alpha(u/a) = F_\alpha(u) + \frac{a^\alpha - 1}{\alpha} - u \frac{a^{\alpha-1} - 1}{\alpha - 1}. \quad (3.4)$$

Using  $\bar{u} = \int_\Omega u(x) dx$  an integration of (3.4) gives

$$\bar{u}^\alpha \int_\Omega F_\alpha(u/\bar{u}) dx = \int_\Omega F_\alpha(u) dx - F_\alpha(\bar{u}) \quad (\text{recall } |\Omega| = 1). \quad (3.5)$$

Crucial for the forthcoming analysis will be the following estimate, which can be seen as a generalization of Poincaré and the log-Sobolev estimates, namely

$$\forall u > 0 \text{ with } u^{\gamma/2} \in H^1(\Omega) : \quad \int_\Omega \frac{|\nabla u|^2}{u^{2-\gamma}} dx \geq \rho(\Omega, \gamma, \alpha) \bar{u}^\gamma \int_\Omega F_\alpha(u/\bar{u}) dx, \quad (3.6)$$

where  $\rho(\Omega, \gamma, \alpha)$  is supposed to be the optimal constant. The monotonicity (3.2) implies

$$0 < \alpha_1 < \alpha_2 \quad \implies \quad \rho(\Omega, \gamma, \alpha_1) \leq \frac{\alpha_1}{\alpha_2} \rho(\Omega, \gamma, \alpha_2).$$

If  $2_d$  is the Lebesgue power for the embedding  $H^1(\Omega) \subset L^{2_d}(\Omega)$  for a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^d$  (namely  $2_d = 2d/(d-2)$  for  $d \geq 3$ ), then  $\rho(\Omega, \gamma, \alpha)$  is positive if  $\gamma 2_d \geq 2$  and  $\alpha \leq \gamma + 2/d$ , see Appendix A. The case  $\alpha = \gamma = 2$  is the *Poincaré estimate*, while  $\alpha = \gamma = 1$  is the *log-Sobolev estimate*, which we will be crucial for the concentrations  $c_i$ .

However, for the internal energy, we want to use an entropy function  $F_\alpha$  with  $\alpha = \gamma \in ]0, 1[$ . Since the entropy functions are equivalent for  $\alpha \in ]0, 1[$  and have a linear growth, a positive constant  $\rho(\Omega, \gamma, \alpha)$  can only exist if  $2_d \gamma \geq 2$ . Subsequently we will often use  $\rho(\Omega, \gamma, 1) > 0$  which certainly holds for  $\gamma 2_d > 2$ . Indeed this is a special and simple subcase of the compactness arguments developed in Section 4.

Considering the case of gas dynamics, where the entropy  $s(u) = s_0 \log u$  is relevant, it is interesting to note that the case  $\alpha = \gamma = 0$  is still possible, but only in dimensions  $d \leq 2$ , see Appendix B.

### 3.2 The general setup of the convexity method

We now want to develop ideas to establish estimate (2.26) for fixed  $U > 0$  and  $(\mathbf{c}, u) \in \widehat{\mathfrak{G}}(U)$ . The main point is that the diffusion part  $\widehat{\mathcal{P}}_\gamma$  of the entropy production has a lower bound in terms of functionals that only depend on the point values  $(\mathbf{c}(x), u(x))$  and no derivatives. These functionals then need to be combined with  $\widehat{\mathcal{P}}_\mathbb{R}$  to find an upper bound for  $\widehat{\mathcal{H}}_U$ . Thus, all the latter functionals only involve point values  $(\mathbf{c}(x), u(x))$  and no derivatives, so it is natural to seek for pointwise estimates of the integrands. However, in general this cannot work, since we need to involve the constraints arising from  $(\mathbf{c}, u) \in \widehat{\mathfrak{G}}(U)$ , namely  $\mathbf{Q}\bar{\mathbf{c}} = \mathbf{Q}\mathbf{w}(U)$  and  $\bar{u} = U$ , and to exploit the ODE decay coefficient  $\widehat{K}_\mathbb{R}(U) := K_\mathbb{R}(\mathbf{Q}\mathbf{w}(U), U)$  from (2.18).

For the densities  $c_i$  we use the classical log-Sobolev estimate with  $\rho_\Omega^{\text{IS}} = \rho(\Omega, 1, 1)$ :

$$\int_\Omega \frac{|\nabla c_i|^2}{c_i} dx \geq \rho_\Omega^{\text{IS}} \mathfrak{B}(c_i | \bar{c}_i) = \rho_\Omega^{\text{IS}} \int_\Omega \bar{c}_i \lambda_B(c_i / \bar{c}_i) dx = \rho_\Omega^{\text{IS}} \left( \int_\Omega \lambda_B(c_i) dx - \lambda_B(\bar{c}_i) \right).$$

For the internal energy the necessary estimate in the general case is  $\int_\Omega \widehat{h}_U''(u) |\nabla u|^2 dx \geq C_* \int_\Omega h_U(u) dx$  for all  $u$  with  $\bar{u} = U$ . Recall the definition of  $h_U$  and  $\widehat{h}_U$  in (2.21). The assumption in Section 2.2 already imposed  $\widehat{h}_U''(u) \geq \delta/u^{2-\gamma}$  for some  $\gamma \in ]0, 1[$ . Hence,

setting  $\rho_\Omega^{\gamma,\alpha} := \rho(\Omega, \gamma, \alpha) > 0$  (cf. (3.6)) for a suitable  $\alpha \geq \gamma$  we can use the following EEP estimate for the internal energy:

$$\forall u \text{ with } \bar{u} = \mathbf{U} : \int_\Omega \frac{|\nabla u|^2}{u^{2-\gamma}} dx \geq \rho_\Omega^{\gamma,\alpha} \mathbf{U}^\gamma \int_\Omega F_\alpha(u/\mathbf{U}) dx. \quad (3.7)$$

For controlling the deviation of  $u$  from  $\bar{u}$ , it would be sufficient to use  $\alpha = \gamma$ ; however, we will use  $F_\alpha$  with  $\alpha \geq 1$ , because a higher value for  $\alpha$  is advantageous for the subsequent convexity condition. We further assume

$$\forall \mathbf{U} \exists C_h(\mathbf{U}) > 0 \forall u > 0 : \mathbf{U}^\gamma F_\alpha(u/\mathbf{U}) \geq C_h(\mathbf{U}) h_{\mathbf{U}}(u). \quad (3.8)$$

For the case  $s(u) = s_0 u^\gamma$  we have  $h_{\mathbf{U}}(u) = s_0 \gamma (1-\gamma) \mathbf{U}^\gamma F_\gamma(u/\mathbf{U})$ , i.e.  $C_h(\mathbf{U}) = 1/(s_0 \gamma (1-\gamma))$  can be chosen independently of  $\mathbf{U}$ .

Combining this with the estimates for the densities we obtain the desired lower bound for  $\widehat{\mathcal{P}}_\gamma$  in terms of derivative-free functionals, namely

$$\widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) \geq p_\Omega \int_\Omega \left( \mathbf{U}^\gamma F_\alpha(u(x)/\mathbf{U}) + \mathfrak{B}(\mathbf{c}(x)|\mathbf{1}) - \mathfrak{B}(\bar{\mathbf{c}}|\mathbf{1}) \right) dx, \quad (3.9)$$

where  $\bar{u} = \mathbf{U}$ ,  $p_\Omega = \min\{\rho_\Omega^{\text{IS}}, \rho_\Omega^{\gamma,\alpha}\}$ , and  $\mathbf{1} = (1, 1, \dots, 1)$ .

Since the desired relative entropy reads  $\widehat{H}_{\mathbf{U}}(\mathbf{c}, u) = h_{\mathbf{U}}(u) + \mathfrak{B}(\mathbf{c}|\mathbf{w}(u))$ , but the ODE decay rate  $\widehat{K}_{\mathbf{R}}(\mathbf{U}) = K_{\mathbf{R}}(\mathbf{Q}\mathbf{w}(\mathbf{U}), \mathbf{U})$  involves the relative Boltzmann entropy  $\mathfrak{B}(\mathbf{c}|\mathbf{w}(\mathbf{U}))$  with respect to the fixed value  $\mathbf{w}(\mathbf{U})$ , we need the following useful lemma.

**Lemma 3.1 (Comparing relative entropies)** *Assume that the functions  $w_i : [0, \infty[ \rightarrow [0, \infty[$  are increasing, concave, and satisfy  $w_i(0) > 0$  and that (3.8) holds, then there exists a constant  $C_H(\mathbf{U})$  such that*

$$\forall (\mathbf{c}, u) \in [0, \infty[^{I+1} : \widehat{H}_{\mathbf{U}}(\mathbf{c}, u) = h_{\mathbf{U}}(u) + \mathfrak{B}(\mathbf{c}|\mathbf{w}(u)) \leq C_H(\mathbf{U}) (\mathbf{U}^\gamma F_\alpha(u/\mathbf{U}) + \mathfrak{B}(\mathbf{c}|\mathbf{w}(\mathbf{U}))).$$

**Proof:** Using (3.8) it suffices to show the assertion with  $\mathbf{U}^\gamma F_\alpha(u/\mathbf{U})$  replaced by  $h_{\mathbf{U}}(u)$ . Hence, we define  $\widetilde{H}(\mathbf{c}, u) = h_{\mathbf{U}}(u) + \mathfrak{B}(\mathbf{c}|\mathbf{w}(\mathbf{U}))$  and see that  $\widetilde{H}$  and  $\widehat{H}_{\mathbf{U}}$  are nonnegative and vanish only at  $(\mathbf{c}, u) = (\mathbf{w}(\mathbf{U}), \mathbf{U})$ . Since both functions behave quadratically near this point, the upper estimates follows by compactness for all bounded domains.

For large  $(\mathbf{c}, u)$  we have  $\widetilde{H}(\mathbf{c}, u) \geq \mathfrak{B}(\mathbf{c}|\mathbf{1}) + \frac{1}{C}(u + |\mathbf{c}|_1) - C$  and  $\widehat{H}_{\mathbf{U}}(\mathbf{c}, u) \leq \mathfrak{B}(\mathbf{c}|\mathbf{1}) + C(1+u+\mu|\mathbf{c}|_1+|\mathbf{w}(u)|_1)$ , where  $\mu = -\min_i \log w_i(0)$ . Since  $|\mathbf{w}(u)|_1 \leq C(1+u)$  the desired result follows also for large  $(\mathbf{c}, u)$ .  $\blacksquare$

Now we are able to state our major result for the convexity method. The resulting decay rate  $\widetilde{K}(\mathbf{U})$  is dominated by  $\kappa_* \widehat{K}_{\mathbf{R}}(\mathbf{U})$ , which is the pure reaction decay and by  $\delta_* p_\Omega$ , which is the decay rate for pure diffusion. However, as in the isothermal case, see [MHM15, Thm. 3.1], the latter coefficient is multiplied by a factor  $\eta = \widehat{K}_{\mathbf{R}}(\mathbf{U})/(\mu(\mathbf{U}) + \widehat{K}_{\mathbf{R}}(\mathbf{U})) \leq 1$  that stands for the interplay of reaction and diffusion, since  $\mu(\mathbf{U})$  measures how much convexity induced by diffusion is needed to convexify the typically non-convex reactive entropy production  $\widehat{P}_{\mathbf{R}}$ .

**Theorem 3.2 (Convexity method)** *Consider  $S(\mathbf{c}, u) = s(u) - \mathfrak{B}(\mathbf{c}|\mathbf{w}(u))$  and let  $h_{\mathbf{U}}$  be defined as above, satisfying  $p_\Omega > 0$  in (3.9),  $C_H(\mathbf{U}) > 0$  in Lemma 3.1, and the unique equilibrium condition (2.17), i.e.  $\widehat{K}_{\mathbf{R}}(\mathbf{U}) > 0$  in (2.18). We define the function*

$$G_{\mu, \mathbf{U}}(\mathbf{c}, u) := \mu (\mathbf{U}^\gamma F_\alpha(u/\mathbf{U}) + \mathfrak{B}(\mathbf{c}|\mathbf{1})) + \widehat{P}_{\mathbf{R}}(\mathbf{c}, u)$$

and assume the following convexity condition:

$$\forall \mathbf{U} \exists \mu(\mathbf{U}) \geq 0 : \quad G_{\mu(\mathbf{U}), \mathbf{U}} : [0, \infty[^{I+1} \rightarrow [0, \infty[ \text{ is convex.} \quad (3.10)$$

Then, we have the following EEP estimate

$$\begin{aligned} \forall (\mathbf{c}, u) \in \widehat{\mathfrak{S}}(\mathbf{U}) : \quad & \delta_* \widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) + \kappa_* \widehat{\mathcal{P}}_R(\mathbf{c}, u) \geq \widetilde{K}(\mathbf{U}) \widehat{\mathcal{H}}_{\mathbf{U}}(\mathbf{c}, u) \\ \text{with } \widetilde{K}(\mathbf{U}) = & \frac{1}{C_H(\mathbf{U})} \min \left\{ \kappa_* \widehat{K}_R(\mathbf{U}), \delta_* p_\Omega \frac{\widehat{K}_R(\mathbf{U})}{\mu(\mathbf{U}) + \widehat{K}_R(\mathbf{U})} \right\}. \end{aligned} \quad (3.11)$$

**Proof:** Defining  $H_B(\mathbf{c}, u) = U^\gamma F_\alpha(u/\mathbf{U}) + \mathfrak{B}(\mathbf{c}|\mathbf{1}) - \mathfrak{B}(\bar{\mathbf{c}}|\mathbf{1})$ , choosing  $\theta \in ]0, 1[$ , and setting  $M(\theta, \mathbf{U}) = \min\{\kappa_*, (1-\theta)\delta_* p_\Omega/\mu(\mathbf{U})\}$  the assumptions yield the estimate

$$\begin{aligned} \delta_* \widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) + \kappa_* \widehat{\mathcal{P}}_R(\mathbf{c}, u) & \geq \int_\Omega \left( \delta_* p_\Omega H_B(\mathbf{c}, u) + \kappa_* \widehat{P}_R(\mathbf{c}, u) \right) dx \\ & \geq \theta \delta_* p_\Omega \int_\Omega H_B(\mathbf{c}, u) dx + M(\theta, \mathbf{U}) \int_\Omega (G_{\mu(\mathbf{U}), \mathbf{U}}(\mathbf{c}, u) - \mu(\mathbf{U}) \mathfrak{B}(\bar{\mathbf{c}}|\mathbf{1})) dx \\ & \stackrel{\text{Jensen}}{\underset{(3.10)}{\geq}} \theta \delta_* p_\Omega \int_\Omega H_B(\mathbf{c}, u) dx + M(\theta, \mathbf{U}) (G_{\mu(\mathbf{U}), \mathbf{U}}(\bar{\mathbf{c}}, \bar{u}) - \mu(\mathbf{U}) \mathfrak{B}(\bar{\mathbf{c}}|\mathbf{1})) \\ & = \theta \delta_* p_\Omega \int_\Omega H_B(\mathbf{c}, u) dx + M(\theta, \mathbf{U}) (\mu(\mathbf{U}) 0 + \widehat{P}_R(\bar{\mathbf{c}}, \mathbf{U})) \end{aligned}$$

where we used  $\bar{u} = \mathbf{U}$  and  $F_\alpha(\bar{u}/\mathbf{U}) = 0$  in the last identity. Employing the ODE estimate (2.18), where we use  $\mathbf{Q}\bar{\mathbf{c}} = \mathbf{Q}\mathbf{w}(\mathbf{U})$  following from  $(\mathbf{c}, u) \in \widehat{\mathfrak{S}}(\mathbf{U})$ , we continue

$$\begin{aligned} \delta_* \widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) + \kappa_* \widehat{\mathcal{P}}_R(\mathbf{c}, u) & \geq \theta \delta_* p_\Omega \int_\Omega H_B(\mathbf{c}, u) dx + M(\theta, \mathbf{U}) \widehat{K}_R(\mathbf{U}) \mathfrak{B}(\bar{\mathbf{c}}|\mathbf{w}(\mathbf{U})) \\ & \geq N(\theta, \mathbf{U}) \int_\Omega (U^\gamma F_\alpha(u/\mathbf{U}) + \mathfrak{B}(\mathbf{c}|\mathbf{1}) - \mathfrak{B}(\bar{\mathbf{c}}|\mathbf{1}) + \mathfrak{B}(\bar{\mathbf{c}}|\mathbf{w}(\mathbf{U}))) dx \\ & =_{(1)} N(\theta, \mathbf{U}) \int_\Omega (U^\gamma F_\alpha(u/\mathbf{U}) + \mathfrak{B}(\mathbf{c}|\mathbf{w}(\mathbf{U}))) dx \geq_{(2)} \frac{N(\theta, \mathbf{U})}{C_H(\mathbf{U})} \int_\Omega (h_{\mathbf{U}}(u) + \mathfrak{B}(\mathbf{c}|\mathbf{w}(u))) dx. \end{aligned}$$

where  $N(\theta, \mathbf{U}) = \min\{\theta \delta_* p_\Omega, M(\theta, \mathbf{U}) \widehat{K}_R(\mathbf{U})\}$ . For “ $=_{(*)}$ ” we used the general identity

$$\mathfrak{B}(\mathbf{c}|\mathbf{1}) - \mathfrak{B}(\mathbf{b}|\mathbf{1}) + \mathfrak{B}(\mathbf{b}|\tilde{\mathbf{w}}) = \mathfrak{B}(\mathbf{c}|\tilde{\mathbf{w}}) + \sum_{i=1}^I (c_i - b_i) \log \tilde{w}_i$$

for  $\mathbf{b} = \bar{\mathbf{c}}$ . Since  $\tilde{\mathbf{w}} = \mathbf{w}(\mathbf{U})$  is spatially constant, the sum on the right-hand side cancels upon integration over  $\Omega$ . For the estimate “ $\geq_{(2)}$ ” we used Lemma 3.1, and the result is established if we choose the optimal  $\theta = \widehat{K}_R/(\mu + \widehat{K}_R)$ .  $\blacksquare$

In the above theorem we have given only the simplest form of the convexity method. As is shown in [MHM15], there are several ways to generalize the method (i) by replacing the convexity assumption by an estimate on the convexification (cf. [MHM15, Sec. 3.3]) or (ii) by using higher entropies  $F_\alpha$  with  $\alpha > 1$  instead of  $F_1 = \lambda_B$  for the densities  $c_i$  (cf. [MHM15, Sec. 3.4]).

Note that we already used the higher-order entropy  $F_\alpha$  for the internal energy, since  $F_\gamma$  and  $h_{\mathbf{U}}$  only have linear growth, which would gives less strict convexity to be exploited for compensating the missing convexity of  $\widehat{P}_R$ . Indeed, we will see in Section 3.3.3 that  $\alpha = 1 > \gamma = 1/2$  is useful. Thus, it is crucial that we have  $\rho(\Omega, \gamma, \alpha) > 0$  for some  $\alpha \geq 1$ ,

which is a well-known fact in finite space dimensions. The optimal constant  $\rho(\Omega, \gamma, \alpha)$  is positive for  $\alpha \in ]0, \gamma+2/d]$  if  $\gamma \in ]0, 1[$  and  $d \leq 2$  and for  $\alpha \in ]0, \gamma+2/d]$  if  $\gamma \in ]1-2/d, 1[$  and  $d \geq 3$ , see e.g. [Mit14] and Appendix A. In particular,  $\alpha = 1$  is always admissible.

In the following we will apply the method to a simple example and leave a general study for future work.

### 3.3 A simple ERDS with scalar $c$

To give a first impression of the convexity method we consider one chemical species with density  $c > 0$  that can react with the background by absorption or generation. This reaction is steered by the local value of the internal energy  $u(t, x)$ .

#### 3.3.1 General description of the simple ERDS

More precisely, we consider the semilinear system

$$\dot{c} = \delta_u \Delta c + \kappa(c, u)(f(w(u)) - f(c)), \quad \dot{u} = \delta_u \Delta u, \quad (3.12)$$

on a bounded Lipschitz domain with no-flux boundary conditions. Here  $w(u)$  is a given function determining the equilibrium concentration  $c = w(u)$  for a given internal energy. The function  $f : [0, \infty[ \rightarrow [0, \infty[$  is assumed to be smooth, strictly increasing and satisfying  $f(0) = 0$ . The positive reaction-rate coefficient  $\kappa = \kappa(c, u)$  gives the strength of the reaction.

We now redo the above estimate on this concrete example and try to make the constants as explicit as possible. For a fixed  $\mathbf{U}$  and an entropy density  $S(c, u) = s(u) - w(u)\lambda_B(c/w(u))$  we construct the relative entropy via

$$\begin{aligned} \mathcal{H}_{\mathbf{U}}(c, u) &= \int_{\Omega} H_{\mathbf{U}}(c(x), u(x)) \, dx, \quad \text{where} \\ H_{\mathbf{U}}(c, u) &= h_{\mathbf{U}}(u) + w(u)\lambda_B\left(\frac{c}{w(u)}\right) \quad \text{with } h_{\mathbf{U}}(u) = s(\mathbf{U}) - s'(u)(u - \mathbf{U}) - s(u). \end{aligned} \quad (3.13)$$

The entropy production terms read

$$\mathcal{P}(c, u) = \delta_c \mathcal{P}_c(c, u) + \delta_u \mathcal{P}_u(c, u) + \mathcal{P}_R(c, u) \quad \text{with} \quad (3.14a)$$

$$\mathcal{P}_c(c, u) = \int_{\Omega} \frac{1}{c} |\nabla c|^2 - \frac{w'(u)}{w(u)} \nabla c \cdot \nabla u \, dx, \quad (3.14b)$$

$$\mathcal{P}_u(c, u) = \int_{\Omega} -\frac{w'(u)}{w(u)} \nabla c \cdot \nabla u + \left( c \frac{w'^2 - w w''}{w^2} + \widehat{h}'' \right) |\nabla u|^2 \, dx, \quad (3.14c)$$

$$\begin{aligned} \mathcal{P}_R(c, u) &= \int_{\Omega} P_R(c(x), u(x)) \, dx \\ &\quad \text{with } P_R(c, u) = \kappa(c, u)(f(c) - f(w(u))) \log\left(\frac{c}{w(u)}\right). \end{aligned} \quad (3.14d)$$

Obviously, we have  $P_R(c, u) \geq 0$ , but  $\mathcal{P}_c$  and  $\mathcal{P}_u$  are only non-negative individually if  $w'(u) \equiv 0$ , which is a manifestation of cross-diffusion.

Proceeding as in Proposition 2.1 we assume that there exists  $\rho \in ]0, 1[$  such that

$$\rho < \frac{4\delta_c \delta_u}{(\delta_c + \delta_u)^2} \leq 1 \quad \text{and} \quad \forall u > 0 : \quad -w(u)w''(u) \geq \frac{1-\rho}{\rho} w'(u)^2. \quad (3.15)$$

Then, the sum is non-negative and has the lower bound

$$\delta_c \mathcal{P}_c(c, u) + \delta_u \mathcal{P}_u(c, u) \geq \int_{\Omega} \left( \theta \delta_c^{\text{red}} \frac{|\nabla c|^2}{c} + \delta_u \widehat{h}_U''(u) |\nabla u|^2 \right) dx,$$

where the reduced diffusion coefficient reads  $\delta_c^{\text{red}} = \delta_c (1 - \rho(\delta_c + \delta_u)^2 / (4\delta_c \delta_u)) > 0$ , and  $\widehat{h}_U(u) = h_U(u) + w(u)$  is as above, e.g. convex.

### 3.3.2 A first simplistic EEP estimate

A very simplistic first result is the following, which assumes that the reactive entropy production  $P_R(c, u)$  dominates the relative entropy  $w(u)\lambda_B(c/w(u))$  for all  $(c, u)$ , which essentially means that the ODE  $\dot{c} = R(c, U)$  has a uniform exponential decay towards the equilibrium  $w(U)$ , independently of  $U$ .

**Proposition 3.3 (Simple decay result)** *If for all  $U > 0$  there exist positive constants  $k_D(U)$  and  $k_R$  such that*

$$\begin{aligned} \forall u \in L^1_{\geq 0}(\Omega), \bar{u} = U : \quad & \int_{\Omega} \widehat{h}''(u) |\nabla u|^2 dx \geq k_D(U) \int_{\Omega} h_U(u) dx, \\ \forall (c, u) \in ]0, \infty[^2 : \quad & P_R(c, u) \geq k_R w(u) \lambda_B(c/w(u)), \end{aligned}$$

then we have the EEP estimate

$$\forall (c, u) \in \widehat{\mathfrak{S}}(U) : \mathcal{P}(c, u) \geq \min\{\delta_u k_D(U), k_R\} \mathcal{H}_U(c, u).$$

Obviously, this result follows simply by estimating  $\int_{\Omega} w(u)\lambda_B(c/w(u)) dx$  by  $\mathcal{P}_R$  and  $\int_{\Omega} h_U(u) dx$  by  $\mathcal{P}_u$  separately, which only works in very special cases.

The result can be applied to the ERDS (3.12) for the choices

$$\begin{aligned} w(u) &= u^\gamma, \quad s(u) = s_0 u^\gamma \text{ with } s_0 > 1 > \gamma > 0, \\ f(c) &= c^\beta \text{ with } \beta \geq 1, \quad \text{and } \kappa(c, u) = \kappa_* u^{\gamma(1-\beta)}. \end{aligned} \tag{3.16}$$

Then, setting  $c = zw(u)$ , we have

$$\frac{P_R(c, u)}{w(u)\lambda_B(c/w(u))} = \kappa_* u^{\gamma(1-\beta)} \frac{(c^\beta - w(u)^\beta) \log(c/w(u))}{w(u)\lambda_B(c/w(u))} = \kappa_* \frac{(z^\beta - 1) \log z}{\lambda_B(z)} \geq \kappa_*$$

Furthermore, we have  $h_U(u) = s_0 \gamma (1-\gamma) U^\gamma F_\gamma(u/U)$  and  $\widehat{h}_U(u) = (1-1/s_0)h_U(u)$ , which makes  $k_D(U)$  independent of  $U$ .

We summarize the result as follows:

**Corollary 3.4 (Simplistic EEP)** *We consider the ERDS (3.12) with the choices (3.16) and set  $S(c, u) = s_0 u^\gamma - u^\gamma \lambda_B(c/u^\gamma)$ . Assume further*

$$1 - 2/d \leq \gamma \leq \frac{4\delta_c \delta_u}{(\delta_c + \delta_u)^2}, \tag{3.17}$$

then we have the following EEP estimate

$$\forall (c, u) \in \widehat{\mathfrak{S}}(U) : \mathcal{P}(c, u) \geq K \mathcal{H}_U(c, u) \text{ with } K = \min\{\kappa_*, \delta_u \frac{s_0 - 1}{s_0} \rho(\Omega, \gamma, \gamma)\} > 0.$$

Of course, in the given case with  $\kappa(c, u) = \kappa_* u^{\gamma(1-\beta)}$  we have uniform decay of  $c$  in the ODE  $\dot{c} = \kappa(c, u)(u^{\gamma\beta} - c^\beta)$ . Hence, it is very easy to show that every solution of  $(c, u)$  of the ERDS (3.12) converges to the equilibrium  $(w(\mathbf{U}), \mathbf{U})$ . Nevertheless the above result is interesting, since it is not so clear that the functional  $\mathcal{H}_{\mathbf{U}}$  satisfies the differential inequality

$$\frac{d}{dt} \mathcal{H}_{\mathbf{U}}(c(t), u(t)) \leq -K \mathcal{H}_{\mathbf{U}}(c(t), u(t))$$

uniformly of all solutions in  $\widehat{\mathfrak{S}}(\mathbf{U})$ .

### 3.3.3 A nontrivial case of the simple ERDS

The strength of the convexity method lies in the fact that it also can be used in some cases where the reactive entropy production  $P_{\mathbf{R}}(\mathbf{c}, u)$  does not provide an upper bound for the relative entropy  $\mathfrak{B}(\mathbf{c} | \mathbf{w}(u))$ . We only give one nontrivial example, namely the special case (3.16), but now specify  $\gamma$  and  $\beta$  as well, namely

$$w(u) = (u+b)^{1/2}, \quad f(c) = c^2, \quad \kappa(c, u) = \kappa_* = \text{const.}, \quad s(u) = 2u^{1/2}, \quad (3.18)$$

for some  $b > 0$ . Thus, the relevant relative entropy reads

$$\begin{aligned} H_{\mathbf{U}}(c, u) &= h_{\mathbf{U}}(u) + (u+b)^{1/2} \lambda_{\mathbf{B}}(c/(u+b)^{1/2}) \text{ with} \\ h_{\mathbf{U}}(u) &= \mathbf{U}^{-1/2} (u^{1/2} - \mathbf{U}^{1/2})^2 = \frac{\mathbf{U}^{1/2}}{2} F_{1/2}(u/\mathbf{U}) \end{aligned} \quad (3.19)$$

However, we now use a higher order entropy for  $u$ , namely  $\alpha = 1 > \gamma = 1/2$ , see also (3.7). Again relying on an enforced version of the restriction (3.17), we now assume

$$d \leq 3 \quad \text{and} \quad \delta_c^{\text{red}} := \delta_c \left( 1 - \frac{(\delta_c + \delta_u)^2}{8\delta_c \delta_u} \right) > 0, \quad (3.20)$$

where the last condition restricts  $\delta_c/\delta_u$  to  $]3-2\sqrt{2}, 3+2\sqrt{2}[$ . Using  $\widehat{h}''(u) = w''(u) - s''(u) \geq \frac{1}{4}u^{-3/2}$ , Proposition 2.2 provides an estimate for the diffusive entropy production, viz.

$$\delta_c \mathcal{P}_c(c, u) + \delta_u \mathcal{P}_u(c, u) \geq \int_{\Omega} \left( p_{\Omega}(\bar{u}) (\lambda_{\mathbf{B}}(u) - \lambda_{\mathbf{B}}(\bar{u})) + q_{\Omega} (\lambda_{\mathbf{B}}(c) - \lambda_{\mathbf{B}}(\bar{c})) \right) dx,$$

where  $p_{\Omega}(\mathbf{U}) = \delta_u \rho(\Omega, 1/2, 1)/(4\mathbf{U}^{1/2})$  and  $q_{\Omega} = \delta_c^{\text{red}} \rho(\Omega, 1, 1)$ . Theorem A.1 shows that both constants are strictly positive for  $d \leq 3$ . On the right-hand side we have now chosen a higher entropy for  $u$ , which leads to an extra factor  $\mathbf{U}^{-1/2}$  in  $p_{\Omega}$ , see (3.5). Note also that we have dropped the term  $\theta \delta_c \bar{c} F_1(c/\bar{c})$ , since we will not need it.

For the total entropy production we now have the lower bound

$$\mathcal{P}(c, u) \geq \int_{\Omega} \left( p_{\Omega}(\bar{u}) (\lambda_{\mathbf{B}}(u) - \lambda_{\mathbf{B}}(\bar{u})) + q_{\Omega} (\lambda_{\mathbf{B}}(c) - \lambda_{\mathbf{B}}(\bar{c})) + \frac{\kappa_*}{2} G(c^2, u+b) \right) dx,$$

where  $G(a, b) = (a-b) \log(a/b) \geq 0$ . Now we can exploit the convexity result from [MHM15, Lem. 4.3] which states that the function  $(c, u) \mapsto L_{a,b}(c, u) = \lambda_{\mathbf{B}}(u) + aG(c^2, u+b)$  is convex for  $b = 0$  if  $0 \leq a \leq a_{\text{crit}} \approx 0.8565$ . Calculating the second derivative gives

$$D^2 L_{a,b}(c, u) = D^2 L_{a,0}(c, u+b) + \left( \frac{1}{u} - \frac{1}{u+b} \right) e_2 \otimes e_2 \geq D^2 L_{a,0}(c, u+b) \geq 0,$$

which shows that  $L_{a,b}$  is convex for all  $b \geq 0$  and  $a \in [0, a_{\text{crit}}]$ . Hence, we choose  $\theta \in ]0, 1[$  and set

$$\mathcal{F}_{\theta, \mathbf{U}}(c, u) := \int_{\Omega} \left( \theta p_{\Omega}(\mathbf{U})(\lambda_{\mathbf{B}}(u) - \lambda_{\mathbf{B}}(\mathbf{U})) + q_{\Omega}(\lambda_{\mathbf{B}}(c) - \lambda_{\mathbf{B}}(\bar{c})) \right) dx.$$

Using  $G(\bar{c}^2, r) \geq r \lambda_{\mathbf{B}}(\bar{c}/r^{1/2})/6$  we obtain, for all  $(c, u) \in \widehat{\mathfrak{S}}(\mathbf{U})$ , the estimate

$$\begin{aligned} \mathcal{P}(c, u) &\geq \mathcal{F}_{\theta, \mathbf{U}}(c, u) + \int_{\Omega} \left( (1-\theta) p_{\Omega}(\mathbf{U})(\lambda_{\mathbf{B}}(u) - \lambda_{\mathbf{B}}(\mathbf{U})) + \frac{\kappa_*}{2} G(c^2, u+b) \right) dx \\ &\geq \mathcal{F}_{\theta, \mathbf{U}}(c, u) + M_{\theta, \mathbf{U}} \int_{\Omega} \left( \lambda_{\mathbf{B}}(u) - \lambda_{\mathbf{B}}(\mathbf{U}) + a_{\text{crit}} G(c^2, u+b) \right) dx \\ &\stackrel{\text{Jensen}}{\geq} \mathcal{F}_{\theta, \mathbf{U}}(c, u) + M_{\theta, \mathbf{U}} \left( \lambda_{\mathbf{B}}(\mathbf{U}) - \lambda_{\mathbf{B}}(\mathbf{U}) + G(\bar{c}^2, \mathbf{U}+b) \right) \\ &\geq \mathcal{F}_{\theta, \mathbf{U}}(c, u) + \frac{1}{6} M_{\theta, \mathbf{U}} (\mathbf{U}+b) \lambda_{\mathbf{B}}(\bar{c}/(\mathbf{U}+b)^{1/2}) \text{ with } M_{\theta, \mathbf{U}} = \min \left\{ \frac{\kappa_*}{2a_{\text{crit}}}, (1-\theta) p_{\Omega}(\mathbf{U}) \right\}. \end{aligned}$$

Using the relation  $\int_{\Omega} (\lambda_{\mathbf{B}}(c) - \lambda_{\mathbf{B}}(\bar{c}) + a \lambda_{\mathbf{B}}(\bar{c}/a)) dx = \int_{\Omega} a \lambda_{\mathbf{B}}(c/a) dx$  for  $a = (\mathbf{U}+b)^{1/2}$  and setting  $N_{\theta, \mathbf{U}} = \min \{ q_{\Omega}, (\mathbf{U}+b)^{1/2} M_{\theta, \mathbf{U}}/6 \}$ , we obtain the lower estimate

$$\begin{aligned} \mathcal{P}(c, u) &\geq \int_{\Omega} \left( \theta p_{\Omega}(\mathbf{U}) \mathbf{U} \lambda_{\mathbf{B}}(u/\mathbf{U}) + N_{\theta, \mathbf{U}} (\mathbf{U}+b)^{1/2} \lambda_{\mathbf{B}}(c/(\mathbf{U}^{1/2}+b)) \right) dx \\ &\geq L_{\theta, \mathbf{U}} \int_{\Omega} \tilde{H}_{\mathbf{U}}(c, u) dx \text{ where } \tilde{H}_{\mathbf{U}}(c, u) = h_{\mathbf{U}}(u) + (\mathbf{U}+b)^{1/2} \lambda_{\mathbf{B}}(c/(\mathbf{U}+b)^{1/2}) \end{aligned}$$

and  $L_{\theta, \mathbf{U}} = \min \{ \theta p_{\Omega}(\mathbf{U}) \mathbf{U}^{1/2}, N_{\theta, \mathbf{U}} \}$ . We finally use a quantitative version of Lemma 3.1 for our specific application.

**Lemma 3.5** *The functions  $H_{\mathbf{U}}$  (cf. (3.19)) and  $\tilde{H}_{\mathbf{U}}$  satisfy the estimate*

$$\forall c, u, \mathbf{U}, b > 0 : \quad \tilde{H}_{\mathbf{U}}(c, u) \geq C_H(b, \mathbf{U}) H_{\mathbf{U}}(c, u) \text{ with } C_H(b, \mathbf{U}) = \frac{8}{9 + 2 \log(1 + \mathbf{U}/b)}.$$

**Proof:** We let  $\beta = b/\mathbf{U}$  and define the three auxiliary functions

$$h_{\beta}(v) = \left( \left( v^2 - \frac{\beta}{1+\beta} \right)^{1/2} - \left( \frac{1}{1+\beta} \right)^{1/2} \right)^2, \quad B_{\beta}(a, v) = h_{\beta}(v) + v \lambda_{\mathbf{B}}(a/v), \quad \tilde{B}_{\beta}(a, v) = h_{\beta}(v) + \lambda_{\mathbf{B}}(a).$$

Inserting  $c = (\mathbf{U}+b)^{1/2} a$  and  $u = (\mathbf{U}+b)v^2 - b$ , we obtain

$$\frac{1}{C(\mathbf{U}, b)} := \sup_{c, u \geq 0} \frac{H_{\mathbf{U}}(c, u)}{\tilde{H}_{\mathbf{U}}(c, u)} = \sup \left\{ \frac{B_{\beta}(a, v)}{\tilde{B}_{\beta}(a, v)} \mid a \geq 0, v \geq \left( \frac{\beta}{1+\beta} \right)^{1/2} \right\} =: g(\beta), \text{ where } \beta = b/\mathbf{U}.$$

We decompose the quotient  $B_{\beta}/\tilde{B}_{\beta}$  into three parts that can be estimated separately.

$$\begin{aligned} \frac{B_{\beta}(a, v)}{\tilde{B}_{\beta}(a, v)} &= 1 + \frac{v-1-\log v}{\tilde{B}_{\beta}(a, v)} + \frac{(1-a) \log v}{\tilde{B}_{\beta}(a, v)} =: 1 + T_1(a, v) + T_2(v) \\ &\leq 1 + \frac{v-1-\log v}{(v-1)^2} + \frac{|1-a| |v-1|}{(v-1)^2 + \frac{1}{2}(a-1)^2} \leq 1 + \frac{1}{2} + 2 = 7/2, \end{aligned}$$

Using  $h_{\beta}(v) \geq (v-1)^2$  we bound  $T_1$  for  $a > 0$  and  $v \geq v_{\beta} := (\beta/(1+\beta))^{1/2} \in ]0, 1[$  via

$$T_1(a, v) \leq \frac{v-1-\log v}{(v-1)^2} \leq \frac{1}{2} + \log(1/v_{\beta}).$$

For  $T_2$  we observe that it is negative for  $(1-a)(v-1) < 0$ . For  $(a, v) \in [0, 1] \times [1, \infty[$  we easily find  $T_2(a, v) \leq 2^{-1/2}$ , where the supremum is attained for  $(a, v) \rightarrow (1, 1)$ . For  $(a, v) \in [1, \infty[ \times ]0, 1]$  one can show numerically that

$$T_2(a, v) \leq \log(1/v) + 2^{-1/2} \leq \log(1/v_\beta) + 2^{-1/2}.$$

Analytically one chooses  $a_1 > 1$ , then for  $a \geq a_1$  we have

$$T_2(a, v) \leq \frac{(a-1) \log(1/v)}{\lambda_B(a)} \leq \frac{a_1-1}{\lambda_B(a_1)} \log(1/v_\beta). \quad (3.21)$$

For  $a \in [1, a_1]$  we have  $\lambda_B(a) \geq \rho_1(a-1)^2$  with  $\rho_1 := \lambda_B(a_1)/(a_1-1)^2 < 1/2$ . For  $v \in [v_\beta, 1]$  we further have  $\log(1/v) \leq (1 + \log(1/v_\beta))(1-v)$  and obtain

$$T_2(a, v) \leq \frac{(a-1)(1 + \log(1/v_\beta))(1-v)}{(v-1)^2 + \rho_1(a-1)^2} \leq \frac{1 + \log(1/v_\beta)}{2\sqrt{\rho_1}}$$

Optimizing the position of  $a_1$  one gets an estimate that is qualitatively the same as (3.21).

Together we found  $g(\beta) \leq 2 \log(1/v_\beta) + 9/4$ , which gives the desired result using  $\log(1/v_\beta) = \frac{1}{2} \log(1+U/b)$ .  $\blacksquare$

We can now summarize the EEP estimate for the simple ERDS (3.12) under the specification (3.18). Inspecting all the above definitions of constants depending on  $\theta$ , we see that the optimal value is  $\theta = 1/7$ . With this choice we obtain the following result.

**Theorem 3.6 (Constructive EEP estimate for (3.12))** *For  $\mathcal{H}_U$  and  $\mathcal{P}$  as defined above via the choices (3.18) and (3.19), we have the following estimate*

$$\forall (c, u) \in \widehat{\mathfrak{S}}(\mathbf{U}) : \quad \mathcal{P}(c, u) \geq K_b(\mathbf{U}) \mathcal{H}_U(c, u), \quad \text{where} \quad (3.22)$$

$$K_b(\mathbf{U}) = \frac{8}{9 + \log(1+U/b)} \min \left\{ \frac{\delta_u}{28} \rho(\Omega, 1/2, 1), \delta_c^{\text{red}} \rho(\Omega, 1, 1), \frac{\kappa_*}{11} (U+b)^{1/2} \right\}.$$

This result is almost quasi-optimal in the sense, that it behaves linearly in the three limiting factors, namely the diffusion in  $c$ , the diffusion in  $u$ , and the convergence in the ODE  $\dot{c} = \kappa_*(U+b-c^2)$ . It remains unclear, whether the slowly degenerating prefactor  $(9 + 2 \log(1+U/b))^{-1}$  is really needed or simply an artefact of our method.

However, we see that the condition  $b > 0$  is really needed to obtain a nontrivial decay rate  $K_b(\mathbf{U})$ . The origin of the dependence on  $b$  is the estimate in Lemma 3.5, and it is open whether the result still holds in the case  $b = 0$ . The following remark indicates a possible way to treat cases with  $w_i(0) = 0$ .

**Remark 3.7 (Case  $w_i(0) = 0$ )** *Throughout our methods we experience difficulties in handling the case  $w_i(0) = 0$ . The point is the  $H_U(\mathbf{c}, u) = h_U(u) + \mathfrak{B}(\mathbf{c} | \mathbf{w}(u))$  does not have a good upper bound if  $w_i(u)$  is very small.*

*A possible way to handle such cases might be to use an entropy  $s$  that is unbounded at  $u = 0$  like  $s(u) = s_0 \log u$  leading to  $h_U(u) = F_0(u/U)$ . We conjecture that for all  $U > 0$  there exists  $C_H(\mathbf{U}) > 0$  such that the estimate*

$$F_0(u/U) + U \lambda_B(u/U) + U^{1/2} \lambda_B(c/U^{1/2}) \geq C_H(\mathbf{U}) \left( F_0(u/U) + U \lambda_B(u/U) + u^{1/2} \lambda_B(c/u) \right)$$

*holds true for all  $c, u > 0$ . However, in that case we are forced to use the generalized log-Sobolev estimate (3.6) for the case  $\gamma = 0$ , which is only possible for space dimensions  $d \leq 2$ , see Appendix B.*

**Remark 3.8** It was shown in [Mie16, Sec. 3.3] that for  $\beta \in [1, 22]$  the function  $(c, u) \mapsto \mu(\beta)\lambda_{\text{B}}(u) + (c^\beta - u)\log(c^\beta/u)$  is convex, if  $\mu(\beta)$  is chosen sufficiently large. Hence, the above convexity method can be adapted to cover the simple ERDS

$$\dot{c} = \delta_c \Delta c + \kappa(w(u)^\beta - c^\beta), \quad \dot{u} = \delta_u \Delta u,$$

if we choose  $w(u) = r(b+u)^{1/\beta}$  with  $r, b > 0$  and a suitable  $s(\cdot)$ , e.g.  $s(u) = s_0 u^\gamma + w(u)$  with  $\gamma > 1 - 2/d$ .

## 4 The Glitzky-Hünlich approach for ERDS

In this section we discuss the quite general result for uniform exponential decay developed in [GIH97] for semiconductor models. There the theory is restricted to space dimensions 1 or 2, because charged particles are considered which are coupled via the Poisson equation. In [Mie16] it is shown that the result transfers to general space dimensions, if electronic charges are not present or can be ignored. Here we generalize the method to the case of non-isothermal systems, where the detailed balance equilibria  $\mathbf{w}(u)$  depend on the internal energy  $u$ . This is nontrivial and leads to dimension restrictions for entropies of the form  $s(u) = cu^\gamma$ , namely we will need  $2_d \gamma > 2$ , where  $2_d > 2$  is the Lebesgue exponent such that  $H^1(\Omega)$  embeds continuously into  $L^{2_d}(\Omega)$ .

### 4.1 General setup and result

We recall the definition of the equilibria  $\mathbf{w}(\mathbf{q}, \mathbf{U})$  which are assumed to be the unique equilibria of the ODEs  $\dot{\mathbf{c}} = \mathbf{R}(\mathbf{c}, \mathbf{U})$  satisfying condition  $\mathbf{c} \in \mathbf{C}_{\mathbf{q}}$ , i.e.  $\mathbf{Q}\mathbf{c} = \mathbf{q}$ . As before we define the relative entropy

$$\mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) = \int_{\Omega} H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}(x), u(x)) \, dx \text{ with } H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) = h_{\mathbf{U}}(u) + \mathfrak{B}(\mathbf{c} | \mathbf{w}(\mathbf{q}, \mathbf{U})).$$

As before we let  $\widehat{h}_{\mathbf{U}}(u) = h_{\mathbf{U}}(u) - I + \sum_{i=1}^I w_i(u)$  and assume that

$$\begin{aligned} 0 &= h_{\mathbf{U}}(\mathbf{U}) < h_{\mathbf{U}}(u) \text{ for } u \neq \mathbf{U}, \quad \widehat{h}_{\mathbf{U}} \text{ is convex,} \\ w_i &\text{ is concave for } i = 1, \dots, I. \end{aligned}$$

As a consequence  $(\mathbf{c}, u) \mapsto H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u)$  is convex, non-negative, and  $H_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u) = 0$  if and only if  $(\mathbf{c}, u) = (\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$ .

Since we are not interested in quantitative results, we can use the simplified form  $\mathcal{P} = \widehat{\mathcal{P}}_{\gamma} + \widehat{\mathcal{P}}_{\text{R}}$  with  $\gamma \in ]0, 1[$  for the entropy production, where we recall

$$\widehat{\mathcal{P}}_{\gamma}(\mathbf{c}, u) = \int_{\Omega} \left( \frac{|\nabla u|^2}{u^{2-\gamma}} + \sum_{i=1}^I \frac{|\nabla c_i|^2}{c_i} \right) dx \text{ and } \widehat{\mathcal{P}}_{\text{R}}(\mathbf{c}, u) := \int_{\Omega} \widehat{P}_{\text{R}}(\mathbf{c}, u) \, dx, \quad (4.1)$$

$$\text{where } \widehat{P}_{\text{R}}(\mathbf{c}, u) = \sum_{r=1}^R G\left(\frac{\mathbf{c}^{\alpha^r}}{\mathbf{w}(u)^{\alpha^r}}, \frac{\mathbf{c}^{\beta^r}}{\mathbf{w}(u)^{\beta^r}}\right) dx \text{ with } G(a, b) = (a-b)\log(a/b).$$

We will use the following basic assumptions on the entropy function  $h_{\mathbf{U}}$  and on the equilibria relations  $c_i = w_i(u)$ . Throughout, we assume that  $(\mathbf{q}, \mathbf{U})$  and  $\gamma \in ]0, 1[$  are fixed,

and all subsequent constants may depend on these values with special indication.

$$\exists C > 0 \forall u \geq 0 : \frac{1}{C} \min\{|u-\mathbf{U}|, (u-\mathbf{U})^2\} \leq h_{\mathbf{U}}(u) \leq C \min\{|u-\mathbf{U}|, (u-\mathbf{U})^2\}, \quad (4.2a)$$

$$\exists C > 0 \forall u \geq 0, i = 1, \dots, I : 0 < w_i(0) \leq w_i(u) \leq C(1+u)^\gamma; \quad (4.2b)$$

$$\forall i = 1, \dots, I : \text{ the function } w_i \text{ is concave}; \quad (4.2c)$$

$$\text{the function } \widehat{h}_{\mathbf{U}} = h_{\mathbf{U}} - I + \sum_{i=1}^I w_i \text{ is convex.} \quad (4.2d)$$

Obviously, the functions  $w_i$  must be increasing, since they are concave and bounded from below by the positive value  $w_i(0)$ .

The main result of this section is the follow EEP estimate that will be obtained by a compactness method that is much more general than the convexity method discussed in the previous section. However, in this result the constant  $K_M(\mathbf{q}, \mathbf{U})$  also depends on the upper bound  $M$  for  $\mathcal{H}_{\mathbf{q}, \mathbf{U}}$ .

**Theorem 4.1 (EEP estimate via compactness)** *Consider a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^d$  and let the functions  $h$  and  $w_i$  satisfy the assumptions (4.2) with*

$$2_d \gamma > 2, \quad \text{where } 2_d = \frac{2d}{d-2} \text{ for } d \geq 3 \text{ and } \infty \text{ else.} \quad (4.3)$$

Moreover, assume the unique equilibrium condition UEC (2.17). Then, for each  $M > 0$  and each  $(\mathbf{q}, \mathbf{U}) \in \mathfrak{Q}$ , there exists a constant  $K_M(\mathbf{q}, \mathbf{U}) > 0$  such that

$$\forall (\mathbf{c}, u) \in \mathfrak{S}(\mathbf{q}, \mathbf{U}) \text{ with } \mathcal{H}(\mathbf{c}, u) \leq M : \widehat{\mathcal{P}}_\gamma(\mathbf{c}, u) + \widehat{\mathcal{P}}_{\mathbf{R}}(\mathbf{c}, u) \geq K_M(\mathbf{q}, \mathbf{U}) \mathcal{H}_{\mathbf{q}, \mathbf{U}}(\mathbf{c}, u). \quad (4.4)$$

The proof of this result is the content of the remaining subsection of Section 4. It relies on two basic facts about functionals  $\mathcal{F}(\lambda, \mathbf{z}) = \int_\Omega F(\lambda, \mathbf{z}(x)) \, dx$ , where  $\Omega$  is an open and bounded subset of  $\mathbb{R}^d$ .

(a) If  $F : [0, 1] \times \mathbb{R}^m \rightarrow \mathbb{R} \cup \{\infty\}$  is non-negative and lower semicontinuous, then

$$(\lambda_n, \mathbf{y}_n) \rightarrow (\lambda^*, \mathbf{y}^*) \text{ in } \mathbb{R} \times L^1(\Omega)^m \implies \mathcal{F}(\lambda^*, \mathbf{z}^*) \leq \liminf_{n \rightarrow \infty} \mathcal{F}(\lambda_n, \mathbf{z}_n). \quad (4.5a)$$

(b) If  $F : [0, 1] \times \mathbb{R}^m \rightarrow \mathbb{R}$  is continuous and satisfies  $0 \leq F(\lambda, \mathbf{z}) \leq C(1+|\mathbf{z}|)^p$  for some  $p \geq 1$  and  $C > 0$ , then

$$(\lambda_n, \mathbf{y}_n) \rightarrow (\lambda^*, \mathbf{y}^*) \text{ in } \mathbb{R} \times L^p(\Omega)^m \implies \mathcal{F}(\lambda_n, \mathbf{z}_n) \rightarrow \mathcal{F}(\lambda^*, \mathbf{z}^*). \quad (4.5b)$$

Clearly, (4.5a) follows from Fatou's lemma and (4.5b) from Lebesgue's dominated convergence theorem, see [FoL07, Lem. 1.83] and [FoL07, Thm. 1.85]. The major point is that we will apply (4.5a) for the functional  $\widehat{\mathcal{P}}_{\mathbf{R}}$ , where the density  $\widehat{\mathcal{P}}_{\mathbf{R}}$  may have arbitrarily high growth because  $\boldsymbol{\alpha}^r, \boldsymbol{\beta}^r \in \mathbb{N}_0^I$  are not restricted. In contrast, the continuity result (4.5b) will be used for  $\mathcal{H}$  or a rescaled version of it, where the growth is fixed and independent of the stoichiometric vectors  $\boldsymbol{\alpha}^r$  and  $\boldsymbol{\beta}^r$ .

In the rest of this section, the value of  $(\mathbf{q}, \mathbf{U}) \in \mathfrak{Q}$  is fixed. Hence, we drop the subscripts at  $\mathcal{H}$  and  $H$  in the sequel.

## 4.2 Some preliminary estimates

Here, we provide upper and lower bounds that are needed to obtain coercivity as well as the upper bound for applying (4.5b).

We first provide a lower bound on the relative entropy density  $H$ . We use that the function  $\lambda_B : z \mapsto z \log z - z + 1$  satisfies the elementary estimates

$$\forall \delta > 0 \exists C_\delta > 0 \forall z \geq 0 : \quad 4(1-\sqrt{z})^2 \leq \lambda_B(z) \leq C_\delta(1-\sqrt{z})^2(1+z)^\delta. \quad (4.6)$$

This implies  $w\lambda_B(c/w) \geq 4(\sqrt{z} - \sqrt{w})^2 \geq 2z - 4w$  and with (4.2a), (4.2b), and  $\gamma < 1$ , we obtain

$$\exists C > 0 \forall \mathbf{c}, u : \quad H(\mathbf{c}, u) \geq \frac{1}{C}u - C + 2 \sum_{i=1}^I c_i. \quad (4.7)$$

Since  $\widehat{\mathcal{P}}_{\text{diff}}$  gives a bound on  $\nabla(u^{\gamma/2})$  it will be important to control  $h$  and  $w_i$  in terms of  $v = u^{\gamma/2}$ . We will use the following results.

**Lemma 4.2** *Assume that  $h$  and  $w_i$  satisfy (4.2) with  $\mathbf{U} > 0$  and  $\gamma \in ]0, 1[$ . Then, there exists  $\widehat{c}_h, C, C_w > 0$  such that*

$$\forall u \geq 0 : \quad \widehat{c}_h |u^{\gamma/2} - \mathbf{U}^{\gamma/2}|^2 \leq h(u) \leq C(|u^{\gamma/2} - \mathbf{U}^{\gamma/2}|^2 + |u^{\gamma/2} - \mathbf{U}^{\gamma/2}|^{2/\gamma}), \quad (4.8a)$$

$$\forall v \geq -\mathbf{U}^{\gamma/2} : \quad \left| (w_i[(\mathbf{U}^{\gamma/2} + v)^{2/\gamma}]^{1/2} - (w_i(\mathbf{U}))^{1/2}) \right| \leq C_w |v|. \quad (4.8b)$$

**Proof:** The lower bound in (4.8a) follows simply by observing that the lower bound  $h$  in (4.2a) and the left-hand side of (4.8a) both behave as  $(u - \mathbf{U})^2$  for  $u$  near  $\mathbf{U}$ . Otherwise  $h$  is strictly positive with linear growth for  $u \rightarrow \infty$ , while the left-hand side in (4.8a) has only growth like  $u^\gamma$  with  $\gamma < 1$ .

For the upper bound we argue similarly that  $h$  and the right-hand side behave like  $(u - \mathbf{U})^2$  for  $u$  near  $\mathbf{U}$ . Otherwise  $h$  and the right-hand side are bounded linearly, so the result follows.

To show (4.8b) we let  $m(v) = (\mathbf{U}^{\gamma/2} + v)^{2/\gamma}$  and  $\widetilde{w}(v) = [w(m(v))]^{1/2}$ . Hence, we have to find  $C$  such that  $|\widetilde{w}(v) - \widetilde{w}(0)| \leq C|v|$ .

*Case  $v \in [-\mathbf{U}^{\gamma/2}, 0]$ :* Concavity and monotonicity of  $w_i$  implies concavity and monotonicity of  $w_i^{1/2}$ . Using  $m(v) \leq m(0) = \mathbf{U}$  we immediately find  $\widetilde{w}(0) \geq \widetilde{w}(v)$ . Thus, it suffices to estimate  $\widetilde{w}(v) - \widetilde{w}(0)$  by  $C_-v$  from below. Using  $m(0) = \mathbf{U}$  we have

$$\begin{aligned} \widetilde{w}(v) - \widetilde{w}(0) &= w(m(v))^{1/2} - w(\mathbf{U})^{1/2} \geq \left(1 - \frac{m(v)}{\mathbf{U}}\right)w(0)^{1/2} + \frac{m(v)}{\mathbf{U}}w(\mathbf{U})^{1/2} - w(\mathbf{U})^{1/2} \\ &= (m(v) - m(0))\frac{1}{\mathbf{U}}(w(\mathbf{U})^{1/2} - w(0)^{1/2}). \end{aligned}$$

Since  $w(\mathbf{U}) \geq w(0)$  it suffices to estimate  $m(v) - m(0)$  from below. Because  $m$  is convex we find  $m(v) - m(0) \geq m'(0)v$ , where  $m'(0) > 0$ . Thus, for  $v \in [-\bar{u}^{\gamma/2}, 0]$  we have

$$\widetilde{w}(v) - \widetilde{w}(0) \geq C_-v \quad \text{with } C_- = 2(w(\mathbf{U})^{1/2} - w(0)^{1/2})/(\gamma \mathbf{U}^{\gamma/2}).$$

*Case  $v \geq 0$ :* We have  $\widetilde{w}'(v) = \frac{w'(m(v))}{2\widetilde{w}(v)}m'(v)$ . Moreover, concavity of  $w$  implies  $0 < w(0) \leq w(u) + w'(u)(0-u)$ , which implies  $w'(u) \leq w(u)/u$ . With  $m'(v) = \frac{2}{\gamma}m(v)^{1-\gamma/2}$  and  $u = m(v)$  we have

$$\widetilde{w}'(v) = \frac{w'(u)}{\gamma w(u)^{1/2}}u^{1-\gamma/2} \leq \frac{w(u)^{1/2}}{\gamma u^{\gamma/2}} \leq \frac{1}{\gamma} \left( \frac{C(1+u^\gamma)}{u^\gamma} \right)^{1/2} \leq C_+ \quad \text{for } u \geq \mathbf{U}.$$

Thus, we conclude  $\widetilde{w}(0) \leq \widetilde{w}(v) \leq \widetilde{w}(0) + C_+v$  for  $v \geq 0$ , and (4.8b) is established with  $C_w = \max\{C_-, C_+\}$ .  $\blacksquare$

### 4.3 Extraction of a converging subsequence

We assume that the result is not true and will generate a contradiction at the end of Section 4.6. Since the result is not true, there is a sequence  $\mathbf{y}_n = (\mathbf{c}_n, u_n)$  such that

$$0 < \mathcal{H}(\mathbf{y}_n) \leq R, \quad \mathbf{Q}\mathbf{y}_n = \mathbf{q}, \quad \widehat{\mathcal{P}}(\mathbf{y}_n) := \widehat{\mathcal{P}}_\gamma(\mathbf{y}_n) + \widehat{\mathcal{P}}_R(\mathbf{y}_n) \leq \frac{1}{n} \mathcal{H}(\mathbf{y}_n) \leq \frac{M}{n}. \quad (4.9)$$

We define the auxiliary functions

$$a_{n,i} := c_{n,i}^{1/2} \text{ for } i = 1, \dots, I \quad \text{and} \quad a_{n,I+1} := u_n^{\gamma/2},$$

and define the sequence of vector-valued function  $\mathbf{a}_n : \Omega \rightarrow [0, \infty]^{I+1}$ . The bound  $\mathcal{H}(\mathbf{y}_n) \leq M$  together with the coercivity (4.7) yields the bound

$$\|\mathbf{a}_n\|_{L^2(\Omega)^{I+1}} \leq C.$$

Moreover,  $\widehat{\mathcal{P}}_\gamma(\mathbf{y}_n) \leq \widehat{\mathcal{P}}(\mathbf{y}_n) \leq M/n$  gives the bound

$$\|\nabla \mathbf{a}_n\|_{L^2(\Omega)^{I+1}} \leq \frac{C}{\sqrt{n}}.$$

Thus, we conclude that, up to the extraction of a subsequence, the sequence  $\mathbf{a}_n$  converges strongly in  $H^1(\Omega)$  to a limit  $\mathbf{a}^*$  with  $\nabla \mathbf{a}^* \equiv 0$ , i.e. each component of the limit is a constant function.

Choosing  $q > 1$  with  $2q \in ]2/\gamma, 2d]$ , which exists by assumption (4.3), we obtain  $\mathbf{a}_n \rightarrow \mathbf{a}^*$  in  $L^{2q}(\Omega)^{I+1}$ . Recalling the definition of  $\mathbf{a}_n$  yields

$$\begin{aligned} c_{n,i} &= (a_{n,i})^2 \rightarrow (a_i^*)^2 =: c_i^* \text{ in } L^q(\Omega), \\ u_n &= (a_{n,I+1})^{2/\gamma} \rightarrow (a_{I+1}^*)^{2/\gamma} =: u^* \text{ in } L^{q\gamma}(\Omega). \end{aligned} \quad (4.10)$$

This defines a limit vector  $\mathbf{y}^* = (\mathbf{c}^*, u^*)$ .

### 4.4 Identification of the limit $\mathbf{a}^*$

Since the functional  $\mathbf{Q}$  is linear, it is continuous in  $L^1(\Omega)$  and we conclude

$$\mathbf{Q}\mathbf{y}^* = \lim_{n \rightarrow \infty} \mathbf{Q}\mathbf{y}_n = \lim_{n \rightarrow \infty} \mathbf{q} = \mathbf{q}.$$

By energy conservation, this means in particular that  $u^* = \mathbf{U}$ .

To identify the other components of  $\mathbf{a}^*$  and thus the limits of  $c_{n,i}$  we use the relation  $\mathcal{P}_R(\mathbf{y}_n) \leq M/n$ . We note that the integrand  $\widehat{\mathcal{P}}_R$  of  $\mathcal{P}_R$  is non-negative and continuous, since  $0 < w_i(0) \leq w_i(u)$ , see (4.2a). Hence,  $\mathcal{P}_R$  is lower semicontinuous with respect to the strong topology in  $L^1(\Omega)^{I+1}$ , see (4.5a). This implies

$$0 \leq \mathcal{P}_R(\mathbf{c}^*, \mathbf{U}) \leq \liminf_{n \rightarrow \infty} \mathcal{P}_R(\mathbf{c}_n, u_n) \leq \lim_{n \rightarrow \infty} M/n = 0.$$

Together with  $\mathbf{Q}\mathbf{y}^* = \mathbf{q}$  and the uniqueness assumption (2.17), we conclude that  $\mathbf{y}^*$  is the unique steady state in the stoichiometric subspace, i.e.  $\mathbf{y}^* = (\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U})$ .

## 4.5 Blowup near the equilibrium

To generate a contradiction we set

$$\lambda_n^2 := \mathcal{H}(\mathbf{y}_n) > 0$$

and first show  $\lambda_n \rightarrow 0$ . This follows by using the fact that the density  $H(\mathbf{c}, u) = h(u) + \sum w_i(u)\lambda_B(c_i/w_i(u))$  is a continuous function and satisfies, for all  $\delta > 0$ , the bound

$$0 \leq H(\mathbf{c}, u) \leq C_\delta \left(1 + u + (1 + |\mathbf{c}| + u^\gamma)(1 + |\mathbf{c}|)^\delta\right)$$

for some  $C_\delta > 0$ . For this we use (4.6), giving  $w_i\lambda_B(c_i/w_i) \leq C_\delta(c_i^{1/2} - w_i^{1/2})^2(1 + c_i/w_i)^\delta \leq \tilde{C}_\delta(c_i + w_i)(1 + c_i)^\delta$ , together with (4.2a) and (4.2b).

Choosing  $\delta \in ]0, \gamma q - 1]$ , where we use (4.3) once again, we see that  $\mathcal{H}$  is strongly continuous on  $L^q(\Omega)^I \times L^{\gamma q}(\Omega)$  such that

$$\lim_{n \rightarrow \infty} \lambda_n^2 = \lim_{n \rightarrow \infty} \mathcal{H}(\mathbf{y}_n) = \mathcal{H}(\mathbf{y}^*) = \mathcal{H}(\mathbf{w}(\mathbf{q}, \mathbf{U}), \mathbf{U}) = 0,$$

as desired.

We now define the blowup functions

$$b_{n,i} = \frac{1}{\lambda_n} (a_{n,i} - a_i^*) \quad \text{for } i = 1, \dots, I+1.$$

Since  $\widehat{\mathcal{P}}_\gamma$  expressed in  $\mathbf{a}$  is exactly quadratic in  $\nabla a_i$ , we obtain

$$\begin{aligned} 1 &= \frac{1}{\lambda_n^2} \mathcal{H}(\mathbf{y}_n) \geq \frac{n}{\lambda_n^2} \widehat{\mathcal{P}}_\gamma(\mathbf{y}_n) = \frac{n}{4\lambda_n^2} \left( \gamma^2 \|\nabla a_{n,I+1}\|_{L^2}^2 + \sum_1^I \|\nabla a_{n,i}\|_{L^2}^2 \right) \\ &= \frac{n}{4} \left( \gamma^2 \|\nabla b_{n,I+1}\|_{L^2}^2 + \sum_1^I \|\nabla b_{n,i}\|_{L^2}^2 \right). \end{aligned} \quad (4.11)$$

To control  $\mathbf{b}_n$  without derivatives we use a rescaled coercivity of  $H$ , namely

$$H(\mathbf{c}, u) \geq \widehat{c}_h |u^{\gamma/2} - \mathbf{U}^{\gamma/2}|^2 + 4 \sum_1^I (c_i^{1/2} - w_i(u)^{1/2})^2,$$

which follows from the lower estimate in (4.8a) and  $w\lambda_B(c/w) \geq 4(c^{1/2} - w^{1/2})^2$ . Inserting  $\mathbf{y}_n = \Phi(\lambda_n \mathbf{b}_n)$  given by

$$\Phi(\lambda \mathbf{b})_i = (w_i(\mathbf{U})^{1/2} + \lambda b_i)^2 \quad \text{and} \quad \Phi(\lambda, \mathbf{b})_{I+1} = (\mathbf{U}^{\gamma/2} + \lambda b_{I+1})^{2/\gamma} \quad (4.12)$$

and choosing any  $\vartheta \in ]0, 4[$  we arrive at the estimate

$$\begin{aligned} \frac{1}{\lambda_n^2} H(\mathbf{c}_n, u_n) &\geq \widehat{c}_h |b_{n,I+1}|^2 + \frac{\vartheta}{\lambda_n^2} \sum_{i=1}^I \left( \lambda_n b_{n,i} + w_i(\mathbf{U})^{1/2} - [w_i((\mathbf{U}^{\gamma/2} + \lambda_n b_{n,I+1})^{2/\gamma})]^{1/2} \right)^2 \\ &\stackrel{(1)}{\geq} \widehat{c}_h |b_{n,I+1}|^2 + \vartheta \sum_{i=1}^I \left( \frac{1}{2} b_{n,i}^2 - C_w^2 b_{n,I+1}^2 \right) \stackrel{(2)}{\geq} \frac{1}{C} \left( |b_{n,I+1}|^2 + \sum_{i=1}^I \frac{1}{2} b_{n,i}^2 \right). \end{aligned}$$

In “ $\stackrel{(1)}{\geq}$ ” we first used  $(a+b)^2 \geq \frac{1}{2}a^2 - b^2$  and then (4.8b), while “ $\stackrel{(2)}{\geq}$ ” follows by choosing  $\vartheta$  suitable, e.g.  $\vartheta = \min\{4, \widehat{c}_h/(2IC_w^2)\}$ . As a result we conclude that

$$1 = \frac{1}{\lambda_n^2} \mathcal{H}(\mathbf{y}_n) \geq \frac{1}{C} \|\mathbf{b}_n\|_{L^2}^2.$$

Together with (4.11) we see that, after choosing a subsequence (not relabeled),

$$\mathbf{b}_n \rightarrow \mathbf{b}^* = \text{const.} \quad \text{in } H^1(\Omega)^{I+1}.$$

## 4.6 The final contradiction

Here and in the sequel we shortly write  $\mathbf{w} = (w_1, \dots, w_I) = \mathbf{w}(\mathbf{q}, \mathbf{U})$ . The mapping  $\mathbf{y} = \Phi(\lambda \mathbf{b})$  introduced in (4.12) satisfies

$$\Phi(0) = (\mathbf{w}, \mathbf{U}) \quad \text{and} \quad D\Phi(0) = \text{diag}(2w_1^{1/2}, \dots, 2w_I^{1/2}, \frac{2}{\gamma} \mathbf{U}^{1-\gamma/2}),$$

which shows that the weights  $w_i(\mathbf{U})^{1/2}$  are relevant. We will show that  $\mathbf{b}^*$  satisfies the following three relations

$$\mathbf{s} := (w_i^{1/2} b_i^*)_{i=1, \dots, I} \in \mathbb{S} \quad \text{and} \quad \mathbf{b}_{I+1}^* = 0; \quad (4.13a)$$

$$\sum_{r=1}^R \left( (\boldsymbol{\alpha}^r - \boldsymbol{\beta}^r) \cdot \mathbf{d} \right)^2 = 0 \quad \text{with} \quad \mathbf{d} := (b_i^*/w_i^{1/2})_{i=1, \dots, I}, \quad \text{giving} \quad \mathbf{d} \in \mathbb{S}^\perp; \quad (4.13b)$$

$$\mathbf{b}^* \neq 0. \quad (4.13c)$$

Obviously, these three relations cannot hold simultaneously, since  $\mathbf{s} \in \mathbb{S}$  and  $\mathbf{d} \in \mathbb{S}^\perp$  implies  $0 = \mathbf{s} \cdot \mathbf{d}$ , but obviously  $\mathbf{s} \cdot \mathbf{d} = |\mathbf{b}^*|^2$ , which contradicts (4.13c).

To derive (4.13a) we observe  $\mathbf{Q}\mathbf{y}_n = \mathbf{q} = \mathbf{Q}\mathbf{y}^*$ , hence  $\mathbf{z}_n = \frac{1}{\lambda_n} (\mathbf{y}_n - \mathbf{y}^*)$  satisfies  $\mathbf{Q}\mathbf{z}_n = 0$ . Moreover, the convergence of  $\mathbf{b}_n$  and the relations (4.12) easily provide the convergence

$$\mathbf{z}_n \rightarrow \mathbf{z}^* := \left( 2w_1^{1/2} b_1^*, \dots, 2w_I^{1/2} b_I^*, \frac{2}{\gamma} \mathbf{U}^{1-\gamma/2} b_{I+1}^* \right) = \left( 2\mathbf{s}, \frac{2}{\gamma} \mathbf{U}^{1-\gamma/2} b_{I+1}^* \right).$$

Thus, (4.13a) is established, since it is equivalent to  $\mathbf{Q}\mathbf{z}^* = 0$ .

To obtain (4.13b) we use the relation

$$1 = \frac{1}{\lambda_n^2} \mathcal{H}(\mathbf{y}_n) \geq \frac{n}{\lambda_n^2} \widehat{\mathcal{P}}_R(\mathbf{y}_n), \quad \text{which implies} \quad \frac{1}{\lambda_n^2} \widehat{\mathcal{P}}_R(\Phi(\lambda_n, \mathbf{b}_n)) \rightarrow 0. \quad (4.14)$$

To exploit this, we define the auxiliary functional

$$\begin{aligned} \mathcal{K}_R(\lambda, \mathbf{b}) &:= \frac{1}{\lambda^2} \widehat{\mathcal{P}}_R(\Phi(\lambda, \mathbf{b})) = \int_{\Omega} K_R(\lambda, \mathbf{b}(x)) \, dx, \quad \text{where} \\ K_R(\lambda, \mathbf{b}) &= \frac{1}{\lambda^2} \widehat{\mathcal{P}}_R(\Phi(\lambda, \mathbf{b})) \quad \text{for } \lambda > 0 \text{ and} \\ 2K_R(0, \mathbf{b}) &= \sum_{r=1}^R \left( (\boldsymbol{\alpha}^r - \boldsymbol{\beta}^r) \cdot \left( \frac{2b_i}{w_i^{1/2}} - \mu \frac{w'_i}{w_i} b_{I+1} \right) \right)^2 \quad \text{with } w'_i = \partial_{\mathbf{U}} w_i(\mathbf{q}, \mathbf{U}), \end{aligned}$$

where  $\mu = 2/(\gamma \mathbf{U}^{\gamma/2-1})$  and  $\widehat{\mathcal{P}}_R$  may be extended by  $\infty$  if  $\mathbf{y} = \Phi(\lambda, \mathbf{b})$  does not lie in  $[0, \infty]^{I+1}$ . By this definition, the function  $K_R$  is non-negative and lower semicontinuous in  $(\lambda, \mathbf{b}) \in$

$[0, \infty[ \times \mathbb{R}^{I+1}$ , hence the functional  $\mathcal{K}_R$  is strongly lower semicontinuous in  $L^2$ , see (4.5a). Thus,

$$\mathbf{b}_n \rightarrow \mathbf{b}^* \text{ implies } \mathcal{K}_R(\lambda_n, \mathbf{b}_n) \rightarrow 0 = \mathcal{K}_R(0, \mathbf{b}^*) = K_R(0, \mathbf{b}^*).$$

Since we already know  $b_{I+1}^* = 0$ , we conclude (4.13b).

For the final relation we introduce the integral density function  $K_H$  via

$$\begin{aligned} \text{for } \lambda \in ]0, 1]: \quad K_H(\lambda, \mathbf{b}) &= \frac{1}{\lambda^2} H(\Phi(\lambda, \mathbf{b})), \\ \text{for } \lambda = 0: \quad K_H(0, \mathbf{b}) &= \frac{4h''(\mathbf{U})}{\gamma^2 \mathbf{U}^{\gamma-2}} b_{I+1}^2 + \sum_{i=1}^I \left( 2b_i - \frac{\mathbf{w}'_i}{\mathbf{w}_i^{1/2}} \mu b_{I+1} \right)^2. \end{aligned}$$

We define  $K_H$  only on the closed subset

$$\text{dom}(K_H) = \{0\} \times \mathbb{R}^{I+1} \cup \{(\lambda, \mathbf{b}) \in ]0, 1] \times \mathbb{R}^{I+1} \mid b_{I+1} \geq -\frac{\mathbf{U}^{\gamma/2}}{\lambda}, b_i \geq -\frac{\mathbf{w}_i^{1/2}}{\lambda} \text{ for } i=1, \dots, I\}.$$

By construction, the function  $K_H$  is non-negative and continuous on its domain. We now establish an upper bound for  $K_H$  using the upper bound for  $h$  in (4.8a) and the upper bound of  $w\lambda_B(c/w) \leq C_\delta(c^{1/2}-w^{1/2})^2(1+c/w)^\delta$ , see (4.6). Using the shorthand  $\Phi_{I+1}(\mathbf{b}) = (\mathbf{U}^{\gamma/2} + \lambda b_{I+1})^{2/\gamma}$  we have

$$\begin{aligned} \frac{h(\Phi_{I+1}(\mathbf{b}))}{\lambda^2} &\leq \frac{C}{\lambda^2} \left( (\lambda b_{I+1})^2 + (\lambda b_{I+1})^{2/\gamma} \right) \leq C(b_{I+1}^2 + b_{I+1}^{2/\gamma}), \\ \frac{w_i(u)\lambda_B\left(\frac{c_i}{w_i(u)}\right)}{\lambda^2} &\leq \frac{C_\delta}{\lambda^2} \left( \mathbf{w}_i^{1/2} + \lambda b_i - w_i(\Phi_{I+1}(\mathbf{b}))^{1/2} \right)^2 \left( 1 + \frac{(\mathbf{w}_i^{1/2} + \lambda b_i)^2}{w_i(\Phi_{I+1}(\mathbf{b}))} \right)^\delta \\ &\stackrel{*}{\leq} C_\delta (|b_i| + C_w |b_{I+1}|)^2 \left( 1 + (\mathbf{w}_i^{1/2} + \lambda b_i)^2 / w_i(0) \right)^\delta \\ &\leq C_{K_H} (1 + |\mathbf{b}|)^{2+2\delta} \quad \text{for all } (\lambda, \mathbf{b}) \in \text{dom}(K_H), \end{aligned}$$

where we used  $\lambda \in ]0, 1]$  and  $\gamma \in ]0, 1[$  and employed (4.8b) and  $w_i(u) \geq w_i(0)$  for “ $\leq^*$ ”.

Of course, the construction of  $K_H$  was such that, for  $\lambda > 0$ , we have

$$\frac{1}{\lambda^2} \mathcal{H}(\Phi(\lambda, \mathbf{b})) = \mathcal{K}_H(\lambda, \mathbf{b}) := \int_{\Omega} K_H(\lambda, \mathbf{b}(x)) \, dx.$$

The upper bound on  $K_H$  and its continuity imply continuity of  $\mathcal{K}_H$ , see (4.5b). More precisely, from  $\mathbf{b}_n \rightarrow \mathbf{b}_*$  in  $H^1(\Omega)^{I+1}$  we obtain strong convergence in  $L^{2q}(\Omega)$  for some  $q > 1$ . Choosing  $\delta \in ]0, q-1]$  we use that for all  $\mathbf{y}_n$  we have  $(\lambda, \mathbf{b}_n(x)) \in \text{dom}(K_H)$  for almost all  $x \in \Omega$ . Then, using (4.5b), we obtain

$$1 = \frac{1}{\lambda_n^2} \mathcal{H}(\mathbf{y}_n) = \mathcal{K}_H(\lambda_n, \mathbf{b}_n) \rightarrow \mathcal{K}_H(0, \mathbf{b}_*) = \left( \frac{4h''(u)}{\gamma^2 \mathbf{U}^{\gamma-2}} (b_{I+1}^*)^2 + \sum_{i=1}^I \left( 2b_i^* - \frac{\mathbf{w}'_i}{\mathbf{w}_i^{1/2}} \mu b_{I+1}^* \right)^2 \right).$$

Thus, (4.13c) is established and the contradiction is complete.

This finishes the proof of Theorem 4.1.

## A Generalized EEP estimates

This appendix gives a proof of the EEP estimate (3.6) for general  $\gamma$  and  $\alpha$ . For a more detailed exposition, we refer to [Mit14]. We will use the Poincaré-Sobolev inequality

$$\|\nabla u\|_p \geq \eta_{PS}(\Omega, p, q) \|u - \bar{u}\|_q \tag{A.1}$$

corresponding to the Sobolev embedding  $W^{1,p}(\Omega) \hookrightarrow L^q(\Omega)$ , where  $\eta_{PS}(\Omega, p, q)$  is the optimal Poincaré-Sobolev constant. This inequality holds true for  $p \geq d$  or for  $p < d$  and  $q \leq dp/(d-p)$ .

**Theorem A.1** *Let  $\gamma > \max\{0, 1-2/d\}$  and  $\alpha \leq \gamma+2/d$  for  $d \geq 3$  or  $\alpha < 1+\gamma$  for  $d \leq 2$ . Then  $\rho(\Omega, \gamma, \alpha)$  is strictly positive and can be bounded by*

$$\rho(\Omega, \gamma, \alpha) \geq \frac{4\alpha(\gamma-2/q)}{\gamma^2(q-1)} \eta_{PS}(\Omega, 2, q)^2 \quad (\text{A.2})$$

with  $q \geq \max\{2/\gamma, 2/(1+\gamma-\alpha)\}$ .

**Proof:** We substitute  $v = u^{\gamma/2}$ . Then

$$\begin{aligned} \bar{u}^\gamma \int_{\Omega} F_{\alpha_1} \left( \frac{u}{\bar{u}} \right) dx &\leq \frac{\alpha_2 \bar{u}^\gamma}{\alpha_1} \int_{\Omega} F_{\alpha_2} \left( \frac{u}{\bar{u}} \right) dx = C \left( \frac{\|v\|_x^x}{\|v\|_y^{x-2}} - \|v\|_y^2 \right) \\ &\leq C(\|v\|_z^2 - \|v\|_y^2) \leq C(z-1) \|v - \bar{v}\|_z^2 \leq \frac{C(z-1)}{\eta_{PS}(\Omega, 2, z)^2} \|\nabla v\|_2^2 \end{aligned}$$

with  $\alpha_2 > 1$ ,  $C = 1/(\alpha_1(\alpha_2-1))$ ,  $x = 2\alpha_2/\gamma$ ,  $y = 2/\gamma$ , and  $z = 2/(1+\gamma-\alpha_2)$ . In the first line we used monotonicity of the map  $\alpha \mapsto \alpha F_\alpha(z)$ , see (3.2). In the last line, we applied the Hölder inequality  $\|v\|_x^x \leq \|v\|_y^{x-2} \|v\|_z^2$ , the Poincaré-Sobolev inequality and the following estimate which holds true for  $x \geq 2$  (see [Mit14] for a proof):

$$\|v\|_x^2 - \|v\|_y^2 \leq \|v\|_x^2 - \|v\|_1^2 \leq (x-1) \|v - \bar{v}\|_x^2$$

This proves the theorem. ■

**Remark A.2** *The above theorem still remains true for the cases  $\gamma = 1 - 2/d$  as well as  $\alpha = \gamma + 1$  for  $d \leq 2$ , see [Mit14]. In the case  $\alpha = 2$  and  $\gamma = 1$ , which will be used below, we have the simple bound*

$$\rho(\Omega, 1, 2) \geq 2 \eta_{PS}(\Omega, 1, 2)^2, \quad (\text{A.3})$$

which follows from the following chain of estimates:

$$\bar{u} \int_{\Omega} \frac{|\nabla u|^2}{u} dx \geq \|\nabla u\|_1^2 \geq \eta_{PS}(\Omega, 1, 2)^2 \|u - \bar{u}\|^2 = 2 \eta_{PS}(\Omega, 1, 2)^2 \cdot \bar{u}^2 \int_{\Omega} F_2(u/\bar{u}) dx.$$

## B The EEP estimate for $\alpha = \gamma = 0$ and $d \leq 2$

This appendix provides a proof that the EEP inequality also holds in the case  $\alpha = \gamma = 0$ . Here we only give a densified version of the full proof and refer to [Mit14] for a detailed exposition, which is based on the approach developed in [CaL92, CCL10]. We note that  $\rho(\Omega, 1, 2)$  is positive only for  $\Omega \subset \mathbb{R}^d$  with  $d \leq 2$ .

**Theorem B.1 (Case  $\alpha = \gamma = 0$  in dimension  $d \leq 2$ )** *We have the estimate  $\rho(\Omega, 0, 0) \geq \varrho := \rho(\Omega, 1, 2) > 0$ , i.e.*

$$\forall u > 0 : \int_{\Omega} \frac{|\nabla u(x)|^2}{u(x)^2} dx \geq \rho(\Omega, 1, 2) \int_{\Omega} F_0(u(x)/U) dx. \quad (\text{B.1})$$

**Proof:** We set  $u = e^v$  and observe that (B.1) is equivalent to

$$H(v) := \int_{\Omega} |\nabla v|^2 dx \geq \varrho G(v) \text{ with } G(v) := \left( \log \left( \int_{\Omega} e^v dx \right) - \int_{\Omega} v dx \right). \quad (\text{B.2})$$

We consider  $H$  and  $G$  as nonnegative, proper, and convex functionals on  $L^2(\Omega)$ . Because of  $\Omega \subset \mathbb{R}^d$  with  $d \leq 2$ , the Moser-Trudinger inequality  $G$  is bounded whenever  $H$  is bounded.

We establish (B.2) by Legendre transform for which we note

$$H^*(\psi) = \frac{1}{4} \int_{\Omega} (\psi - \bar{\psi})(-\Delta)^{-1}(\psi - \bar{\psi}) dx \text{ and } G^*(\psi) = \begin{cases} \int_{\Omega} \Gamma(\psi(x)) dx & \text{if } \bar{\psi} = 0, \\ \infty & \text{else,} \end{cases}$$

where  $\Gamma(\psi) = \lambda_B(\psi+1)$  for  $\psi \geq -1$  and  $\infty$  otherwise.

We now define  $J(\psi) = G^*(\psi) - \varrho H^*(\psi)$  and show  $J(\psi_0) \geq 0$  for all  $\psi_0$ . For this we note  $J(0) = 0$  and then consider  $J$  along the solutions  $\psi_t := e^{t\Delta}\psi_0$  of the diffusion equation  $\dot{\psi} = \Delta\psi$ . Setting  $\psi_t = c_t - 1$  with  $\bar{c}_t = 1$  and  $c_t \geq 0$ . By differentiating along solutions and using  $\dot{c}_t = \Delta c_t$  (with Neumann boundary conditions) we obtain

$$\frac{d}{dt} J(c_t - 1) = - \int_{\Omega} \frac{|\nabla c_t|^2}{c_t} dx + \frac{\varrho}{2} \int_{\Omega} (c_t - 1)^2 dx \geq 0,$$

since  $\varrho = \rho(\Omega, 1, 2)$  which is the optimal constant for the last estimate. Hence  $J(\psi_0) \geq J(\psi_t) \geq J(0) = 0$  since  $\psi_t \rightarrow 0$ .

Now we reverse the Legendre transform using  $H^*(\varrho\psi) = \varrho^2 H^*(\psi)$ , hence

$$\begin{aligned} H(v) &= \sup (\langle v, \psi \rangle - H^*(\psi)) \stackrel{\psi = \varrho\tilde{\psi}}{=} \sup (\langle v, \varrho\tilde{\psi} \rangle - \varrho^2 H^*(\tilde{\psi})) \\ &\stackrel{J \geq 0}{=} \varrho \sup (\langle v, \tilde{\psi} \rangle - G^*(\tilde{\psi})) = \varrho G(v). \end{aligned}$$

This proves the assertion. ■

## References

- [AGH02] G. ALBINUS, H. GAJEWSKI, and R. HÜNLICH. Thermodynamic design of energy models of semiconductor devices. *Nonlinearity*, 15(2), 367–383, 2002.
- [BD\*10] M. BURGER, M. DI FRANCESCO, J.-F. PIETSCHMANN, and B. SCHLAKE. Nonlinear cross-diffusion with size exclusion. *SIAM J. Math. Analysis*, 42(6), 2842–2871, 2010.
- [BoD15] D. BOTHE and W. DREYER. Continuum thermodynamics of chemically reacting fluid mixtures. *Acta Mech.*, 226, 1757–1805, 2015.
- [CaL92] E. A. CARLEN and M. LOSS. Competing symmetries, the logarithmic hls inequality and onofri’s inequality on  $\mathbf{S}^n$ . *Geometric & Functional Analysis GFA*, 2(1), 90–104, 1992.
- [CCL10] E. A. CARLEN, J. A. CARRILLO, and M. LOSS. Hardy-littlewood-sobolev inequalities via fast diffusion flows. *Proc. Natl. Acad. Sci. U.S.A.*, 107(46), 19696–701, 2010.
- [DD\*16] W. DREYER, P.-É. DRUET, P. GAJEWSKI, and C. GUHLKE. Existence of weak solutions for improved nernst–planck–poisson models of compressible reacting electrolytes. *WIAS preprint 2291*, 2016.
- [DeF06] L. DESVILLETES and K. FELLNER. Exponential decay toward equilibrium via entropy methods for reaction-diffusion equations. *J. Math. Anal. Appl.*, 319(1), 157–176, 2006.
- [DFT16] L. DESVILLETES, K. FELLNER, and B. Q. TANG. Trend to equilibrium for reaction-diffusion systems arising from complex balanced chemical reaction networks. *arXiv:1604.04536*, 2016.

- [Fei73] M. FEINBERG. Complex balancing in general kinetic systems. *Arch. Rational Mech. Anal.*, 49, 187–194, 1972/73.
- [FeT16] K. FELLNER and B. Q. TANG. Explicit exponential convergence to equilibrium for nonlinear reaction-diffusion systems with detailed balance condition. *arXiv:1601.05992*, 2016.
- [Fis15] J. FISCHER. Global existence of renormalized solutions to entropy-dissipating reaction-diffusion systems. *Arch. Rational Mech. Anal.*, 218(1), 553–587, 2015.
- [FoL07] I. FONSECA and G. LEONI. *Modern Methods in the Calculus of Variations:  $L^p$  spaces*. Springer, 2007.
- [GliH97] A. GLITZKY and R. HÜNLICH. Energetic estimates and asymptotic for electro-reaction-diffusion systems. *Z. angew. Math. Mech. (ZAMM)*, 77(11), 823–832, 1997.
- [HH\*16] S. HITTMER, J. HASKOVEC, P. A. MARKOWICH, and A. MIELKE. Decay to equilibrium for energy-reaction-diffusion systems. *SIAM J. Math. Analysis*, 2016. Submitted. WIAS preprint 2233.
- [JKO98] R. JORDAN, D. KINDERLEHRER, and F. OTTO. The variational formulation of the Fokker-Planck equation. *SIAM J. Math. Analysis*, 29(1), 1–17, 1998.
- [JüM06] A. JÜNGEL and D. MATTHES. An algorithmic construction of entropies in higher-order nonlinear PDEs. *Nonlinearity*, 19(3), 633–659, 2006.
- [Jün15] A. JÜNGEL. The boundedness-by-entropy method for cross-diffusion systems. *Nonlinearity*, 28, 1963–2001, 2015.
- [LiM13] M. LIERO and A. MIELKE. Gradient structures and geodesic convexity for reaction-diffusion systems. *Phil. Trans. Royal Soc. A*, 371(2005), 20120346, 28, 2013.
- [LiY99] E. H. LIEB and J. YNGVASON. The physics and mathematics of the second law of thermodynamics. *Phys. Rep.*, 310(1), 96, 1999.
- [MHM15] A. MIELKE, J. HASKOVEC, and P. A. MARKOWICH. On uniform decay of the entropy for reaction-diffusion systems. *J. Dynam. Diff. Eqns.*, 27(3-4), 897–928, 2015.
- [Mie11a] A. MIELKE. Formulation of thermoelastic dissipative material behavior using GENERIC. *Contin. Mech. Thermodyn.*, 23(3), 233–256, 2011.
- [Mie11b] A. MIELKE. A gradient structure for reaction-diffusion systems and for energy-drift-diffusion systems. *Nonlinearity*, 24, 1329–1346, 2011.
- [Mie13] A. MIELKE. Thermomechanical modeling of energy-reaction-diffusion systems, including bulk-interface interactions. *Discr. Cont. Dynam. Systems Ser. S*, 6(2), 479–499, 2013.
- [Mie16] A. MIELKE. Uniform exponential decay for reaction-diffusion systems with balanced mass-action kinetics. *In preparation*, 2016.
- [Mit14] M. MITTENZWEIG. Entropy production inequalities for heat equations. Master’s thesis, Freie Universität and WIAS Berlin, November 2014.
- [Ons31] L. ONSAGER. Reciprocal relations in irreversible processes, I+II. *Physical Review*, 37, 405–426, 1931. (part II, 38:2265–2279).
- [Ott01] F. OTTO. The geometry of dissipative evolution equations: the porous medium equation. *Comm. Partial Differential Equations*, 26, 101–174, 2001.