



IUCrData

ISSN 2414-3146

Received 27 May 2021

Accepted 1 June 2021

Edited by W. T. A. Harrison, University of Aberdeen, Scotland

Keywords: crystal structure; borate; amino acid.

CCDC reference: 2087276


Structural data: full structural data are available from iucrdata.iucr.org

(S)-Alanine ethyl ester tetracyanidoborate, (C₅H₁₂NO)[B(CN)₄]


Tim Peppel^{a*} and Martin Köckerling^b
^aLeibniz-Institut für Katalyse e.V. (LIKAT), Heterogene Photokatalyse, Albert-Einstein-Str. 29a, D-18059 Rostock, Germany, and ^bUniversität Rostock, Institut für Chemie, Anorganische Festkörperchemie, Albert-Einstein-Str. 3a, D-18059 Rostock, Germany. *Correspondence e-mail: tim.peppel@catalysis.de

The title molecular salt, C₅H₁₂NO⁺·C₄BN₄[−] or (C₅H₁₂NO)[B(CN)₄], was obtained as single crystals by slow evaporation of a solution of the compound in acetonitrile over several weeks. The asymmetric unit contains two (*S*)-alanine ethyl ester cations and two tetracyanidoborate anions, which are linked by N—H···N hydrogen bonds. The compound exhibits a relatively low melting point of 110°C and shows a solid–solid phase transition near room temperature (*T*_{s–s} = 29°C) on the basis of DSC measurements.

3D view



Chemical scheme



Structure description

For more than 20 years, ionic liquids as salts with low melting points have attracted great interest because of their unique properties and applications. These properties include for instance large liquid ranges, broad electrochemical windows as well as low vapour pressures (Hallett & Welton, 2011; Welton, 1999). The title compound acts as a first example of a low-melting chiral substance in our ongoing efforts to investigate tetracyanidoborate-based ionic liquids (Bernsdorf *et al.*, 2009; Flemming *et al.*, 2010; Siegesmund *et al.*, 2017).

The asymmetric unit of the title compound consists of two (*S*)-alanine ethyl ester cations and two tetracyanidoborate anions (Fig. 1). The conformations of the cations about the stereogenic centres (C10 and C15) are almost the same, as indicated by the C9—C10—C11—O2 and C14—C15—C16—O4 torsion angles of −61.9 (3) and −63.0 (3)°, respectively, but the conformations of the ethyl side chains differ substantially: C11—O2—C12—C13 = −86.1 (3) and C16—O4—C17—C18 = 136.5 (3)°. Otherwise, all bond lengths and angles within the cation are in the expected ranges (Dimitrijević *et al.*, 2013). The geometry around the B atoms is close to tetrahedral with C—B—C angles ranging from 107.8 (2) to 111.2 (2)°.



OPEN ACCESS

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N9-H9F\cdots N6^i$	0.91	2.16	2.920 (3)	141
$N10-H10B\cdots N4^{ii}$	0.91	2.05	2.953 (3)	174
$N10-H10D\cdots N1$	0.91	2.07	2.961 (3)	166
$N9-H9E\cdots N8^{iii}$	0.91	2.14	3.001 (3)	158
$N10-H10C\cdots N2^{iv}$	0.91	2.15	3.015 (3)	159
$N9-H9D\cdots N5^v$	0.91	2.14	3.017 (3)	161
$N9-H9F\cdots N3^{vi}$	0.91	2.64	3.147 (3)	116

Symmetry codes: (i) $x - \frac{1}{2}, y + \frac{1}{2}, z - 1$; (ii) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (iii) $x, y, z - 1$; (iv) $x, y + 1, z$; (v) $x - \frac{1}{2}, y - \frac{1}{2}, z - 1$; (vi) $-x + \frac{1}{2}, y + \frac{1}{2}, -z$.

In the extended structure, the shortest hydrogen-bond contacts are found between the N-bonded H atoms of the cations (N9 and N10) and the N atoms of the tetra-cyanidoborate anions: the shortest N $\cdots$ N donor-acceptor distance is 2.920 (3) Å (Table 1). Fig. 2 shows the packing of the ions within and around the unit cell.

Synthesis and crystallization

The title compound, (C₅H₁₂NO)[B(CN)₄], was obtained in high purity as a colorless solid on a multi-gram scale from the salt metathesis of (*S*)-alanine ethyl ester hydrochloride and K[B(CN)₄] in acetonic solution at room temperature. (*S*)-Alanine ethyl ester hydrochloride (2.0 g, 13.0 mmol) was added in one portion to a vigorously stirred solution of K[B(CN)₄] (2.2 g, 14.3 mmol) in 100 ml acetone at room temperature and was further stirred overnight. The precipitate was filtered off and the solvent of the filtrate was removed in vacuum. The residue was dissolved in a minimum amount of dichloromethane, filtered again and the solvent was removed in vacuum. The final product was obtained as a colourless solid in high yield (2.8 g, 91%); m.p. = 110°C, T_{s-s} = 29°C. The thermal behaviour was determined by means of differential scanning calorimetry (DSC) in the temperature range from –100 to 200°C with a heating rate of 10 K min^{–1}. Analytical data for C₉H₁₂BN₅O₂ % (calc.): C 46.43 (46.39); H 5.25 (5.19); N 26.53 (30.05).




Figure 1
The asymmetric unit of (C₅H₁₂NO)[B(CN)₄] with atom labelling.

Table 2

Experimental details.

Crystal data	
Chemical formula	C ₅ H ₁₂ NO ₂ ⁺ ·C ₄ N ₄ B [–]
M_r	233.05
Crystal system, space group	Monoclinic, C2
Temperature (K)	173
a, b, c (Å)	17.059 (1), 8.7467 (4), 18.855 (1)
β (°)	111.468 (4)
V (Å ³)	2618.2 (3)
Z	8
Radiation type	Mo $K\alpha$
μ (mm ^{–1})	0.09
Crystal size (mm)	0.27 × 0.18 × 0.15
Data collection	
Diffractometer	Bruker APEXII CCD
Absorption correction	Multi-scan (SADABS; Bruker, 2017)
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	11976, 7354, 5158
R_{int}	0.038
$(\sin \theta/\lambda)_{max}$ (Å ^{–1})	0.725
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.053, 0.133, 1.00
No. of reflections	7354
No. of parameters	307
No. of restraints	1
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{max}, \Delta\rho_{min}$ (e Å ^{–3})	0.38, –0.24
Absolute structure	Flack x determined using 1751 quotients $[(I^+) - (I^-)] / [(I^+) + (I^-)]$ (Parsons et al., 2013)
Absolute structure parameter	0.2 (8)

Computer programs: APEX2 and SAINT (Bruker, 2017), SHELXS (Sheldrick, 2015b), SHELXT (Sheldrick, 2015a), DIAMOND (Brandenburg & Putz, 2019) and publCIF (Westrip, 2010).

Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. Sixteen reflections were omitted from the refinement because their intensities were affected by the beam stop. Details can be found in the refine_special_




Figure 2
A view of the unit-cell contents in projection down the b axis.

details field in the CIF. The refined value of the Flack absolute structure parameter of 0.2 (8) was ambiguous, and the absolute structure was assigned on the basis of the enantiomeric pure (*S*)-alanine ethyl ester hydrochloride used in the synthesis.

Acknowledgements

The authors thank Dr A. Villinger (Universität Rostock) for maintaining the functionality of the X-ray facilities. The publication of this article was funded by the Open Access Fund of the Leibniz Association.

Funding information

Funding for this research was provided by: Deutsche Forschungsgemeinschaft (grant No. KO-1616-4 to MK).

References

- Bernsdorf, A., Brand, H., Hellmann, R., Köckerling, M., Schulz, A., Villinger, A. & Voss, K. (2009). *J. Am. Chem. Soc.* **131**, 8958–8970.
- Brandenburg, K. & Putz, H. (2019). *Diamond*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2017). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dimitrijević, D. P., Novaković, S. B., Radić, G. R., Jevtić, V. V., Menéndez-Taboada, L., García-Granda, S. & Trifunović, S. R. (2013). *J. Serb. Chem. Soc.* **78**, 1531–1537.
- Flemming, A., Hoffmann, M. & Köckerling, M. (2010). *Z. Anorg. Allg. Chem.* **636**, 562–568.
- Hallett, J. P. & Welton, T. (2011). *Chem. Rev.* **111**, 3508–3576.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Siegesmund, A., Topp, A., Nitschke, C. & Köckerling, M. (2017). *ChemistrySelect*, **2**, 11328–11335.
- Welton, T. (1999). *Chem. Rev.* **99**, 2071–2084.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

full crystallographic data

IUCrData (2021). **6**, x210562 [https://doi.org/10.1107/S2414314621005629]

(S)-Alanine ethyl ester tetracyanidoborate, (C₅H₁₂NO)[B(CN)₄]

Tim Peppel and Martin Köckerling

1-Ethoxy-1-oxopropan-2-aminium tetracyanoborate

Crystal data

C₅H₁₂NO₂⁺·C₄N₄B[−]

M_r = 233.05

Monoclinic, *C*2

a = 17.059 (1) Å

b = 8.7467 (4) Å

c = 18.855 (1) Å

β = 111.468 (4)°

V = 2618.2 (3) Å³

Z = 8

F(000) = 976

D_x = 1.182 Mg m^{−3}

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 2564 reflections

θ = 4.3–25.5°

μ = 0.09 mm^{−1}

T = 173 K

Block, colourless

0.27 × 0.18 × 0.15 mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: microfocus sealed tube

φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2017)

7354 independent reflections

5158 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.038

θ_{max} = 31.0°, θ_{min} = 4.4°

h = −24→24

k = −10→12

l = −25→26

11976 measured reflections

Refinement

Refinement on *F*²

Least-squares matrix: full

R [*F*² > 2σ(*F*²)] = 0.053

wR(*F*²) = 0.133

S = 1.00

7354 reflections

307 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0649*P*)²]

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} < 0.001

Δρ_{max} = 0.38 e Å^{−3}

Δρ_{min} = −0.24 e Å^{−3}

Absolute structure: Flack *x* determined using

1751 quotients [(*F*⁺)(*F*)]/[(*F*⁺)(*F*)] (Parsons et
al., 2013)

Absolute structure parameter: 0.2 (8)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
B1	0.4401 (2)	0.0236 (3)	0.3575 (2)	0.0267 (6)
C1	0.4986 (2)	0.1670 (3)	0.3916 (2)	0.0327 (6)
N1	0.5418 (2)	0.2686 (3)	0.4167 (2)	0.0494 (7)
C2	0.4971 (2)	−0.1273 (3)	0.3745 (2)	0.0281 (5)
N2	0.5371 (2)	−0.2346 (3)	0.3855 (1)	0.0405 (6)
C3	0.3940 (2)	0.0381 (3)	0.2677 (2)	0.0326 (6)
N3	0.3621 (2)	0.0434 (3)	0.2032 (2)	0.0505 (7)
C4	0.3729 (2)	0.0061 (3)	0.3968 (2)	0.0343 (6)
N4	0.3242 (2)	−0.0082 (4)	0.4246 (2)	0.0524 (7)
B2	0.4421 (2)	0.4179 (3)	0.8626 (2)	0.0288 (6)
C5	0.4893 (2)	0.5759 (3)	0.8893 (2)	0.0378 (6)
N5	0.5235 (2)	0.6898 (3)	0.9064 (2)	0.0574 (8)
C6	0.5081 (2)	0.2830 (3)	0.8927 (2)	0.0345 (6)
N6	0.5555 (2)	0.1866 (3)	0.9158 (2)	0.0505 (7)
C7	0.4013 (2)	0.4174 (4)	0.7723 (2)	0.0396 (7)
N7	0.3728 (2)	0.4224 (4)	0.7078 (2)	0.068 (1)
C8	0.3709 (2)	0.3968 (3)	0.8974 (2)	0.0304 (5)
N8	0.3198 (2)	0.3818 (3)	0.9222 (2)	0.0423 (6)
C9	0.1899 (2)	0.2660 (3)	0.0774 (2)	0.0485 (8)
H9A	0.1933	0.2721	0.1304	0.073*
H9B	0.2436	0.2290	0.0762	0.073*
H9C	0.1447	0.1952	0.0491	0.073*
C10	0.1714 (2)	0.4213 (3)	0.0417 (1)	0.0284 (5)
H10A	0.1168	0.4582	0.0436	0.034*
N9	0.1651 (1)	0.4193 (2)	−0.0392 (1)	0.0251 (4)
H9D	0.1236	0.3536	−0.0663	0.038*
H9E	0.2150	0.3881	−0.0414	0.038*
H9F	0.1529	0.5149	−0.0592	0.038*
C11	0.2398 (2)	0.5319 (3)	0.0848 (1)	0.0290 (5)
O1	0.2858 (1)	0.5932 (2)	0.0589 (1)	0.0390 (5)
O2	0.2429 (1)	0.5455 (2)	0.1560 (1)	0.0417 (5)
C12	0.3156 (2)	0.6236 (4)	0.2100 (2)	0.0534 (9)
H12A	0.3004	0.6699	0.2511	0.064*
H12B	0.3340	0.7064	0.1838	0.064*
C13	0.3855 (2)	0.5119 (6)	0.2431 (2)	0.069 (1)
H13A	0.4347	0.5646	0.2791	0.103*
H13B	0.4003	0.4663	0.2022	0.103*
H13C	0.3674	0.4313	0.2699	0.103*
C14	0.6152 (2)	0.5843 (4)	0.5570 (2)	0.0447 (7)
H14A	0.6329	0.6107	0.6111	0.067*
H14B	0.5816	0.4904	0.5470	0.067*
H14C	0.5813	0.6678	0.5261	0.067*
C15	0.6923 (2)	0.5598 (3)	0.5365 (1)	0.0262 (5)
H15A	0.7255	0.6569	0.5462	0.031*
N10	0.6667 (1)	0.5199 (2)	0.4547 (1)	0.0228 (4)

H10B	0.7135	0.5058	0.4430	0.034*
H10C	0.6352	0.5971	0.4258	0.034*
H10D	0.6358	0.4323	0.4451	0.034*
C16	0.7477 (2)	0.4339 (3)	0.5836 (1)	0.0246 (5)
O3	0.7610 (1)	0.3169 (2)	0.5581 (1)	0.0429 (5)
O4	0.7763 (2)	0.4724 (3)	0.6559 (1)	0.0500 (6)
C17	0.8324 (2)	0.3692 (4)	0.7118 (2)	0.0477 (8)
H17A	0.8557	0.2922	0.6864	0.057*
H17B	0.8015	0.3151	0.7397	0.057*
C18	0.9012 (2)	0.4620 (6)	0.7653 (2)	0.065 (1)
H18A	0.9405	0.3952	0.8037	0.097*
H18B	0.8776	0.5372	0.7904	0.097*
H18C	0.9313	0.5153	0.7371	0.097*

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
B1	0.024 (1)	0.028 (1)	0.028 (2)	−0.001 (1)	0.010 (1)	0.001 (1)
C1	0.035 (1)	0.033 (1)	0.030 (1)	−0.001 (1)	0.012 (1)	0.001 (1)
N1	0.057 (2)	0.042 (1)	0.045 (2)	−0.017 (1)	0.014 (1)	−0.006 (1)
C2	0.027 (1)	0.031 (1)	0.029 (1)	−0.001 (1)	0.013 (1)	0.004 (1)
N2	0.038 (1)	0.041 (1)	0.045 (2)	0.007 (1)	0.018 (1)	0.009 (1)
C3	0.032 (1)	0.031 (1)	0.033 (2)	−0.001 (1)	0.010 (1)	0.002 (1)
N3	0.051 (2)	0.060 (2)	0.033 (1)	−0.004 (1)	0.006 (1)	0.003 (1)
C4	0.029 (1)	0.038 (1)	0.037 (2)	0.001 (1)	0.014 (1)	0.000 (1)
N4	0.041 (1)	0.070 (2)	0.055 (2)	0.002 (1)	0.028 (1)	−0.001 (2)
B2	0.028 (1)	0.028 (1)	0.029 (2)	0.004 (1)	0.009 (1)	0.001 (1)
C5	0.027 (1)	0.036 (1)	0.048 (2)	0.002 (1)	0.011 (1)	0.006 (1)
N5	0.039 (1)	0.041 (2)	0.081 (2)	−0.006 (1)	0.009 (1)	0.003 (2)
C6	0.037 (1)	0.035 (1)	0.029 (2)	0.006 (1)	0.010 (1)	−0.002 (1)
N6	0.053 (2)	0.046 (2)	0.043 (2)	0.019 (1)	0.007 (1)	−0.003 (1)
C7	0.040 (2)	0.045 (2)	0.035 (2)	0.010 (1)	0.014 (1)	0.007 (1)
N7	0.070 (2)	0.098 (3)	0.034 (2)	0.023 (2)	0.015 (1)	0.012 (2)
C8	0.029 (1)	0.028 (1)	0.030 (1)	0.002 (1)	0.006 (1)	−0.002 (1)
N8	0.036 (1)	0.047 (1)	0.046 (2)	0.001 (1)	0.019 (1)	0.001 (1)
C9	0.077 (2)	0.033 (2)	0.034 (2)	−0.009 (2)	0.018 (2)	0.002 (1)
C10	0.028 (1)	0.032 (1)	0.026 (1)	−0.002 (1)	0.012 (1)	0.002 (1)
N9	0.026 (1)	0.0231 (9)	0.025 (1)	−0.0002 (8)	0.0083 (8)	−0.0007 (8)
C11	0.035 (1)	0.028 (1)	0.024 (1)	−0.001 (1)	0.011 (1)	0.002 (1)
O1	0.047 (1)	0.041 (1)	0.033 (1)	−0.0185 (9)	0.0192 (9)	−0.0055 (8)
O2	0.054 (1)	0.047 (1)	0.026 (1)	−0.016 (1)	0.0161 (8)	−0.0048 (8)
C12	0.073 (2)	0.056 (2)	0.026 (2)	−0.030 (2)	0.012 (2)	−0.010 (1)
C13	0.054 (2)	0.100 (3)	0.046 (2)	−0.014 (2)	0.010 (2)	−0.022 (2)
C14	0.049 (2)	0.060 (2)	0.030 (1)	0.027 (2)	0.020 (1)	0.009 (1)
C15	0.035 (1)	0.022 (1)	0.021 (1)	0.004 (1)	0.0097 (9)	0.0004 (9)
N10	0.0255 (9)	0.0230 (9)	0.021 (1)	0.0000 (8)	0.0092 (7)	0.0005 (7)
C16	0.025 (1)	0.027 (1)	0.022 (1)	0.0028 (9)	0.0096 (9)	0.0022 (9)
O3	0.059 (1)	0.032 (1)	0.032 (1)	0.0192 (9)	0.0095 (9)	−0.0016 (8)

O4	0.068 (1)	0.049 (1)	0.023 (1)	0.026 (1)	0.0046 (9)	0.0007 (9)
C17	0.052 (2)	0.062 (2)	0.026 (2)	0.027 (2)	0.010 (1)	0.013 (1)
C18	0.041 (2)	0.104 (3)	0.048 (2)	0.015 (2)	0.014 (2)	0.014 (2)

Geometric parameters (Å, °)

B1—C4	1.585 (4)	C11—O2	1.330 (3)
B1—C1	1.585 (4)	O2—C12	1.455 (3)
B1—C3	1.591 (4)	C12—C13	1.490 (6)
B1—C2	1.600 (4)	C12—H12A	0.9900
C1—N1	1.142 (4)	C12—H12B	0.9900
C2—N2	1.135 (3)	C13—H13A	0.9800
C3—N3	1.137 (3)	C13—H13B	0.9800
C4—N4	1.138 (4)	C13—H13C	0.9800
B2—C7	1.585 (4)	C14—C15	1.514 (4)
B2—C6	1.586 (4)	C14—H14A	0.9800
B2—C5	1.586 (4)	C14—H14B	0.9800
B2—C8	1.590 (4)	C14—H14C	0.9800
C5—N5	1.140 (4)	C15—N10	1.482 (3)
C6—N6	1.139 (4)	C15—C16	1.510 (3)
C7—N7	1.134 (4)	C15—H15A	1.0000
C8—N8	1.138 (3)	N10—H10B	0.9100
C9—C10	1.497 (4)	N10—H10C	0.9100
C9—H9A	0.9800	N10—H10D	0.9100
C9—H9B	0.9800	C16—O3	1.188 (3)
C9—H9C	0.9800	C16—O4	1.312 (3)
C10—N9	1.489 (3)	O4—C17	1.451 (3)
C10—C11	1.505 (4)	C17—C18	1.479 (5)
C10—H10A	1.0000	C17—H17A	0.9900
N9—H9D	0.9100	C17—H17B	0.9900
N9—H9E	0.9100	C18—H18A	0.9800
N9—H9F	0.9100	C18—H18B	0.9800
C11—O1	1.192 (3)	C18—H18C	0.9800
C4—B1—C1	110.0 (2)	C13—C12—H12A	109.8
C4—B1—C3	110.2 (2)	O2—C12—H12B	109.8
C1—B1—C3	111.2 (2)	C13—C12—H12B	109.8
C4—B1—C2	108.5 (2)	H12A—C12—H12B	108.2
C1—B1—C2	109.0 (2)	C12—C13—H13A	109.5
C3—B1—C2	107.8 (2)	C12—C13—H13B	109.5
N1—C1—B1	178.8 (3)	H13A—C13—H13B	109.5
N2—C2—B1	179.0 (3)	C12—C13—H13C	109.5
N3—C3—B1	177.5 (3)	H13A—C13—H13C	109.5
N4—C4—B1	179.1 (3)	H13B—C13—H13C	109.5
C7—B2—C6	111.0 (2)	C15—C14—H14A	109.5
C7—B2—C5	108.4 (2)	C15—C14—H14B	109.5
C6—B2—C5	108.9 (2)	H14A—C14—H14B	109.5
C7—B2—C8	110.0 (2)	C15—C14—H14C	109.5

C6—B2—C8	108.3 (2)	H14A—C14—H14C	109.5
C5—B2—C8	110.2 (2)	H14B—C14—H14C	109.5
N5—C5—B2	177.8 (3)	N10—C15—C16	108.8 (2)
N6—C6—B2	178.7 (3)	N10—C15—C14	110.3 (2)
N7—C7—B2	177.6 (4)	C16—C15—C14	111.6 (2)
N8—C8—B2	179.8 (3)	N10—C15—H15A	108.7
C10—C9—H9A	109.5	C16—C15—H15A	108.7
C10—C9—H9B	109.5	C14—C15—H15A	108.7
H9A—C9—H9B	109.5	C15—N10—H10B	109.5
C10—C9—H9C	109.5	C15—N10—H10C	109.5
H9A—C9—H9C	109.5	H10B—N10—H10C	109.5
H9B—C9—H9C	109.5	C15—N10—H10D	109.5
N9—C10—C9	112.0 (2)	H10B—N10—H10D	109.5
N9—C10—C11	108.2 (2)	H10C—N10—H10D	109.5
C9—C10—C11	110.3 (2)	O3—C16—O4	126.0 (2)
N9—C10—H10A	108.8	O3—C16—C15	124.2 (2)
C9—C10—H10A	108.8	O4—C16—C15	109.8 (2)
C11—C10—H10A	108.8	C16—O4—C17	119.4 (2)
C10—N9—H9D	109.5	O4—C17—C18	107.6 (3)
C10—N9—H9E	109.5	O4—C17—H17A	110.2
H9D—N9—H9E	109.5	C18—C17—H17A	110.2
C10—N9—H9F	109.5	O4—C17—H17B	110.2
H9D—N9—H9F	109.5	C18—C17—H17B	110.2
H9E—N9—H9F	109.5	H17A—C17—H17B	108.5
O1—C11—O2	125.8 (2)	C17—C18—H18A	109.5
O1—C11—C10	124.3 (2)	C17—C18—H18B	109.5
O2—C11—C10	109.8 (2)	H18A—C18—H18B	109.5
C11—O2—C12	117.2 (2)	C17—C18—H18C	109.5
O2—C12—C13	109.4 (3)	H18A—C18—H18C	109.5
O2—C12—H12A	109.8	H18B—C18—H18C	109.5
N9—C10—C11—O1	−7.2 (3)	N10—C15—C16—O3	−6.3 (3)
C9—C10—C11—O1	115.7 (3)	C14—C15—C16—O3	115.6 (3)
N9—C10—C11—O2	175.3 (2)	N10—C15—C16—O4	175.1 (2)
C9—C10—C11—O2	−61.9 (3)	C14—C15—C16—O4	−63.0 (3)
O1—C11—O2—C12	−9.5 (4)	O3—C16—O4—C17	2.5 (4)
C10—C11—O2—C12	168.1 (2)	C15—C16—O4—C17	−178.9 (3)
C11—O2—C12—C13	−86.1 (3)	C16—O4—C17—C18	136.5 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N9—H9F $\cdots$ N6 ⁱ	0.91	2.16	2.920 (3)	141
N10—H10B $\cdots$ N4 ⁱⁱ	0.91	2.05	2.953 (3)	174
N10—H10D $\cdots$ N1	0.91	2.07	2.961 (3)	166
N9—H9E $\cdots$ N8 ⁱⁱⁱ	0.91	2.14	3.001 (3)	158
N10—H10C $\cdots$ N2 ^{iv}	0.91	2.15	3.015 (3)	159

N9—H9 <i>D</i> ···N5 ^v	0.91	2.14	3.017 (3)	161
N9—H9 <i>F</i> ···N3 ^{vi}	0.91	2.64	3.147 (3)	116

Symmetry codes: (i) $x-1/2, y+1/2, z-1$; (ii) $x+1/2, y+1/2, z$; (iii) $x, y, z-1$; (iv) $x, y+1, z$; (v) $x-1/2, y-1/2, z-1$; (vi) $-x+1/2, y+1/2, -z$.