

2,4-Bis(diphenylphosphanyl)-1,1,2,3,3,4-hexaphenyl-1,3-diphospha-2,4-diboracyclobutane tetrahydrofuran sesqui-solvate

Normen Peulecke,* Bernd H. Müller, Anke Spannenberg and Uwe Rosenthal

Leibniz-Institut für Katalyse e.V. an der Universität Rostock, Albert-Einstein-Strasse 29a, 18059 Rostock, Germany

Correspondence e-mail: normen.peulecke@catalysis.de


Received 5 March 2012; accepted 15 March 2012

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in solvent or counterion; R factor = 0.041; wR factor = 0.096; data-to-parameter ratio = 19.5.

In the title compound, $\text{C}_{60}\text{H}_{50}\text{B}_2\text{P}_4 \cdot 1.5\text{C}_4\text{H}_8\text{O}$, the diphosphadiborane molecule lies on an inversion centre, whereas the disordered tetrahydrofuran solvent molecule is in a general position with a partial occupancy of 0.75. The diphosphadiborane molecule consists of an ideal planar four-membered B_2P_2 ring with an additional phenyl and a $-\text{PPh}_2$ group attached to each B atom.

Related literature

For the structure of a monomeric diphosphaborane molecule, see: Bartlett *et al.* (1988). For assumed monomeric $\text{PhB}(\text{PPh}_2)_2$, see: Coates & Livingstone (1961). For the structures of other dimeric boron-bridged bisphosphine compounds, see: Herdtweck *et al.* (1997); Kaufmann *et al.* (1997); Nöth (1987).



Experimental

Crystal data

$\text{C}_{60}\text{H}_{50}\text{B}_2\text{P}_4 \cdot 1.5\text{C}_4\text{H}_8\text{O}$
 $M_r = 1024.66$
 Orthorhombic, $Pbca$
 $a = 19.2421$ (4) Å
 $b = 11.6938$ (2) Å
 $c = 24.9769$ (5) Å

$V = 5620.13$ (19) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.18$ mm⁻¹
 $T = 150$ K
 $0.35 \times 0.28 \times 0.20$ mm

Data collection

Stoe IPDS II diffractometer
 Absorption correction: numerical
 ($X\text{-SHAPE}$ and $X\text{-RED32}$; Stoe & Cie, 2005)
 $T_{\min} = 0.927$, $T_{\max} = 0.986$

91233 measured reflections
 6705 independent reflections
 4392 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.072$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.096$
 $S = 0.87$
 6705 reflections
 343 parameters

9 restraints
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.64$ e Å⁻³
 $\Delta\rho_{\min} = -0.29$ e Å⁻³

Data collection: $X\text{-AREA}$ (Stoe & Cie, 2005); cell refinement: $X\text{-AREA}$; data reduction: $X\text{-RED32}$ (Stoe & Cie, 2005); program(s) used to solve structure: $SHELXS97$ (Sheldrick, 2008); program(s) used to refine structure: $SHELXL97$ (Sheldrick, 2008); molecular graphics: XP in $SHELXTL$ (Sheldrick, 2008); software used to prepare material for publication: $SHELXTL$.

This work was supported by the Leibniz-Institut für Katalyse e.V. an der Universität Rostock.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YK2048).

References

- Bartlett, R. U., Dias, H. V. R. & Power, P. P. (1988). *Inorg. Chem.* **27**, 3919–3922.
 Coates, G. E. & Livingstone, J. G. (1961). *J. Chem. Soc.* pp. 5053–5055.
 Herdtweck, E., Jäcke, F. & Wagner, M. (1997). *Organometallics*, **16**, 4737–4745.
 Kaufmann, B., Jetzfellner, R., Leissring, E., Issleib, K., Nöth, H. & Schmidt, M. (1997). *Chem. Ber.* **130**, 1677–1692.
 Nöth, H. (1987). *Z. Anorg. Allg. Chem.* **555**, 79–84.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Stoe & Cie (2005). $X\text{-SHAPE}$, $X\text{-RED32}$ and $X\text{-AREA}$. Stoe & Cie, Darmstadt, Germany.

supporting information

Acta Cryst. (2012). E68, o1125 [https://doi.org/10.1107/S1600536812011361]

2,4-Bis(diphenylphosphanyl)-1,1,2,3,3,4-hexaphenyl-1,3-diphospha-2,4-dibora-cyclobutane tetrahydrofuran sesquisolvate

Normen Peulecke, Bernd H. Müller, Anke Spannenberg and Uwe Rosenthal

S1. Comment

We became interested in such a class of compounds, because the boron-bridged bisphosphines could be potential ligands for the chromium catalyzed selective oligomerization of ethene, like PNP. Unfortunately the synthesis of $\text{PhB}(\text{PPh}_2)_2$, according to the literature (Coates & Livingstone, 1961), failed and always ended up with insoluble polymer. Therefore we changed the procedure using LiPPh_2 instead of HPPh_2 . Examples of structurally characterized borone-bridged bisphosphines are known (Herdtweck *et al.*, 1997; Kaufmann *et al.*, 1997; Nöth, 1987). Only bulky substituents at the boron lead to a monomeric structure (Bartlett *et al.*, 1988). In the present publication, we report on the formation of the dimeric $\text{C}_{60}\text{H}_{50}\text{B}_2\text{P}_4$. In the structure of the title compound, the diphosphadiborane molecule occupies the position at an inversion center, whereas the solvent molecule of tetrahydrofuran lies in general position with partial occupancy equal to 0.75. In the tetrahydrofuran molecule the O atom is disordered over two sites with occupancies of 0.341 (9): 0.409 (9). All P—B distances in the four-membered ring are essentially identical [B1—P2 = 2.030 (3) Å and B1—P2ⁱ = 2.036 (2) Å], and also the B1—P1 bond distance of 2.043 (2) Å is not significantly different. In the B_2P_2 ring, angles of nearly 90° were observed [P2—B1—P2ⁱ = 87.24 (9)° and B1—P2—B1ⁱ = 92.75°].

S2. Experimental

PhBCl_2 (0.817 ml, 6.3 mmol) was added to a solution of 25 mL Ph_2PLi (0.5M in thf) in 20 ml of thf at -40°C and the resulting solution was stirred at room temperature for 48 h. Subsequently, the formed light brown solution was filtered, reduced to the half, over-layered with *n*-hexane and stored at 0°C. Crystals of the title compound appeared, which were suitable for crystal structure analysis. The white compound was fully characterized by standard analytical methods *e.g.* ^{31}P -NMR: (C_6D_6): -10.7(br), -42.2(tr) p.p.m..

S3. Refinement

H atoms were placed in idealized positions with $d(\text{C—H}) = 0.95$ Å (CH), 0.99 Å (CH_2) and refined using a riding model with $U_{\text{iso}}(\text{H})$ fixed at 1.2 $U_{\text{eq}}(\text{C})$.




Figure 1

The structure of the diphosphadiborane molecule showing the atom-labelling scheme. Hydrogen atoms are omitted for clarity. Displacement ellipsoids are drawn at the 30% probability level.

2,4-Bis(diphenylphosphanyl)-1,1,2,3,3,4-hexaphenyl- 1,3-diphospha-2,4-diboracyclobutane tetrahydrofuran sesquisolvate

Crystal data

$C_{60}H_{50}B_2P_4 \cdot 1.5C_4H_8O$

$M_r = 1024.66$

Orthorhombic, *Pbca*

$a = 19.2421$ (4) Å

$b = 11.6938$ (2) Å

$c = 24.9769$ (5) Å

$V = 5620.13$ (19) Å³

$Z = 4$

$F(000) = 2160$

$D_x = 1.211$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6569 reflections

$\theta = 1.6\text{--}28.0^\circ$

$\mu = 0.18$ mm⁻¹

$T = 150$ K

Prism, colourless

$0.35 \times 0.28 \times 0.20$ mm

Data collection

Stoe IPDS II

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: numerical

(*X-SHAPE* and *X-RED32*; Stoe & Cie, 2005)

$T_{\min} = 0.927$, $T_{\max} = 0.986$

91233 measured reflections

6705 independent reflections

4392 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.072$
 $\theta_{\text{max}} = 28.0^\circ$, $\theta_{\text{min}} = 2.2^\circ$
 $h = -25 \rightarrow 25$

$k = -15 \rightarrow 15$
 $l = -32 \rightarrow 32$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.096$
 $S = 0.87$
 6705 reflections
 343 parameters
 9 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0536P)^2 + 0.0P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.002$
 $\Delta\rho_{\text{max}} = 0.64 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.29 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1A	0.1872 (2)	0.6123 (6)	0.2276 (2)	0.061 (2)*	0.341 (9)
O1B	0.1940 (2)	0.5614 (5)	0.24577 (17)	0.0613 (19)*	0.409 (9)
C31	0.2198 (2)	0.5279 (4)	0.1944 (2)	0.0981 (17)	0.75
H31A	0.1909	0.5086	0.1629	0.118*	0.341 (9)
H31B	0.2305	0.4572	0.2147	0.118*	0.341 (9)
H31C	0.1834	0.5419	0.1672	0.118*	0.409 (9)
H31D	0.2297	0.4448	0.1948	0.118*	0.409 (9)
C32	0.2832 (2)	0.5897 (4)	0.17909 (16)	0.0913 (16)	0.75
H32A	0.2749	0.6378	0.1471	0.110*	0.75
H32B	0.3214	0.5356	0.1711	0.110*	0.75
C33	0.3004 (2)	0.6610 (4)	0.22580 (13)	0.0766 (13)	0.75
H33A	0.3059	0.7420	0.2151	0.092*	0.75
H33B	0.3441	0.6346	0.2427	0.092*	0.75
C34	0.24158 (17)	0.6479 (4)	0.26295 (13)	0.0627 (10)	0.75
H34A	0.2513	0.5891	0.2905	0.075*	0.341 (9)
H34B	0.2301	0.7211	0.2808	0.075*	0.341 (9)
H34C	0.2595	0.6279	0.2989	0.075*	0.409 (9)
H34D	0.2167	0.7217	0.2659	0.075*	0.409 (9)
C1	1.06155 (9)	−0.01162 (16)	0.34268 (7)	0.0246 (4)	
C2	1.00019 (10)	0.02724 (18)	0.31988 (7)	0.0307 (4)	
H2	0.9719	0.0795	0.3392	0.037*	
C3	0.97947 (11)	−0.00883 (19)	0.26950 (8)	0.0364 (5)	

H3	0.9371	0.0185	0.2547	0.044*
C4	1.01996 (11)	−0.0841 (2)	0.24080 (8)	0.0369 (5)
H4	1.0057	−0.1089	0.2063	0.044*
C5	1.08146 (11)	−0.12344 (19)	0.26268 (8)	0.0369 (5)
H5	1.1099	−0.1749	0.2430	0.044*
C6	1.10164 (10)	−0.08786 (17)	0.31321 (7)	0.0308 (4)
H6	1.1437	−0.1161	0.3280	0.037*
C7	1.13848 (10)	0.16430 (16)	0.39173 (7)	0.0257 (4)
C8	1.20183 (10)	0.18941 (18)	0.41622 (7)	0.0299 (4)
H8	1.2214	0.1365	0.4408	0.036*
C9	1.23674 (11)	0.29054 (19)	0.40521 (9)	0.0376 (5)
H9	1.2794	0.3070	0.4228	0.045*
C10	1.20972 (12)	0.36709 (19)	0.36884 (9)	0.0414 (5)
H10	1.2332	0.4369	0.3617	0.050*
C11	1.14820 (11)	0.34169 (18)	0.34279 (9)	0.0383 (5)
H11	1.1302	0.3933	0.3169	0.046*
C12	1.11277 (10)	0.24195 (18)	0.35416 (8)	0.0319 (4)
H12	1.0704	0.2259	0.3362	0.038*
C13	0.98211 (9)	0.19751 (16)	0.43708 (7)	0.0240 (4)
C14	1.01833 (11)	0.29722 (16)	0.45140 (7)	0.0304 (4)
H14	1.0579	0.2909	0.4740	0.036*
C15	0.99832 (12)	0.40388 (18)	0.43376 (9)	0.0390 (5)
H15	1.0236	0.4696	0.4448	0.047*
C16	0.94210 (12)	0.41587 (19)	0.40029 (9)	0.0427 (6)
H16	0.9291	0.4892	0.3874	0.051*
C17	0.90491 (12)	0.32050 (19)	0.38570 (8)	0.0382 (5)
H17	0.8659	0.3282	0.3627	0.046*
C18	0.92386 (10)	0.21293 (17)	0.40430 (7)	0.0280 (4)
H18	0.8967	0.1484	0.3946	0.034*
C19	0.86269 (9)	−0.06475 (16)	0.46070 (6)	0.0237 (4)
C20	0.82663 (9)	0.03413 (16)	0.47541 (7)	0.0263 (4)
H20	0.8517	0.1014	0.4844	0.032*
C21	0.75461 (10)	0.03493 (18)	0.47697 (8)	0.0314 (4)
H21	0.7306	0.1022	0.4876	0.038*
C22	0.71758 (10)	−0.06199 (19)	0.46314 (8)	0.0344 (4)
H22	0.6682	−0.0612	0.4639	0.041*
C23	0.75253 (10)	−0.15974 (18)	0.44826 (8)	0.0328 (4)
H23	0.7271	−0.2259	0.4381	0.039*
C24	0.82462 (10)	−0.16246 (17)	0.44795 (7)	0.0280 (4)
H24	0.8482	−0.2313	0.4390	0.034*
C25	0.98791 (10)	−0.18291 (16)	0.42454 (7)	0.0243 (4)
C26	0.95519 (11)	−0.21012 (17)	0.37607 (7)	0.0295 (4)
H26	0.9138	−0.1714	0.3660	0.035*
C27	0.98292 (11)	−0.29323 (18)	0.34282 (8)	0.0359 (5)
H27	0.9600	−0.3124	0.3104	0.043*
C28	1.04372 (12)	−0.34830 (17)	0.35668 (8)	0.0385 (5)
H28	1.0624	−0.4054	0.3338	0.046*
C29	1.07740 (11)	−0.32094 (17)	0.40355 (8)	0.0337 (5)

H29	1.1197	−0.3581	0.4126	0.040*
C30	1.04967 (10)	−0.23923 (16)	0.43754 (7)	0.0268 (4)
H30	1.0729	−0.2213	0.4700	0.032*
B1	1.01311 (10)	0.07717 (18)	0.45461 (7)	0.0212 (4)
P1	1.09614 (2)	0.02764 (4)	0.409099 (17)	0.02250 (11)
P2	0.95699 (2)	−0.06707 (4)	0.467327 (17)	0.02053 (10)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C31	0.079 (3)	0.089 (4)	0.126 (4)	−0.017 (3)	−0.022 (3)	−0.053 (3)
C32	0.098 (4)	0.128 (4)	0.048 (2)	−0.003 (3)	0.015 (2)	−0.035 (3)
C33	0.081 (3)	0.080 (3)	0.069 (3)	−0.027 (2)	0.015 (2)	−0.019 (2)
C34	0.048 (2)	0.089 (3)	0.0510 (19)	−0.013 (2)	0.0032 (16)	−0.0253 (19)
C1	0.0280 (9)	0.0259 (10)	0.0200 (8)	−0.0036 (8)	0.0021 (7)	0.0005 (7)
C2	0.0311 (10)	0.0355 (11)	0.0256 (9)	0.0051 (9)	−0.0015 (8)	−0.0013 (8)
C3	0.0369 (11)	0.0464 (13)	0.0260 (9)	0.0042 (10)	−0.0062 (8)	0.0006 (9)
C4	0.0420 (12)	0.0480 (13)	0.0207 (9)	−0.0042 (10)	−0.0003 (8)	−0.0058 (9)
C5	0.0399 (12)	0.0422 (13)	0.0285 (10)	0.0029 (10)	0.0061 (8)	−0.0082 (9)
C6	0.0287 (10)	0.0343 (11)	0.0294 (9)	0.0007 (9)	0.0010 (8)	−0.0022 (8)
C7	0.0261 (9)	0.0268 (10)	0.0243 (9)	−0.0006 (8)	0.0043 (7)	−0.0015 (7)
C8	0.0293 (10)	0.0337 (11)	0.0267 (9)	−0.0032 (8)	0.0019 (8)	0.0003 (8)
C9	0.0321 (11)	0.0420 (12)	0.0387 (11)	−0.0092 (9)	0.0044 (9)	−0.0064 (10)
C10	0.0395 (12)	0.0311 (12)	0.0535 (13)	−0.0086 (10)	0.0155 (10)	0.0006 (10)
C11	0.0370 (12)	0.0322 (12)	0.0458 (12)	0.0028 (9)	0.0100 (10)	0.0113 (9)
C12	0.0271 (10)	0.0330 (11)	0.0355 (10)	−0.0004 (8)	0.0028 (8)	0.0068 (9)
C13	0.0284 (9)	0.0237 (9)	0.0199 (8)	0.0006 (8)	0.0053 (7)	0.0009 (7)
C14	0.0389 (11)	0.0246 (10)	0.0276 (9)	−0.0031 (9)	0.0080 (8)	0.0011 (8)
C15	0.0532 (13)	0.0237 (11)	0.0401 (11)	−0.0005 (10)	0.0192 (10)	0.0018 (9)
C16	0.0525 (13)	0.0302 (12)	0.0455 (12)	0.0149 (10)	0.0232 (10)	0.0159 (10)
C17	0.0359 (11)	0.0462 (13)	0.0324 (10)	0.0167 (10)	0.0111 (9)	0.0150 (9)
C18	0.0290 (10)	0.0321 (10)	0.0229 (8)	0.0056 (8)	0.0054 (7)	0.0043 (8)
C19	0.0248 (8)	0.0265 (9)	0.0198 (8)	−0.0023 (8)	−0.0019 (7)	0.0022 (7)
C20	0.0278 (9)	0.0252 (10)	0.0257 (8)	−0.0013 (8)	−0.0036 (7)	0.0019 (7)
C21	0.0294 (10)	0.0317 (11)	0.0331 (10)	0.0033 (8)	−0.0045 (8)	0.0013 (8)
C22	0.0230 (9)	0.0424 (12)	0.0379 (10)	−0.0015 (9)	−0.0057 (8)	0.0018 (9)
C23	0.0290 (10)	0.0330 (11)	0.0363 (10)	−0.0093 (9)	−0.0073 (8)	0.0008 (9)
C24	0.0289 (10)	0.0267 (10)	0.0285 (9)	−0.0021 (8)	−0.0035 (8)	0.0012 (8)
C25	0.0300 (10)	0.0207 (9)	0.0221 (8)	−0.0051 (8)	0.0038 (7)	−0.0002 (7)
C26	0.0335 (10)	0.0294 (10)	0.0255 (9)	−0.0058 (9)	0.0008 (8)	−0.0008 (7)
C27	0.0471 (13)	0.0333 (11)	0.0271 (9)	−0.0123 (10)	0.0023 (9)	−0.0075 (8)
C28	0.0524 (13)	0.0265 (11)	0.0367 (11)	−0.0033 (10)	0.0130 (10)	−0.0102 (9)
C29	0.0394 (11)	0.0244 (10)	0.0375 (11)	0.0005 (8)	0.0080 (9)	0.0003 (8)
C30	0.0306 (10)	0.0236 (10)	0.0263 (9)	−0.0034 (8)	0.0037 (8)	0.0015 (7)
B1	0.0235 (9)	0.0220 (10)	0.0182 (9)	−0.0021 (8)	−0.0006 (7)	0.0005 (7)
P1	0.0239 (2)	0.0231 (2)	0.0206 (2)	0.0002 (2)	−0.00034 (18)	0.00079 (18)
P2	0.0229 (2)	0.0195 (2)	0.01925 (19)	−0.00212 (19)	−0.00134 (18)	−0.00060 (17)

Geometric parameters (Å, °)

O1A—C34	1.431 (3)	C12—H12	0.9500
O1A—C31	1.433 (4)	C13—C18	1.400 (3)
O1B—C34	1.429 (3)	C13—C14	1.405 (3)
O1B—C31	1.430 (4)	C13—B1	1.590 (3)
C31—C32	1.470 (3)	C14—C15	1.378 (3)
C31—H31A	0.9900	C14—H14	0.9500
C31—H31B	0.9900	C15—C16	1.374 (3)
C31—H31C	0.9900	C15—H15	0.9500
C31—H31D	0.9900	C16—C17	1.374 (3)
C32—C33	1.471 (3)	C16—H16	0.9500
C32—H32A	0.9900	C17—C18	1.390 (3)
C32—H32B	0.9900	C17—H17	0.9500
C33—C34	1.472 (3)	C18—H18	0.9500
C33—H33A	0.9900	C19—C24	1.394 (3)
C33—H33B	0.9900	C19—C20	1.398 (3)
C34—H34A	0.9900	C19—P2	1.8222 (18)
C34—H34B	0.9900	C20—C21	1.386 (3)
C34—H34C	0.9900	C20—H20	0.9500
C34—H34D	0.9900	C21—C22	1.383 (3)
C1—C2	1.387 (3)	C21—H21	0.9500
C1—C6	1.390 (3)	C22—C23	1.377 (3)
C1—P1	1.8455 (18)	C22—H22	0.9500
C2—C3	1.386 (3)	C23—C24	1.388 (3)
C2—H2	0.9500	C23—H23	0.9500
C3—C4	1.377 (3)	C24—H24	0.9500
C3—H3	0.9500	C25—C30	1.397 (3)
C4—C5	1.382 (3)	C25—C26	1.401 (2)
C4—H4	0.9500	C25—P2	1.8251 (19)
C5—C6	1.385 (3)	C26—C27	1.385 (3)
C5—H5	0.9500	C26—H26	0.9500
C6—H6	0.9500	C27—C28	1.380 (3)
C7—C8	1.395 (3)	C27—H27	0.9500
C7—C12	1.396 (3)	C28—C29	1.376 (3)
C7—P1	1.8454 (19)	C28—H28	0.9500
C8—C9	1.388 (3)	C29—C30	1.385 (3)
C8—H8	0.9500	C29—H29	0.9500
C9—C10	1.377 (3)	C30—H30	0.9500
C9—H9	0.9500	B1—P2	2.028 (2)
C10—C11	1.383 (3)	B1—P2 ⁱ	2.0361 (18)
C10—H10	0.9500	B1—P1	2.045 (2)
C11—C12	1.381 (3)	P2—B1 ⁱ	2.0363 (18)
C11—H11	0.9500		
C34—O1A—C31	103.7 (3)	C9—C10—H10	120.2
C34—O1B—C31	104.0 (3)	C11—C10—H10	120.2
O1B—C31—C32	112.7 (3)	C12—C11—C10	120.5 (2)

O1A—C31—C32	100.1 (4)	C12—C11—H11	119.8
O1B—C31—H31A	125.6	C10—C11—H11	119.8
O1A—C31—H31A	111.8	C11—C12—C7	120.84 (19)
C32—C31—H31A	111.8	C11—C12—H12	119.6
O1B—C31—H31B	81.0	C7—C12—H12	119.6
O1A—C31—H31B	111.8	C18—C13—C14	116.06 (17)
C32—C31—H31B	111.8	C18—C13—B1	125.14 (17)
H31A—C31—H31B	109.5	C14—C13—B1	118.58 (16)
O1B—C31—H31C	109.0	C15—C14—C13	122.1 (2)
O1A—C31—H31C	88.5	C15—C14—H14	119.0
C32—C31—H31C	109.0	C13—C14—H14	119.0
H31B—C31—H31C	129.5	C16—C15—C14	120.5 (2)
O1B—C31—H31D	109.0	C16—C15—H15	119.8
O1A—C31—H31D	138.9	C14—C15—H15	119.8
C32—C31—H31D	109.0	C15—C16—C17	119.23 (19)
H31A—C31—H31D	83.9	C15—C16—H16	120.4
H31C—C31—H31D	107.8	C17—C16—H16	120.4
C31—C32—C33	105.0 (3)	C16—C17—C18	120.6 (2)
C31—C32—H32A	110.8	C16—C17—H17	119.7
C33—C32—H32A	110.8	C18—C17—H17	119.7
C31—C32—H32B	110.8	C17—C18—C13	121.5 (2)
C33—C32—H32B	110.8	C17—C18—H18	119.3
H32A—C32—H32B	108.8	C13—C18—H18	119.3
C32—C33—C34	105.6 (3)	C24—C19—C20	118.51 (16)
C32—C33—H33A	110.6	C24—C19—P2	122.12 (14)
C34—C33—H33A	110.6	C20—C19—P2	118.87 (14)
C32—C33—H33B	110.6	C21—C20—C19	120.61 (18)
C34—C33—H33B	110.6	C21—C20—H20	119.7
H33A—C33—H33B	108.8	C19—C20—H20	119.7
O1B—C34—C33	112.1 (3)	C22—C21—C20	120.17 (19)
O1A—C34—C33	101.7 (3)	C22—C21—H21	119.9
O1B—C34—H34A	80.7	C20—C21—H21	119.9
O1A—C34—H34A	111.4	C23—C22—C21	119.75 (18)
C33—C34—H34A	111.4	C23—C22—H22	120.1
O1B—C34—H34B	127.1	C21—C22—H22	120.1
O1A—C34—H34B	111.4	C22—C23—C24	120.57 (18)
C33—C34—H34B	111.4	C22—C23—H23	119.7
H34A—C34—H34B	109.3	C24—C23—H23	119.7
O1B—C34—H34C	109.2	C23—C24—C19	120.35 (18)
O1A—C34—H34C	138.2	C23—C24—H24	119.8
C33—C34—H34C	109.2	C19—C24—H24	119.8
H34B—C34—H34C	82.8	C30—C25—C26	118.43 (17)
O1B—C34—H34D	109.2	C30—C25—P2	119.41 (13)
O1A—C34—H34D	86.8	C26—C25—P2	121.88 (15)
C33—C34—H34D	109.2	C27—C26—C25	120.28 (19)
H34A—C34—H34D	130.1	C27—C26—H26	119.9
H34C—C34—H34D	107.9	C25—C26—H26	119.9
C2—C1—C6	117.71 (17)	C28—C27—C26	120.24 (19)

C2—C1—P1	126.48 (14)	C28—C27—H27	119.9
C6—C1—P1	115.81 (14)	C26—C27—H27	119.9
C3—C2—C1	121.20 (19)	C29—C28—C27	120.29 (19)
C3—C2—H2	119.4	C29—C28—H28	119.9
C1—C2—H2	119.4	C27—C28—H28	119.9
C4—C3—C2	120.30 (19)	C28—C29—C30	120.0 (2)
C4—C3—H3	119.9	C28—C29—H29	120.0
C2—C3—H3	119.9	C30—C29—H29	120.0
C3—C4—C5	119.42 (18)	C29—C30—C25	120.69 (18)
C3—C4—H4	120.3	C29—C30—H30	119.7
C5—C4—H4	120.3	C25—C30—H30	119.7
C4—C5—C6	120.05 (19)	C13—B1—P2	125.41 (13)
C4—C5—H5	120.0	C13—B1—P2 ⁱ	114.90 (12)
C6—C5—H5	120.0	P2—B1—P2 ⁱ	87.27 (8)
C5—C6—C1	121.32 (19)	C13—B1—P1	113.00 (12)
C5—C6—H6	119.3	P2—B1—P1	105.52 (9)
C1—C6—H6	119.3	P2 ⁱ —B1—P1	107.17 (9)
C8—C7—C12	117.86 (18)	C7—P1—C1	99.40 (8)
C8—C7—P1	117.71 (14)	C7—P1—B1	103.32 (8)
C12—C7—P1	124.38 (15)	C1—P1—B1	106.76 (8)
C9—C8—C7	121.03 (19)	C19—P2—C25	106.41 (9)
C9—C8—H8	119.5	C19—P2—B1	120.25 (8)
C7—C8—H8	119.5	C25—P2—B1	110.61 (8)
C10—C9—C8	120.13 (19)	C19—P2—B1 ⁱ	111.67 (8)
C10—C9—H9	119.9	C25—P2—B1 ⁱ	115.19 (8)
C8—C9—H9	119.9	B1—P2—B1 ⁱ	92.73 (8)
C9—C10—C11	119.6 (2)		

Symmetry code: (i) $-x+2, -y, -z+1$.