

LETTER • OPEN ACCESS

Implications of climate mitigation for future agricultural production

To cite this article: Christoph Müller *et al* 2015 *Environ. Res. Lett.* **10** 125004

View the [article online](#) for updates and enhancements.

Recent citations

- [LPJmL4 – a dynamic global vegetation model with managed land – Part 1: Model description](#)
Sibyll Schaphoff *et al*
- [Climate and development: enhancing impact through stronger linkages in the implementation of the Paris Agreement and the Sustainable Development Goals \(SDGs\)](#)
Luis Gomez-Echeverri
- [Adaptation of global land use and management intensity to changes in climate and atmospheric carbon dioxide](#)
Peter Alexander *et al*

Environmental Research Letters



LETTER

Implications of climate mitigation for future agricultural production

OPEN ACCESS

RECEIVED
19 February 2015

REVISED
10 November 2015

ACCEPTED FOR PUBLICATION
11 November 2015

PUBLISHED
25 November 2015

Content from this work
may be used under the
terms of the [Creative
Commons Attribution 3.0
licence](#).

Any further distribution of
this work must maintain
attribution to the
author(s) and the title of
the work, journal citation
and DOI.



Christoph Müller¹, Joshua Elliott^{2,3}, James Chryssanthacopoulos³, Delphine Deryng^{4,5}, Christian Folberth⁶, Thomas A M Pugh⁷ and Erwin Schmid⁸

¹ Potsdam Institute for Climate Impact Research, Potsdam, Germany

² University of Chicago Computation Institute, Chicago, IL, USA

³ Columbia University Center for Climate Systems Research, New York, NY, USA

⁴ School of Environmental Sciences, University of East Anglia, Norwich, UK

⁵ Grantham Research Institute on Climate Change and the Environment, London School of Economics and Political Science, London, UK

⁶ International Institute for Applied Systems Analysis, Laxenburg, Austria

⁷ Karlsruhe Institute of Technology, IMK-IFU, Garmisch-Partenkirchen, Germany

⁸ University of Natural Resources and Life Sciences, Vienna, Austria

E-mail: cmueller@pik-potsdam.de and jelliott@ci.uchicago.edu

Keywords: climate change, climate mitigation, agriculture, crop model, AgMIP, ISI-MIP, carbon dioxide

Supplementary material for this article is available [online](#)

Abstract

Climate change is projected to negatively impact biophysical agricultural productivity in much of the world. Actions taken to reduce greenhouse gas emissions and mitigate future climate changes, are thus of central importance for agricultural production. Climate impacts are, however, not unidirectional; some crops in some regions (primarily higher latitudes) are projected to benefit, particularly if increased atmospheric carbon dioxide is assumed to strongly increase crop productivity at large spatial and temporal scales. Climate mitigation measures that are implemented by reducing atmospheric carbon dioxide concentrations lead to reductions both in the strength of climate change and in the benefits of carbon dioxide fertilization. Consequently, analysis of the effects of climate mitigation on agricultural productivity must address not only regions for which mitigation is likely to reduce or even reverse climate damages. There are also regions that are likely to see increased crop yields due to climate change, which may lose these added potentials under mitigation action. Comparing data from the most comprehensive archive of crop yield projections publicly available, we find that climate mitigation leads to overall benefits from avoided damages at the global scale and especially in many regions that are already at risk of food insecurity today. Ignoring controversial carbon dioxide fertilization effects on crop productivity, we find that for the median projection aggressive mitigation could eliminate ~81% of the negative impacts of climate change on biophysical agricultural productivity globally by the end of the century. In this case, the benefits of mitigation typically extend well into temperate regions, but vary by crop and underlying climate model projections. Should large benefits to crop yields from carbon dioxide fertilization be realized, the effects of mitigation become much more mixed, though still positive globally and beneficial in many food insecure countries.

Introduction

The discussion and debate around climate mitigation is generally focused on energy mix options, technical and economic feasibility, and associated losses in economic performance measured in percentage of gross domestic product (Kriegler *et al* 2014). Much less attention has been paid to avoided damages through reduced climate change impacts if global warming is

constrained to low levels, such as the 2° target (but see e.g. Warren *et al* 2013).

Recent model intercomparison projects such as the Agricultural Model Intercomparison and Improvement Project (AgMIP) (Rosenzweig *et al* 2013) and the Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP) (Warszawski *et al* 2014) have coordinated simulations of climate change impacts under different future climate projections and

highlighted the significant uncertainty in agricultural impact models, mainly focusing on the high-end emission scenario under the Representative Concentration Pathway (RCP) 8.5 (Moss *et al* 2010, Riahi *et al* 2011). However, the participating crop models also covered a broader set of climate scenarios, including the low-end representative concentration pathway (RCP2.6), representing a potential future world with strong climate mitigation policies (van Vuuren *et al* 2011a).

The objective of this study is to analyze the effects of such strong climate mitigation on biophysical agricultural productivity, which includes both reduced potential benefits from climate change in currently temperature-limited areas (e.g. in higher latitudes) as well as avoided damages in regions that are negatively affected by climate change (e.g. tropics). Hence, we assess the effect of climate mitigation on biophysical agricultural productivity by comparing simulations of the global gridded crop models (GGCMs) as submitted to the ISI-MIP archive (<http://esg.pik-potsdam.de/esgf-web-fe/>) for the end of the 21st century.

While moderate climate change is often found to have positive effects on agricultural productivity in higher latitudes, where agricultural production is often constrained by low temperatures, the tropics are typically projected to experience detrimental effects of climate change on agricultural productivity even at low levels of warming (Rosenzweig and Parry 1994, Funk *et al* 2008, Rosenzweig *et al* 2014). Carbon dioxide (CO₂) fertilization, i.e. the stimulation of photosynthesis in C3 crops (such as wheat, rice and soy) and reduced water requirements in all crops, has been shown to be able to compensate for some part of the detrimental effects of climate change on agricultural productivity, especially if assumed to not to be constrained by down-regulating mechanisms or nutrient limitation (Zavala *et al* 2008, Leakey *et al* 2009, Ribeiro *et al* 2012). The effectiveness of CO₂ fertilization has been assessed in various modeling and experimental studies but still constitutes a major uncertainty in future crop yield projections (Long *et al* 2006, Tubiello *et al* 2007, Ainsworth *et al* 2008) and biosphere modeling in general (Schimel *et al* 2015). This is not only because of the uncertainty in the implementation in the models, but also because experimental studies have identified several mechanisms that respond negatively to CO₂ fertilization and may prevent that CO₂ benefits on photosynthesis can be exploited in the form of higher crop yields (e.g. Ribeiro *et al* 2012). As anthropogenic climate change is mainly driven by increasing atmospheric concentrations of carbon dioxide (abbreviated as [CO₂] in the following), along with other greenhouse gases (Myhre *et al* 2013), climate mitigation does not only cause a decrease in global warming and associated changes in atmospheric circulation and precipitation patterns, but also reductions in the potential benefits of CO₂ fertilization.

We address uncertainty in patterns of climate change and climate sensitivity as well as in crop models and parameterizations by analyzing an ensemble of five General Circulation Models (GCMs) and six GGCMs for RCPs 8.5 and 2.6. Owing to the given data availability, we focus our analysis on simulations where CO₂ fertilization is assumed to be fully effective, for which all 30 GCM × GGCM combinations are available. Simulations with the assumption of ineffective CO₂ fertilization are only available for one GCM (HadGEM2-ES) and all six GGCMs, which we analyze as well to ensure the robustness of our conclusions to the on-going uncertainty regarding the effectiveness of CO₂ fertilization in enhancing crop yields.

Methods

Simulation ensembles

We here consider the broadest set of consistent future crop yield simulations available to date, which is publicly available in the ISI-MIP fast-track data archive (Warszawski *et al* 2014) as described by Rosenzweig *et al* (2014). While Rosenzweig *et al* (2014) focused on the high-end emission scenario (RCP8.5), we here compare these results against a low emission scenario (RCP2.6), which represents strong climate mitigation efforts (van Vuuren *et al* 2011a). This data set covers various sources of uncertainty: (a) the uncertainty in climate change amplitude and patterns as represented by five different GCM implementations of each RCP emission scenario from the CMIP5 data archive (Taylor *et al* 2012); (b) the uncertainty from crop model response to climate change and CO₂ fertilization as represented by six different GGCMs (Rosenzweig *et al* 2014); and (c) the uncertainty from the effectiveness of CO₂ fertilization on agricultural productivity as represented by two contrasting simulations with static and dynamic [CO₂]. The ten future climate change projections have been supplied to crop modelers as global daily fields of bias-corrected weather variables, including daily minimum and maximum temperature, precipitation, shortwave radiation, vapor pressure deficit and others (Hempel *et al* 2013).

The five GCMs considered and bias-corrected in ISI-MIP are: HadGEM2-ES (Jones *et al* 2011), IPSL-CM5A-LR (Dufresne *et al* 2013), MIROC-ESM-CHEM (Watanabe *et al* 2011), GFDL-ESM2M (Dunne *et al* 2013a, 2013b), and NorESM1-M (Bentsen *et al* 2013, Iversen *et al* 2013), and were selected by earliest availability in the CMIP5 data archive (Taylor *et al* 2012). The GCMs translate the greenhouse gas emission pathways (RCP 2.6 and 8.5) into spatial and temporal fields of temperatures, precipitation and other atmospheric variables. These climate scenarios were supplied to crop modelers in bias-corrected form (Hempel *et al* 2013).

Crops and crop models

We consider maize, wheat, rice, and soy for this comparison. These four major staple foods were selected as the top priority crops in the ISI-MIP fast-track, and together account for 57% of all vegetable food calories, 53% of all animal feed calorie supply and 50% of harvested area in 2010 worldwide (FAO 2015). The six GGCMs include site-based models (EPIC (Williams 1995, Izaurralde *et al* 2006), GEPIC (Williams *et al* 1990, Liu *et al* 2007), pDSSAT (Jones *et al* 2003, Elliott *et al* 2014b)) and agro-ecosystem models (LPJ-GUESS (Smith *et al* 2001, Bondeau *et al* 2007, Lindeskog *et al* 2013), LPJmL (Bondeau *et al* 2007, Fader *et al* 2010, Waha *et al* 2012, Schaphoff *et al* 2013), PEGASUS (Deryng *et al* 2011, Deryng *et al* 2014)), see table S1 and Rosenzweig *et al* (2014) for a comprehensive description of models, setups and simulation protocols as well as tables S1 and S2 in the appendix of this paper. The IMAGE-AEZ model, which also contributed data to the ISI-MIP archive, was excluded as not all scenarios considered here are available from that model.

These six crop models differ in the parameters and mechanisms assumed to represent agricultural management, as well as in how plant processes are implemented. pDSSAT, for example, implements CO₂ effects in form of a scaling factor on the radiation use efficiency factor (RUE) that is used to compute net primary productivity as a function of intercepted light energy. PEGASUS, EPIC and GEPIC also scale RUE and additionally scale transpiration efficiency with increasing [CO₂], while LPJ-GUESS and LPJmL explicitly compute stomata conductance (for the exchange of water and CO₂ with the atmosphere) and photosynthesis as constrained by CO₂ availability, light energy and Rubisco activity (see table S2). Management systems are represented by each model's default setting, ranging from a uniform assumption of crops grown under stress-free automatic fertilization everywhere (EPIC) to pixel-specific calibration of reference period yields (PEGASUS). Similarly, sowing dates and crop varieties follow model-specific parameterizations, and can differ substantially. This diversity, however, also reflects the great variability in production systems within regions that is typically simplified to representative systems in simulations.

CO₂ fertilization

We analyze the full set of 30 GGCM × GCM combinations here under the assumption that CO₂ fertilization is fully effective. The central response is reported in terms of the ensemble median. The effectiveness of CO₂ fertilization in the long run and at the scale of agricultural production systems is highly uncertain (Zavala *et al* 2008, Leakey *et al* 2009, Ribeiro *et al* 2012). For this reason, we also consider simulations with fixed [CO₂] (table S2), representing the

assumption that CO₂ fertilization is ineffective at these scales. According to the given data availability in the ISI-MIP data archive for GGCM simulations (Rosenzweig *et al* 2014), we focus on the HadGEM2-ES climate model (Jones *et al* 2011) implementation of the different RCPs for the case of ineffective CO₂ fertilization.

Aggregation

We present results aggregated to FPU as well as globally. For the spatial aggregation we assume static current crop-specific and irrigation system specific harvested areas per simulated half-degree grid cell based on MIRCA2000 (Portmann *et al* 2010). Crop production (prod) is measured in peta calories (Pcal = 10¹⁵ cal) per FPU (or globally) which is computed as the sum over all grid cells p in that spatial unit, multiplying area (area), the fractional use (frac) per crop c and irrigation system i (rainfed, irrigated), the caloric density (cal) per crop and simulated changes in productivity (γ) for each time step t , according to equation (1)

$$\text{prod}_t = \sum_{p=1, c=1, i=1}^n \left(\text{area}_p * \text{frac}_{p,c,i} * \text{cal}_c * \gamma_{t,c,i} \right). \quad (1)$$

Ensemble corrections

In order to make absolute changes comparable between model simulations, we bias-correct simulated crop production of maize, wheat, rice and soy with gridded crop yields from Iizumi *et al* (2014). For that, we interpolate the Iizumi data from their original 1.125° resolution to the GGCM resolution of 0.5° and aggregate these to the level of Food Production Units (an intersection of major river basins and national boundaries; FPU, see figure S1) (Cai and Rosegrant 2002, Kummu *et al* 2010) and global levels. A model-specific correction factor is then derived per FPU to scale GGCM simulations so that simulated FPU-level production averaged over the reference period matches the observed levels. This approach is analogous to the bias-correction commonly applied to GCM results.

The PEGASUS model does not simulate rice. Therefore, when analyzing changes in total calorie production of the four crops, we use the ensemble median of rice simulations from the other five models in place of the missing PEGASUS rice results. In this way, total calorie projections of PEGASUS are not biased compared to the other GGCMs that include rice as a third C3 crop. For more details of this approach see Elliott *et al* (2014a). PEGASUS was omitted in the rice-only analyses presented here.

Metrics

All changes in agricultural production are represented as absolute changes of the bias-corrected 30-year period at the end of the 21st century (2070–2099) with

respect to the baseline period (1980–2009), see equation (2)

$$\Delta y = \text{prod}_{\text{future}} - \text{prod}_{\text{baseline}}. \quad (2)$$

Here $\widetilde{\Delta y}$ depicts the median of all Δy (as computed in equation (2)) from all GGCM \times GCM combinations considered. Even under strong climate change impact scenarios (here represented by the five RCP8.5 scenarios), climate change impacts on agriculture are not uniformly negative and do not necessarily scale unidirectionally with climate change. The high latitudes, which currently experience strong low-temperature limitations, typically profit from climate change, especially if full effectiveness of CO₂ fertilization is assumed (Rosenzweig *et al* 2014). Under climate mitigation, both the impacts of climate change alone, and the combined impacts of climate change and CO₂ fertilization effects, can be not only dampened or fortified versions of the impact under non-mitigated climate change, but they can also differ in sign. That is, crop-region combinations that see positive impacts from climate change (or from the combined climate change and CO₂ fertilization effects) under non-mitigated climate change may see negative impacts under climate mitigation scenarios (or vice versa).

To quantify the effects of climate mitigation on agricultural productivity, we thus need to consider both cases: (a) avoided damage due to mitigation (β) and (b) lost potentials due to mitigation (λ). Avoided damage metrics (β) are computed for all crop-region combinations, which are projected to see net declining productivity under non-mitigated climate change (RCP8.5 scenarios). Lost potentials metrics (λ) are computed for all crop-region combinations which are projected to see net increasing productivity under non-mitigated climate change (RCP8.5 scenarios). These two cases are mutually exclusive and can thus be presented in maps. We represent the metrics in relative form, i.e. percentage of damages avoided and percentage of potentials lost to allow for a direct comparison between FPUs. At the global aggregation, this is, however, weighted by the total production per FPU to avoid overly emphasizing regions with little contribution to overall crop production in the figures shown. We show maps of absolute changes in production per FPU in the appendix (figure S2). The metrics are defined as follows

$$\beta = \left(1 - \frac{\widetilde{\Delta y_{2.6}}}{\widetilde{\Delta y_{8.5}}} \right) * 100\% \text{ if } \widetilde{\Delta y_{8.5}} < 0 \quad (3)$$

and

$$\lambda = \left(\frac{\widetilde{\Delta y_{2.6}}}{\widetilde{\Delta y_{8.5}}} - 1 \right) * 100\% \text{ if } \widetilde{\Delta y_{8.5}} > 0. \quad (4)$$

As such, the avoided damages of mitigation (β) range from $-\infty$ and $+\infty$, with positive values indicating benefits of mitigation from avoided damages

and negative values indicating that mitigation would make negative impacts more negative. Similarly, the lost potentials from mitigation (λ) range from $-\infty$ and $+\infty$, with positive values indicating that the benefit of climate change is greater after mitigation and negative values indicating that the benefit of climate change is reduced by mitigation.

Quantifying uncertainties

In the presentation of results, we focus on the ensemble median of the 30 simulation sets (5GCMs \times 6GGCMs) available for RCP8.5 and for RCP2.6. We dissect the uncertainties in the global climate change impact assessment induced from GGCMs and CO₂ fertilization by conducting an ANOVA analysis using the `aov` function in R (R Development Core Team 2014). To understand the role of CO₂ fertilization, we focus on the HadGEM2-ES scenarios, as data is not available for the other GCMs for the assumption of ineffective CO₂ fertilization.


Results

Impacts of climate change and CO₂ fertilization

Unmitigated climate change, as e.g. represented by the RCP8.5 emission scenario (Riahi *et al* 2011, van Vuuren *et al* 2011b) poses a significant threat to agricultural production, especially in lower latitudes even if CO₂ fertilization is assumed to be fully effective in enhancing yields (figure 1). Of the four individual crops analyzed here, maize (figure 1(B)) shows the strongest negative impact of climate change. As for all C4 crops, the photosynthesis of maize is not directly stimulated by elevated [CO₂]. Higher latitudes are projected to respond positively to climate change and CO₂ fertilization. At the global scale, unmitigated climate change under the RCP8.5 scenario leads to a median reduction in crop production of the four major crops wheat, maize, rice and soy by 94 Pcal, or just less than 2% of total production. Among the full set of GCM \times GGCM combinations with fully effective CO₂ fertilization, the global RCP8.5 impact ranges from a caloric increase by 2565 Pcal to a decrease by 1047 Pcal (blue bars in figure S3).

Climate change mitigation, i.e. substantially lower [CO₂] and other greenhouse gases by 2100, is represented by the RCP2.6 scenario (van Vuuren *et al* 2011a), which peaks in around 2050 at 443 ppmv [CO₂] and declines to 421 ppmv by 2100. Under climate mitigation, agricultural production is not only subject to less severe increases in temperature, but also to less CO₂ fertilization (if assumed to be effective) and circulation patterns and associated changes in precipitation can be different from both current day and RCP8.5 patterns.

Nonetheless, climate change impacts on agricultural production display similar large-scale



patterns, with stronger impacts in the lower latitudes and beneficial effects in the higher latitudes, where crop growth is often temperature limited in at least parts of the year. However, impacts on agricultural productivity are generally less pronounced (figure 2), and may also change sign in some FPUs, both for individual crops (e.g maize in eastern USA) and for overall agricultural productivity (e.g in Europe).

Under strong mitigation efforts (RCP2.6), combined climate change and CO₂ fertilization effects lead to a small median increase in agricultural production at the global scale of 73 Pcal (1%, blue in figure S4), while climate change alone (assuming no CO₂ fertilization) leads to a slight reduction of global agricultural production (red bars in figure S4).

Impacts of GHG mitigation on agricultural productivity

There are many regions that profit from climate mitigation (i.e. lower [CO₂] and less climate change) either because projected damages from climate change on agricultural productivity can be (partially) avoided (green areas in figure 3) or because yield increases under climate mitigation (RCP2.6) are projected to exceed the projected increases in the unmitigated climate change scenarios (RCP8.5, blue in figure 3). These regions often coincide with regions that are under high or extreme risk of food insecurity (Rosen *et al* 2015), such as many African countries and India, but also cover main producer areas, such as the United States of America (figure 2). Other regions see detrimental effects of climate mitigation on




Figure 2. As figure 1 but for RCP 2.6, representing a strong climate mitigation scenario.

agricultural productivity, mainly because yields are projected to increase more strongly under unmitigated climate change (RCP8.5) than under climate mitigation (RCP2.6, red areas in figure 2(a)).


Crop models indicate that climate mitigation can alleviate all negative impacts on agriculture at the global scale and even supply a small bonus of 2% if CO_2 effects can be fully realized (table 1). There are also substantial regional differences between the individual crops (figures 3(B)–(E)). Maize, a C4 crop that sees no direct stimulation of photosynthesis under elevated $[\text{CO}_2]$, displays the strongest overall response to climate change and has the largest potential to profit from climate mitigation, while the other three crops display mixed responses.

Climate mitigation efforts do not only reduce the median damage projections for agricultural productivity, but the sign of changes is also uncertain in projections of crop productivity under climate change and CO_2 fertilization. As such, climate mitigation (i.e.

RCP2.6 versus RCP8.5) also makes changes to lower agricultural productivity ($\Delta y_{\text{future}} < 0$) less likely (figure 4 (A) versus (B)).

Uncertainty in future projections of agricultural productivity

At the global level, positive and negative impacts in the different regions compensate each other and global change impacts are thus projected to be relatively small for the median case. However, the simulation results of changes in global agricultural productivity are subject to large uncertainties from differences in RCPs, GCMs and GGCMs. With assumed full effectiveness in CO_2 fertilization, total agricultural productivity on currently cultivated areas increases by 112 Pcal from 1980–2009 to 2070–2099 under climate mitigation (RCP 2.6, figure S4) and decreases by about 37 Pcal under unmitigated climate change (RCP 8.5, figure S3).




If CO₂ fertilization is assumed to be ineffective, crop production in almost all regions profits from climate mitigation (figure S5).

If CO₂ fertilization is assumed to be ineffective, projections of global crop productivity are subject to uncertainties from the emission pathway (RCP2.6 versus RCP8.5) and the GGCM used. The role of different GGCMs cannot be investigated here, so for this analysis is based on HadGEM2-ES only. In this setting, the RCP determines the amplitude of climate change while CO₂ concentrations are assumed to be constant. We find that the GGCMs contribute relatively little to overall uncertainty, except for maize (56% of overall variance), whereas the emission pathway (RCP) and thus the strength of climate change strongly affects the

Table 1. ANOVA results for HadGEM2-ES, assuming no CO₂ fertilization, describing the shares of overall variance explained by the GGCMs, the RCPs and their interaction in percent. The standard deviation (Pcal) indicates how variable projections are across RCPs and GGCMs.

	All crops	Maize	Wheat	Rice	Soy
GGCM share (%)	29.6	56.3	15.5	17.5	21.9
RCP share (%)	66.9	36.4	71.8	80.9	74.8
GGCM × RCP interaction share (%)	3.5	7.3	12.8	1.6	3.2
Standard deviation (Pcal)	751	393	192	158	85



projected changes in agricultural productivity (table 1).

If CO₂ fertilization is assumed to be fully effective (table 2), the relative shares are reversed (except for maize). The RCP-induced variance becomes significantly smaller across all crops and the GGCM-induced variance becomes dominant. Also the GGCM \times RCP interaction contributes larger shares to the overall variance, which is higher when CO₂ fertilization is included for the combination of all four crops, and rice and soy separately, but lower for maize and wheat.

If considering the three sources of uncertainty CO₂ fertilization, RCP and GGCM in the HadGEM2-ES scenario ensemble ($n = 24$), the attribution of variance to the different drivers becomes more mixed. Interaction of the different sources of uncertainty can make up to 38% of overall uncertainty (rice, table S3). Across the different FPUs, this pattern is largely consistent, although shares vary across FPUs, but not necessarily in clear patterns (figure S6).

Without CO₂ effects, climate change impacts are more severe but can be alleviated significantly (avoiding damage of ~ 1200 Pcal or 81%) by climate mitigation at the global scale. While the largest uncertainty is clearly the magnitude of CO₂ fertilization, which often changes the sign of climate change impacts in many regions (figure S7). GCMs also contribute to overall uncertainty despite the use of bias correction (table S4), but their contribution is relatively small compared to the contribution of GGCMs. The effect of GCM uncertainty cannot be directly compared to the effect of CO₂ uncertainty, as the two assumptions on CO₂ fertilization are only available for one GCM (HadGEM2-ES). Across the different FPUs, the GGCM uncertainty is typically dominant (compared to the GCM- and RCP-induced uncertainty), but there are

Table 2. As table 1, but assuming full effectiveness of CO₂ fertilization.

	All crops	Maize	Wheat	Rice	Soy
GGCM share (%)	80.9	72.1	79.3	75.1	82.3
RCP share (%)	1.5	15.1	0.6	1.6	0.1
GGCM \times RCP interaction share (%)	17.6	12.8	20.0	23.4	17.6
Standard Deviation (Pcal)	822	357	143	303	105

various FPUs, where the GGCM-induced uncertainty contributes a third or less to overall uncertainty, especially for many wheat areas (figure S8).

Discussion

Data caveats

We use data of the ISI-MIP fast-track data archive, which is, because of its comprehensiveness and global coverage, the best data base available for studying the effects climate mitigation on agricultural productivity. However, the available data is not as comprehensive as desirable and is subject to some constrictions that need to be considered in the interpretation.

The general assumption in the ISI-MIP fast-track data is that agricultural systems do not adapt to climate change, often implemented in the GGCMs as static sowing dates and static parameters for crop varieties and management, although GEPIC, PEGASUS and LPJ-GUESS allow adaptation of some of these variables (Rosenzweig *et al* 2014). As such, the adaptive capacity in the flexibility of agricultural production (e.g. cropping seasons, soil management, varieties) and the continuous efforts to develop better

technological means (e.g. breeding new varieties) are likely substantially underestimated in the analyzed data set.

Many aspects that were found to significantly affect agricultural production are currently not considered in these simulations, such as direct heat stress (Teixeira *et al* 2013, Deryng *et al* 2014, Siebert *et al* 2014), ozone damage (Fuhrer 2003, Hatfield *et al* 2011, Pleijel and Uddling 2012), altered pressure from weeds, pests and diseases (Dermody *et al* 2008, Zavala *et al* 2008, Hatfield *et al* 2011). A notable exception is the PEGASUS model (Deryng *et al* 2011, Deryng *et al* 2014), which explicitly accounts for heat stress and which typically projects more significant reductions in agricultural productivity than the rest of the model ensemble (Rosenzweig *et al* 2014).

Also, we assume static current day cropping patterns (Portmann *et al* 2010) in the analysis, while also adjustments in land-use patterns would lead to reduced climate change impacts (Nelson *et al* 2014a, 2014b). The present analysis thus represents a comparison of the climate change pressure on the adaptive capacity of the agricultural sector rather than a projection of how well agricultural production systems will perform under climate change. The effects of climate mitigation as analyzed here are thus also indicative of how production challenges can be reduced through reduced GHG emissions and subsequent reduced climate change.

Drivers of uncertainty

There is large variance across the different projections of climate change impacts on agricultural production, constituting the uncertainty in these projections. The attribution of this uncertainty to the different drivers is complicated by incomplete data coverage as the assumption on ineffective CO₂ fertilization was only simulated for one GCM (HadGEM2-ES). The overall variance (shown as the standard deviation in tables 1, 2 and S3, which is the square root of the variance but is expressed in the same unit as the data, Pcal) increases once CO₂ fertilization effects are included, except for maize and wheat. However, the attribution is mostly reversed under the opposite assumptions on the effectiveness of CO₂ fertilization. The importance of the emission path (RCP) for agricultural productivity is high if no CO₂ fertilization is assumed, because climate change is the only driver and the 2 RCPs compared are very different. However, once CO₂ fertilization is included, it compensates or even overcompensates much of the climate change impacts, so that the two RCPs are not that different. The strong climate change scenario (RCP8.5) is combined with strong CO₂ fertilization, whereas the moderate climate change scenario (RCP2.6) is combined with moderate CO₂ fertilization. Whereas the variance over the RCPs declines, variance over the GGCMs increases as new processes (stimulation of photosynthesis in C3 plants,

reduced water consumption in all plants) and feedbacks (e.g. soil moisture) come into play. This increase in GGCM-induced variance is stronger than the reduction in RCP-induced variance for rice, soy and the combination of the four crops, but not for maize and wheat. The increasing contribution of the interaction of RCP and GGCM to overall variance (ranging from 12% for maize to 23% for rice, table 2) compared to the assumption of ineffective CO₂ fertilization (ranging from 2% for rice to 13% for wheat, table 1) indicates that the CO₂ effects are implemented differently in the GGCMs and substantial parts of the overall variance can be attributed to the RCP × GGCM combination. The uncertainty from GCMs in the data sample used here (five bias-corrected GCMs, selected by first availability, see Hempel *et al* 2013) is relatively small (2% for rice, 13% for wheat) if CO₂ effects are considered (appendix table S4). This, however, includes the compensation of climate effects with CO₂ fertilization effects so that the RCP-induced variance is mostly important in the interaction of RCP × GGCM. This indicates that the individual GGCMs respond differently to the strength of climate change and the (partially) compensating effects of CO₂ fertilization. These patterns are variable in space, as there are some regions with greater disagreement between GCM projections and others with very similar projections. Overall, however, the GGCM-induced uncertainty outweighs the other sources of uncertainty.

The strong positive effect of high [CO₂] on simulated crop yields does not only constitute a strong source of uncertainty in the GGCM projections. The assumption on no effectiveness in increasing agricultural yields is justified as there is also substantial uncertainty in the overall effects of CO₂ fertilization. This includes co-limitations from water and nutrients, altered competition with weeds and susceptibility to pests and diseases (Dermody *et al* 2008, Zavala *et al* 2008) and the uncertainty of how well stimulated photosynthesis can actually be translated into higher yields. This includes downregulation of photosynthesis (Ainsworth *et al* 2002, 2008, Long *et al* 2006, Tubiello *et al* 2007), co-limitation through environment-controlled limitations to growth (Fatichi *et al* 2014) or altered chemical composition and subsequent partitioning of plant biomass (Ribeiro *et al* 2012). All these factors are currently not included in crop model simulations other than in aggregate parameters (e.g. scaling factors for RUE, see table S2) and thus render the CO₂ effect estimated here as optimistic. However, as some of these effects can be managed (e.g. pest control) or addressed in breeding strategies (e.g. partitioning) it is impossible to assess the magnitude of these effects. From a nutritional perspective, elevated [CO₂] could also lead to reduced micro-nutrient concentration, possibly posing a threat to nutrition security, especially in less developed regions (Müller *et al* 2014, Myers *et al* 2014).

All six GGCMs have been applied and evaluated in climate change impact studies before (table S1), but the models' performance has not been tested in a consistent evaluation framework. The three ecosystem-type models (LPJ-GUESS, LPJmL, PEGASUS) typically show more pronounced responses than the site-based models (EPIC, GEPIC, pDSSAT). LPJ-GUESS and LPJmL, which often show considerably stronger responses to CO₂ fertilization than the other models, also do not account for changing levels of nutrient limitations under evolving climate and [CO₂]. As such, these two models assume an adaptation to changed conditions through increased fertilizer supply and are therefore not directly comparable to the other models. LPJ-GUESS and PEGASUS also assume adaptation in varieties (growing season length), and PEGASUS and GEPIC also assume adaptation in sowing dates not only within a fixed planting window as EPIC, pDSSAT and LPJ-GUESS do. As such, assumptions on management and on the flexibility of agricultural production systems also constitute a significant source of differences between models' projections. PEGASUS, despite allowing for adaptation in sowing dates and varieties, is typically the most pessimistic (Rosenzweig *et al* 2014).

The diversity in model setups in the ISI-MIP model ensemble employed here with respect to growing seasons or management systems (e.g. potential versus actual yields, explicit nutrient stress, calibrated productivity levels, see table S1) reflects part of the diversity in actual production systems. However, it may also introduce confounding dynamics such as additional water stress if intensity levels are overestimated or if the growing season is not well parameterized. As such, the assumption that model biases can be corrected by a linear correction factor may not hold, but will have to be addressed by improved and harmonized data on management in future simulations (e.g. Elliott *et al* 2015). A global crop model benchmark system is currently under development in AgMIP GGCM (Elliott *et al* 2015) and will facilitate structured model evaluation and subsequent improvements. We thus focus on the model ensemble median here, which has been shown to be a robust estimate for crop model projections (Bassu *et al* 2014, Martre *et al* 2014, Asseng *et al* 2015).

Despite these important qualifiers, GGCM simulations are the best tool available to assess climate change impacts and the role of mitigation efforts for agricultural production systems. The large model uncertainty is largely a product of the multiple complex interactions in soils and plants as well as uncertainties in the parameterization of management practices and crop varieties. This highlights the value of such complex models and the diversity of these in the present modeling ensemble. The GGCM Intercomparison (Elliott *et al* 2015) of the AgMIP (Rosenzweig *et al* 2013) has set out to better assess current model skills, identify deficiencies and to improve on

these, where data availability and process understanding allow for it.

Implications of climate mitigation for agricultural productivity

Agricultural productivity under given management is driven by various aspects of climate change. This includes changes in temperatures (min, max, mean), precipitation (amount and temporal distribution), and in cloudiness, which subsequently affects the available energy from radiation and the corresponding evaporative demand of the atmosphere. Atmospheric CO₂ concentrations do not only drive this climate change and but also stimulate photosynthesis in C3 plants (such as wheat, rice and soy) and decrease water requirements. These climate-change driven alterations of growing conditions can thus lead to diverse changes, depending on their direction (e.g. dryer or wetter), amplitude, and the starting conditions (e.g. cold- or heat-limited growing season).

As such, climate mitigation efforts, which lead to lower greenhouse gas concentrations in the atmosphere (largely realized through lower [CO₂]), have mixed effects on agricultural productivity. In summary, climate change and CO₂ fertilization lead to damages to agricultural productivity in the lower latitudes and to new production potentials through higher yields in higher latitudes as well as in water-limited regions where water is used more efficiently by plants (Deryng *et al* under review). Consequently, climate mitigation measures through reduction of [CO₂] typically reduce the damages in warmer regions and leads to losses of new production potentials in cooler regions. Even if impacts are relatively small at the global aggregation, climate mitigation would strongly benefit many less developed and food insecure countries, as e.g. in Africa and Asia (Rosen *et al* 2015), which would counteract the growing asymmetries between developed and developing countries (Fischer *et al* 2005).

Conclusions

Climate change impacts on agricultural production can be greatly reduced by climate mitigation at the global scale and also in most regions for maize, wheat, rice and soy. However, associated reductions in [CO₂] also lead to reduced positive effects in CO₂ fertilization if these can be materialized by farmers in the fields. Based on the ensemble median of five climate models and six GGCMs, we find that climate mitigation has case-specific effects on agricultural productivity. While overall slightly positive at the global aggregation level, individual regions display very different responses to climate mitigation. Agricultural productivity in many regions shows a positive response to climate mitigation, either through avoided damages or through higher yield increases than under the

unmitigated climate change scenarios. These regions include many regions that are currently food insecure (Rosen *et al* 2015) or are major food producers (e.g. USA). Increased agricultural productivity in other regions, partly through warmer temperatures in cold-limited regions but mainly through the projected positive effects of CO₂ fertilization, cannot directly compensate the overall reduction and would also require substantial changes in production patterns globally in order to exploit their potential. The diversity in response across regions and crops to climate mitigation reflects the diversity in climate change impacts. This analysis of biophysical benefits and losses in agricultural productivity through climate mitigation demonstrates the potential and importance of climate mitigation, especially for many less-developed countries. Developing a comprehensive assessment of climate mitigation effects on the agricultural sector will not only require accounting for changes in production and consumption patterns (Nelson *et al* 2014a, 2014b) but also to better understand and project how agricultural productivity can be secured or enhanced through changes in agricultural management, including assessments of the associated economic costs. This task remains a significant challenge and will have to be addressed by global crop modelers also in cooperation agricultural economists, as e.g., in AgMIP.

Acknowledgments

CM acknowledges financial support from the MACMIT project (01LN1317A) funded through the German Federal Ministry of Education and Research (BMBF). The contribution by TP was funded by the European Commission's 7th Framework Programme, under Grant Agreement numbers 282672 (EMBRACE) and 603542 (LUC4C) and was supported, in part, by the German Federal Ministry of Education and Research (BMBF), through the Helmholtz Association and its research program ATMO.

References

- Ainsworth E A *et al* 2002 A meta-analysis of elevated [CO₂] effects on soybean (Glycine max) physiology, growth and yield *Glob. Change Biol.* **8** 695–709
- Ainsworth E A, Leakey A D B, Ort D R and Long S P 2008 FACE-ing the facts: inconsistencies and interdependence among field, chamber and modeling studies of elevated [CO₂] impacts on crop yield and food supply *New Phytologist* **179** 5–9
- Asseng S *et al* 2015 Rising temperatures reduce global wheat production *Nat. Clim. Change* **5** 143–7
- Bassu S *et al* 2014 Do various maize crop models give the same responses to climate change factors? *Glob. Change Biol.* **20** 2301–20
- Bentsen M *et al* 2013 The Norwegian earth system model, NorESM1-M: I. Description and basic evaluation of the physical climate *Geosci. Model Dev.* **6** 687–720
- Bondeau A *et al* 2007 Modelling the role of agriculture for the 20th century global terrestrial carbon balance *Glob. Change Biol.* **13** 679–706
- Cai X M and Rosegrant M W 2002 Global water demand and supply projections: I. A modeling approach *Water Int.* **27** 159–69
- Dermod O, O'Neill B F, Zangerl A R, Berenbaum M R and DeLucia E H 2008 Effects of elevated CO₂ and O₃ on leaf damage and insect abundance in a soybean agroecosystem *Arthropod-Plant Interact.* **2** 125–35
- Deryng D, Conway D, Ramankutty N, Price J and Warren R 2014 Global crop yield response to extreme heat stress under multiple climate change futures *Environ. Res. Lett.* **9** 034011
- Deryng D *et al* Regional disparities in the beneficial effects of rising CO₂ emissions on crop water productivity *Nat. Clim. Change* submitted
- Deryng D, Sacks W J, Barford C C and Ramankutty N 2011 Simulating the effects of climate and agricultural management practices on global crop yield *Glob. Biogeochem. Cycles* **25** GB2006
- Dufresne J L *et al* 2013 Climate change projections using the IPSL-CM5 earth system model: from CMIP3 to CMIP5 *Clim. Dyn.* **40** 2123–65
- Dunne J P *et al* 2013a GFDL's ESM2 global coupled climate-carbon earth system models: I. Physical formulation and baseline simulation characteristics *J. Clim.* **25** 6646–65
- Dunne J P *et al* 2013b GFDL's ESM2 global coupled climate-carbon earth system models: II. Carbon system formulation and baseline simulation characteristics* *J. Clim.* **26** 2247–67
- Elliott J *et al* 2014a Constraints and potentials of future irrigation water availability on agricultural production under climate change *Proc. Natl Acad. Sci.* **111** 3239–44
- Elliott J, Kelly D, Chrysanthacopoulos J, Glotter M, Jhunhnuwala K, Best N, Wilde M and Foster I 2014b The parallel system for integrating impact models and sectors (pSIMS) *Environ. Model. Softw.* **62** 509–16
- Elliott J *et al* 2015 The global gridded crop model intercomparison: data and modeling protocols for phase 1 (v1.0) *Geosci. Model Dev.* **8** 261–77
- Fader M, Rost S, Müller C, Bondeau A and Gerten D 2010 Virtual water content of temperate cereals and maize: present and potential future patterns *J. Hydrol.* **384** 218–31
- FAO 2015 FAOSTAT (<http://faostat.fao.org/>) (accessed 17 February 2015)
- Fatichi S, Leuzinger S and Körner C 2014 Moving beyond photosynthesis: from carbon source to sink-driven vegetation modeling *New Phytologist* **201** 1086–95
- Fischer G, Shah M, Tubiello F N and van Velhuizen H 2005 Socio-economic and climate change impacts on agriculture: an integrated assessment, 1990–2080 *Phil. Trans. R. Soc. B* **360** 2067–83
- Fuhrer J 2003 Agroecosystem responses to combinations of elevated CO₂, ozone, and global climate change *Agric. Ecosyst. Environ.* **97** 1–20
- Funk C, Dettinger M D, Michaelsen J C, Verdin J P, Brown M E, Barlow M and Hoell A 2008 Warming of the Indian Ocean threatens eastern and southern African food security but could be mitigated by agricultural development *Proc. Natl Acad. Sci. USA* **105** 11081–6
- Hatfield J L, Boote K J, Kimball B A, Ziska L H, Izaurralde R C, Ort D, Thomson A M and Wolfe D 2011 Climate impacts on agriculture: implications for crop production *Agronomy J.* **103** 351–70
- Hempel S, Frieler K, Warszawski L, Schewe J and Piontek F 2013 A trend-preserving bias correction—the ISI-MIP approach *Earth Syst. Dyn.* **4** 219–36
- Iizumi T, Yokozawa M, Sakurai G, Travasso M I, Romanerikov V, Oettli P, Newby T, Ishigooka Y and Furuya J 2014 Historical changes in global yields: major cereal and legume crops from 1982 to 2006 *Glob. Ecology Biogeography* **23** 346–57
- Iversen T *et al* 2013 The Norwegian earth system model, NorESM1-M: II. Climate response and scenario projections *Geosci. Model Dev.* **6** 389–415
- Izaurralde R C, Williams J R, McGill W B, Rosenberg N J and Jakas M C Q 2006 Simulating soil C dynamics with EPIC: model description and testing against long-term data *Ecological Model.* **192** 362–84

- Jones C D *et al* 2011 The HadGEM2-ES implementation of CMIP5 centennial simulations *Geosci. Model Dev.* **4** 543–70
- Jones J W, Hoogenboom G, Porter C H, Boote K J, Batchelor W D, Hunt L A, Wilkens P W, Singh U, Gijsman A J and Ritchie J T 2003 The DSSAT cropping system model *Eur. J. Agronomy* **18** 235–65
- Kriegler E *et al* 2014 The role of technology for achieving climate policy objectives: overview of the EMF 27 study on global technology and climate policy strategies *Clim. Change* **123** 353–67
- Kummu M, Ward P J, Moel H D and Varis O 2010 Is physical water scarcity a new phenomenon? Global assessment of water shortage over the last two millennia *Environ. Res. Lett.* **5** 034006
- Leakey A D B, Ainsworth E A, Bernacchi C J, Rogers A, Long S P and Ort D R 2009 Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE *J. Exp. Bot.* **60** 2859–76
- Lindeskog M, Arneth A, Bondeau A, Waha K, Seaquist J, Olin S and Smith B 2013 Implications of accounting for land use in simulations of ecosystem carbon cycling in Africa *Earth Syst. Dyn.* **4** 385–407
- Liu J G, Williams J R, Zehnder A J B and Yang H 2007 GEPIC—modelling wheat yield and crop water productivity with high resolution on a global scale *Agric. Syst.* **94** 478–93
- Long S P, Ainsworth E A, Leakey A D B, Nosberger J and Ort D R 2006 Food for thought: lower-than-expected crop yield stimulation with rising CO₂ concentrations *Science* **312** 1918–21
- Martre P *et al* 2014 Multimodel ensembles of wheat growth: many models are better than one *Glob. Change Biol.* **21** 911–25
- Moss R H *et al* 2010 The next generation of scenarios for climate change research and assessment *Nature* **463** 747–56
- Müller C, Elliott J and Levermann A 2014 Food security: fertilizing hidden hunger *Nat. Clim. Change* **4** 540–1
- Myers S S *et al* 2014 Increasing CO₂ threatens human nutrition *Nature* **510** 139–42
- Myhre G *et al* 2013 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge, UK: Cambridge University Press)
- Nelson G C *et al* 2014a Climate change effects on agriculture: economic responses to biophysical shocks *Proc. Natl Acad. Sci. USA* **111** 3274–9
- Nelson G C *et al* 2014b Agriculture and climate change in global scenarios: why don't the models agree *Agric. Econ.* **45** 85–101
- Pleijel H and Uddling J 2012 Yield versus quality trade-offs for wheat in response to carbon dioxide and ozone *Glob. Change Biol.* **18** 596–605
- Portmann F T, Siebert S and Döll P 2010 MIRCA2000—Global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and hydrological modeling *Glob. Biogeochemical Cycles* **24** Gb1011
- R Development Core Team 2014 *R: A language and Environment for Statistical Computing* (Vienna, Austria: R Foundation for Statistical Computing) (<http://R-project.org/>)
- Riahi K, Rao S, Krey V, Cho C, Chirkov V, Fischer G, Kindermann G, Nakicenovic N and Rafaj P 2011 RCP 8.5—a scenario of comparatively high greenhouse gas emissions *Clim. Change* **109** 33–57
- Ribeiro D M, Araújo W L, Fernie A R, Schippers J H M and Mueller-Roeber B 2012 Action of gibberellins on growth and metabolism of arabidopsis plants associated with high concentration of carbon dioxide *Plant Physiol.* **160** 1781–94
- Rosen S, Meade B and Murray A 2015 *International Food Security Assessment 2015–2025* p 59
- Rosenzweig C *et al* 2013 The agricultural model intercomparison and improvement project (AgMIP): protocols and pilot studies *Agric. Forest Meteorol.* **170** 166–82
- Rosenzweig C *et al* 2014 Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison *Proc. Natl Acad. Sci.* **111** 3268–73
- Rosenzweig C and Parry M L 1994 Potential impact of climate change on world food supply *Nature* **367** 133–8
- Schaphoff S, Heyder U, Ostberg S, Gerten D, Heinke J and Lucht W 2013 Contribution of permafrost soils to the global carbon budget *Environ. Res. Lett.* **8** 014026
- Schimel D, Stephens B B and Fisher J B 2015 Effect of increasing CO₂ on the terrestrial carbon cycle *Proc. Natl Acad. Sci. USA* **112** 436–41
- Siebert S, Ewert F, Rezaei E E, Kage H and Graß R 2014 Impact of heat stress on crop yield—on the importance of considering canopy temperature *Environ. Res. Lett.* **9** 044012
- Smith B, Prentice I C and Sykes M T 2001 Representation of vegetation dynamics in the modelling of terrestrial ecosystems: comparing two contrasting approaches within European climate space *Glob. Ecology Biogeography* **10** 621–37
- Taylor K E, Stouffer R J and Meehl G A 2012 An overview of CMIP5 and the experiment design *Bull. Am. Meteorol. Soc.* **93** 485–98
- Teixeira E I, Fischer G, van Velthuisen H, Walter C and Ewert F 2013 Global hot-spots of heat stress on agricultural crops due to climate change *Agric. Forest Meteorol.* **170** 206–15
- Tubiello F N, Amthor J S, Boote K J, Donatelli M, Easterling W, Fischer G, Gifford R M, Howden M, Reilly J and Rosenzweig C 2007 Crop response to elevated CO₂ and world food supply—a comment on ‘food for thought...’ by Long *et al* *Science* **312**: 1918–1921, 2006 *Euro. J. Agronomy* **26** 215–23
- van Vuuren D *et al* 2011a RCP2.6: exploring the possibility to keep global mean temperature increase below 2 °C *Clim. Change* **109** 95–116
- van Vuuren D P *et al* 2011b The representative concentration pathways: an overview *Clim. Change* **109** 5–31
- Waha K, van Bussel L G J, Müller C and Bondeau A 2012 Climate-driven simulation of global crop sowing dates *Glob. Ecology Biogeography* **21** 247–59
- Warren R *et al* 2013 Quantifying the benefit of early climate change mitigation in avoiding biodiversity loss *Nat. Clim. Change* **3** 678–82
- Warszawski L, Frieler K, Huber V, Piontek F, Serdeczny O and Schewe J 2014 The inter-sectoral impact model intercomparison project (ISI-MIP): project framework *Proc. Natl Acad. Sci. USA* **111** 3228–32
- Watanabe S *et al* 2011 MIROC-ESM 2010: model description and basic results of CMIP5-20c3m experiments *Geosci. Model Dev.* **4** 845–72
- Williams J R 1995 The EPIC model *Computer Models of Watershed Hydrology* ed V P Singh (Littleton, CO: Water Resources Publications) pp 909–1000
- Williams J R, Dyke P T, Fuchs W W, Benson V W, Rice O W and Taylor E D 1990 *EPIC—Erosion/Productivity Impact Calculator* US Department of Agriculture
- Zavala J A, Casteel C L, DeLucia E H and Berenbaum M R 2008 Anthropogenic increase in carbon dioxide compromises plant defense against invasive insects *Proc. Natl Acad. Sci. USA* **105** 5129–33