

LETTER • OPEN ACCESS

## Nighttime light data reveal lack of full recovery after hurricanes in Southern US

To cite this article: Kelsey Barton-Henry and Leonie Wenz 2022 *Environ. Res. Lett.* **17** 114015

View the [article online](#) for updates and enhancements.

You may also like

- [Evaluating nighttime lights and population distribution as proxies for mapping anthropogenic CO<sub>2</sub> emission in Vietnam, Cambodia and Laos](#)  
Andrea E Gaughan, Tomohiro Oda, Alessandro Sorichetta et al.
- [Non-Thickness-Limited Growth of Anodic Oxide Films on Tantalum](#)  
Y.-M. Li and L. Young
- [Monitoring hydropower reliability in Malawi with satellite data and machine learning](#)  
Giacomo Falchetta, Chisomo Kasamba and Simon C Parkinson

ENVIRONMENTAL RESEARCH  
LETTERS

## LETTER

## Nighttime light data reveal lack of full recovery after hurricanes in Southern US

## OPEN ACCESS

RECEIVED  
27 July 2022REVISED  
23 September 2022ACCEPTED FOR PUBLICATION  
12 October 2022PUBLISHED  
24 October 2022

Original content from  
this work may be used  
under the terms of the  
[Creative Commons  
Attribution 4.0 licence](#).

Any further distribution  
of this work must  
maintain attribution to  
the author(s) and the title  
of the work, journal  
citation and DOI.

Kelsey Barton-Henry<sup>1,2</sup> and Leonie Wenz<sup>1,3,\*</sup> <sup>1</sup> Potsdam Institute for Climate Impact Research, Potsdam, Germany<sup>2</sup> Technische Universität, Berlin, Germany<sup>3</sup> Mercator Research Institute on Global Commons and Climate Change, Berlin, Germany

\* Author to whom any correspondence should be addressed.

E-mail: [leonie.wenz@pik-potsdam.de](mailto:leonie.wenz@pik-potsdam.de)**Keywords:** hurricanes, recovery dynamics, nighttime light data, feature importance analysis, quasi-experimental methodsSupplementary material for this article is available [online](#)**Abstract**

As the climate warms, many areas of the world are experiencing more frequent and extreme weather events. Hurricanes carry some of the costliest short-term socioeconomic repercussions via economic losses and people displaced. There is, however, little quantitative evidence regarding medium- to long-term effects, nor factors moderating recovery. Here we show that areas affected by hurricanes of category 4 or 5 in the southern US between 2014 and 2020 generally do not demonstrate full recovery in the longer term. Utilizing Visible Infrared Imaging Radiometer Suite nighttime light (NTL) data as a proxy for economic activity and population density, we build a timeline of recovery via NTL radiance levels. We exploit the difference in the eligibility for aid from the Federal Emergency Management Agency (FEMA) to apply a quasi-experimental method to identify changes in NTL radiance attributable to hurricanes. We find that after three years, affected areas demonstrate a reduction in NTL radiance levels of between 2% and 14% compared to the pre-disaster period. Combining these results with machine learning techniques, we are able to investigate those factors that contribute to recovery. We find counties demonstrating smaller reductions in NTL radiance levels in the months following the hurricane are buoyed by the amount of FEMA aid received, but that this aid does not foster a longer-term return to normal radiance levels. Investigating areas receiving FEMA aid at the household and individual level, we find age and employment are more important than other demographic factors in determining hurricane recovery over time. These findings suggest that aid may be more important in motivating short-term recovery for public entities than for individuals but is not sufficient to guarantee complete recovery in the longer term.

**1. Introduction**

Weather extremes can affect ecological and socioeconomic systems profoundly [1–4]. Hurricanes are among the costliest natural disasters in the United States. Between 2005 and 2020, the total estimated damages from tropical cyclones amounted to \$595 billion; 43% of that total occurred since 2015 [5]. With continued climate change, such events are becoming more intense and frequent in many parts of the world. It is therefore increasingly important to understand the consequences of these events, and our

ability to recover from them, as well as possible linkages to adaptation behaviors such as migration.

Studies on the economic consequences of hurricanes have already suggested the temporal complexity of recovery patterns [6] especially in the context of climate change-adaptive behaviors [7]. There is evidence that out-migration from hurricane-prone areas may already constrain recovery in areas affected by large hurricanes [8]. Most studies have investigated the effects of one or two hurricanes in isolation and over short time lines, often spanning only months, and conclusions of empirical studies on longer-term

effects are ambiguous. Some find that disaster affected areas in both rich and poor countries do not return to pre-disaster income levels even 20 years post-disaster [9] whereas others find no long-lasting significant effects using nighttime light intensity [10–13] or even a rebound in brightness after a disaster [14].

The determinants of post-disaster recovery also remain poorly understood, despite their potential to inform disaster aid policy, motivate recovery, and enable better quantification of the costs of climate change. Studies have pinpointed emergency aid as a mitigator to out-migration from disaster-affected areas [15] and its slow dispersal as an impediment to recovery [16]. For developed countries, post-disaster migration has been shown to be complex, with the variance in migration across affected areas often explained by the presence of disadvantaged populations, lower overall ex-ante levels of development, or rural or urban status [17, 18]. Federal Emergency Management Agency (FEMA) aid may also act as a moderator of post-disaster migration for low-income households [19]. Others find that factors of social vulnerability are less important in motivating migration than the severity of the disaster [9, 20].

We aim to fill both these gaps by investigating the evolution of hurricane response dynamics over time and identifying the comparative importance of socioeconomic, demographic, and aid-related factors that could drive adaptation responses. To this end, we build a combined data set matching indicators on the reception of FEMA post-disaster aid to monthly geolocated visible infrared imaging radiometer suite day/night band nighttime light (VIIRS NTL) data, urbanization data from the National Center for Health Statistics, and county-level demographic and socioeconomic data from the US Census Bureau and Bureau of Economic Analysis for all category 4 and 5 hurricanes which made landfall in the United States between 2014 and 2020 (see data & methods section of the supplementary materials for detail).

VIIRS NTL data may offer crucial insights into storm recovery dynamics across time and socioeconomic conditions, given their highly granular temporal and geographic resolution and significant correlations to both gross domestic product [21–23] and population density [24–27] as well as their ability to effectively capture patterns of urbanization around the world [28, 29]. These data have previously been used to study time trends of light recovery correlated with one or two hurricanes, or over short time scales [25, 30–34].

Here, we study the effects of many hurricanes at once on a monthly basis, and for the oldest storms, over longer time spans than previously studied. We further build on correlative, time-series based work by employing a quasi-experimental method, a difference-in-differences (DiD) model, to estimate the plausibly causal effects of hurricanes on changes

in night light intensity. We then apply tree-based machine learning models in conjunction with the Shapley Additive explanations algorithm (SHAP) to identify those variables most important in explaining differences in hurricane responses across different counties. SHAP is already widely used in combination with tree methodologies to explain endogenous trends in climate-related human behavior [35, 36]. It has been found to produce local feature importance estimations that are both generalizable to the global model, and well in line with human intuition [37, 38].

## 2. Method overview

### 2.1. Data

We build our main dataset by matching monthly VIIRS NTL radiance data [39] and data on Individual and Publicly available FEMA post-hurricane disaster assistance at the county level [40] for seven hurricanes [41]. The FEMA disaster assistance data identifies those counties in which individuals and households (referred to as Individual Aid hereafter), public entities and nonprofits, or both groups (referred to as Combined Aid hereafter), were eligible for disaster-related relief aid from FEMA. We use this information to identify counties affected by each hurricane (the ‘treatment’ group for that hurricane) and unaffected counties in the same state (the ‘control’). We define two treatment groups, evaluated separately, based on the type of FEMA aid for which they were eligible: Individual or Combined aid. Maps of the areas receiving each type of aid as well as unaffected areas can be found in supplementary figures 1–7.

For further controls, supplementary data on rural and urban areas [42], as well as geographic county boundaries [43], and dollar amounts of total aid provided by FEMA, are added to these data [40]. Last, we include socioeconomic [44] and demographic data [45] on the county level. For details on each dataset, please see supplementary materials: Data & Methods, along with a conceptual overview of the methodology employed in supplementary figure 8.

### 2.2. Empirical strategy

In order to gain insight into the potential causal effects of hurricanes on NTL intensity changes, we employ a DiD methodology. This method first assesses the difference in the level of nighttime radiance between the pre-hurricane period and the post-hurricane period for both the group of counties affected and the group of counties unaffected by the hurricane, separately (first difference). It then subtracts the average change in NTL levels in the treatment group from that of the control group (second difference). If the change in radiance after the storm was only assessed for counties affected by the hurricane, the estimation may be biased due to the effect of trends and events other than the hurricane on the

level of NTL. The second difference removes these biases. Given that the same time period is being compared in both the treatment and control groups, the data does not need to be preprocessed to remove seasonality, as seasonal trends affect both the treatment and control. Similarly, if the difference in NTL radiance between treated and untreated counties was only assessed for the post-hurricane period, there could be biases due to already existing differences between the two groups of counties. The first difference removes those biases. Taken together, the two differences yield a suitable counterfactual, thus enabling the estimation of a plausibly causal effect of a hurricane on the level of radiance in the affected counties.

For each hurricane, we run two DiD models for each storm, one for each treatment variable based on the type of FEMA aid received. For all storms, the entire timeline available is used, meaning earlier storms have fewer months of ‘pre-storm’ data entries (with the minimum being 32 months for the earliest storm) and later storms have fewer post-storm data entries (with a minimum of 16 months). As we assess recovery over time, we shorten the timeline of data month by month, relative to when the storm hit.

The difference-in-differences specification follows:

$$Y_{ct} = \beta_0 + \beta_1(\text{post} \times \text{treatment})_{ct} + \beta_2\text{post}_t + \beta_3\text{treatment}_c + \alpha X_c + \varepsilon_{ct};$$

where  $Y_{ct}$  is the log of the NTL radiance level in county  $c$  at time  $t$  (with  $t$  being a month in a year).  $\text{post}_t$  is a binary variable, taking the value of 1 if  $t$  falls in the period after the hurricane (post-period) and 0 if it falls in the period before the storm. The binary  $\text{treatment}_c$  indicates whether county  $c$  is in the treatment group (=1) or in the control group (=0). Accordingly, the interaction term,  $(\text{post} \times \text{treatment})_{ct}$  takes the value 1 only if the outcome was observed in the treatment group and in the post period, and is zero in all other cases. The difference-in-difference effect,  $\beta_1$ , is our coefficient of interest, denoting by how much the difference in NTL radiance between the treatment and control group changed after the hurricane. It can hence be interpreted as the percent change in NTL radiance in the affected counties after the hurricane over the whole post-disaster period up to time  $t$ .  $X_c$  is a vector of economic and demographic controls on the county level  $c$ , and  $\varepsilon_{ct}$  is the idiosyncratic error term on the month and county level. Running the DiD model for different post-disaster time horizons provides us with a recovery time series indicating the plausibly causal effect of a storm on NTL radiance for different time periods.

Our identification strategy rests on the validity of the ‘common trends’ assumption: in the absence of the hurricane, the treatment and control areas would have followed the same trend in nighttime

**Table 1.** Interpretation of the results of the ‘leave-one-out’ DiD method. The leftmost column of table 1 indicates the sign of the estimation resultant from the DiD model. The middle column indicates the sign of difference between that estimation (here, ‘average change’) and the estimation resultant from the DiD model when that county is omitted. The rightmost column provides our interpretation of these signs for the county omitted.

Average change in radiance	(Average change)— (change when county omitted)	Interpretation for county being left out
–	–	More radiance reduction than average
–	+	Less radiance reduction than average (‘better off’)
+	–	Less radiance increase than average
+	+	More radiance increase than average

light radiance [46]. To assess this assertion, we compare NTL radiance trends in both groups prior to the hurricane. We find the treatment and control follow a common trend, suggesting that the divergence in light we see post-hurricane is indeed due to the hurricane. Graphical representations for each hurricane are found in supplementary figures 9–15.

### 2.3. Identifying county-level effects using a leave-one-out method

The DiD methodology requires an aggregate comparison between the treatment and control groups, and the pre- and post-storm periods. The use of this counterfactual between treatment and control ensures that trends affecting both groups are differentiated out, strengthening the inference of causality. As a consequence, changes in NTL radiance can only be assessed as an average across all counties and not at the individual county level. In order to gain insights into a hurricane’s effects at the county level and possible differences between counties, we combine the DiD method with a ‘leave-one-out’ approach. Specifically, we iteratively re-estimate the DiD model, excluding one county at a time, for the full time period after the storm available for each storm. After subtracting these estimations from the DiD estimation with all counties included, we are left with the difference due to each county’s exemption from the model (measured in percentage points). For clarity, we have included the possible outcomes from comparing two estimations and their interpretations in table 1. We are principally interested in those counties that demonstrated a smaller reduction in lights than the overall reduction in lights for that storm, which we term ‘better off’.

## 2.4. Specifying motivators of differential overall response to hurricanes via SHAP

Focusing on those hurricanes where we find an overall reduction in NTL due to the storm, we aim to identify those factors most important in explaining differences in hurricane recovery across counties, i.e. whether a county experienced more or less negative change in light after a hurricane (examining the whole time period). To this end, we train a tree-based machine learning classifier to predict the likelihood of each county having demonstrated a smaller reduction in light than the overall reduction in light for that storm (being ‘better off’), based on a variety of socioeconomic, demographic, and FEMA-aid based factors at four time stages: three months, six months, and one year after the storm, as well as for the full available timeline. We train three tree-based machine learning classification models first to predict the likelihood of a county being better off than the average county hit by the same hurricane: Random Forest, Extreme Gradient Boosting (XGBoost), and Light Gradient Boosting Machine. The model is trained on 70% of the data and tested on the remaining 30%. For each time period and treatment variable, we employ the best performing algorithm based on predictive accuracy. Supplementary table 1 gives a description of all variables, supplementary table 2 provides prediction accuracy for each algorithm, and the Data & Methods section of the supplementary material describes the classifiers in more detail. We back out the importance of each variable in predicting the outcome via the SHAP feature importance analysis methodology. Using the trained model, SHAP assigns a value of importance to each feature in the model, associated with a particular prediction value in the test set. For each of these predicted values, the contribution of each feature to that prediction is calculated, based on the magnitude of that feature’s contribution alone and in combination with other features. We average these local feature importance values to derive a globally generalizable importance score for each feature.

## 3. Results

### 3.1. Lack of full recovery three years after storms

Using the DiD specification, we compute a storm’s average effect on NTL for each hurricane. Separate models are constructed for the two treatment variables, and include NTL values for all time steps (months) available after the storm.

All storms for which there is three years or more of data after the storm demonstrate negative changes in light due to the storm (table 2). As a general trend, hurricanes that occurred earlier (and for which a longer timeline is therefore available) exhibit stronger reductions in NTL at lower  $p$ -values, the latter of which is likely due to the increase in data points in the post-hurricane period. For hurricanes with

statistically significant negative effects in areas receiving combined aid, we also see a decrease in NTL intensity associated with being in areas receiving individual-level aid alone.

These results suggest a lack of full recovery over two dimensions: over time, and geographically, in counties affected by the storm. We break down our analysis by time, re-estimating the DiD model while limiting the period after the storm month by month during the subsequent two years. Hence, we are able to compare the short-term changes in light month by month between storms, regardless of when the storm occurred.

For the oldest storms, Hermine, Matthew, Harvey, and Irma in the individual-aid case, the overall effect remains negative over time, for both types of treatment variables (figure 1). This is even more remarkable as the initial effect after a few months was an increase in light for Hermine and Matthew in the case of combined aid, and for Matthew also in the case of individual aid. For Michael, we find a strong decrease in radiance levels in the months after the storms hit with radiance levels then returning to pre-storm levels after about a year. For the most recent storms the overall effect is positive, though the time frame is likely too short to draw robust conclusions as also reflected by the respective  $p$ -values for Combined Aid (table 2). In both cases, the post-period of the hurricane is defined as all months following the storm. We find a similar lack of recovery when the post-period is defined as just the respective month, excluding the months between the storm and the month being evaluated, as shown in supplementary figure 16.


### 3.2. Hurricane response differs across affected counties

The previous two analyses suggest that at longer time frames, areas affected by hurricanes generally do not return to their previous levels of radiance. These results are in the geographic aggregate, however, and we may therefore miss any potential disparate geographic effects. Employing a ‘leave-one-out’ approach where the DiD model is re-estimated for all but one county, we are able to isolate the contribution of that county to the overall effect. That is, we can assess whether the exclusion of that county increases or decreases the overall change in NTL and, consequently, whether that country did comparatively better or worse than the average county affected by the hurricane (consult Methods for details).

Figure 2 portrays the direction and magnitude of the difference between the overall light change for a storm, and the light change when each county is omitted from the model. In the legend, we interpret the direction of these values as presented previously in table 1. In the cases of Hermine, Matthew, Harvey, and Dorian, which demonstrate overall negative changes in radiance due to the storm (portrayed on

**Table 2.** Average effect of individual hurricanes on nighttime light intensity in those areas eligible for combined storm-related FEMA aid, and individual-level only FEMA aid. Coefficients on the two variables of interest (the provision of combined, and individual-level only aid) are presented. These coefficients indicate the average percent change in nighttime light intensity in areas receiving the respective types of aid and are calculated separately for each hurricane. Their corresponding *p*-values are presented in the following columns. For these models, the entire available time series is considered. The number of months after the hurricane is presented in the ‘Months Post’ column.

Hurricane	Combined Aid	Individual Aid	<i>p</i> -Value, Combined	<i>p</i> -Value, Individ.	Months Post
Hermine	−2.926	−4.034	0.612	0.596	52
Matthew	−12.127	−14.747	0.000	0.000	51
Harvey	−1.846	−1.575	0.001	0.013	41
Irma	—	−11.868	—	0.000	40
Florence	0.901	3.616	0.501	0.025	28
Michael	3.428	7.018	0.232	0.018	27
Dorian	−9.829	—	0.001	—	16




the red/blue spectrum), we see a diversity of county-level responses to the storm when we omit each county from the empirical model. There is a similar diversity in response to the storm on the county level for those storms demonstrating an increase in NTL after the storm (purple/green spectrum). In both cases, within the treatment group, counties perform diversely, even in storms where the overall effect of the storm is of a large magnitude. There do not appear to be geographic patterns in the dispersion of positive or negative county-level influence. Results from the models including counties eligible for just individual-level aid are included in supplementary figure 17.

### 3.3. Time and FEMA aid designation affect factors facilitating recovery

We next investigate those socioeconomic and demographic factors that most strongly determine the likelihood of a county having experienced a smaller reduction in NTL compared to the average (being ‘better off’). Employing a tree-based machine learning approach, we back out the feature importance scores for the features used to predict each county’s likelihood of being better off.

In figure 3, we provide the comparative importance of each feature in determining if a county was better off, and the direction of their effect for each of four time periods and each of the




two treatment variables, combined public and individual aid and individual aid. As a robustness check, we also conduct the feature importance analyses using exclusively each of the tree-based algorithms tested and all time-cutoffs, regardless of performance. We find that while the specific point values differ, the general trends of how feature importances change as timelines increase, are preserved. These figures are included as supplementary figures 18–20.

In the case of combined aid shown in Panel (a) of figure 3, the total amount of FEMA aid is the most important feature across nearly all time horizons. After three months, the direction of its influence switches from positive to negative, where it remains, indicating that a larger total cost for a storm makes it more likely in the very short-term (up to three months) and less likely thereafter that a county is better off. It is intuitive there may be a negative correlation between aid and being better off, due to the fact that the provision of aid is likely linked to stronger

storm impact, though this may be ambiguous given that FEMA aid is provided with the explicit purpose to aid in the recovery of areas affected by disasters. In our data, we see that combined aid seems to fulfill this aim in the short term, as a larger amount of aid is associated with a higher likelihood of an area being better off. However, this effect reverses as the timeline continues, suggesting that aid may not be sufficient to motivate full recovery.

In the case of individual-level aid alone, the importance score of the cost of individual aid to FEMA is consistently negative across all time scales, indicating that a county with larger cost will be more likely to have a stronger reduction in NTL than the average for that storm. Additionally, employment, income, and age become increasingly important factors in the individual-level aid case, suggesting that they may support recovery and mitigate the effect of hurricanes. Results for those counties which experienced an increase in light after the storm are available in supplementary figure 21.



**Figure 3.** Comparative importance of different factors explaining why a county demonstrates a smaller reduction in lights than the aggregate reduction. SHAP importance measures the average impact each feature has on the model output (expressed as a percentage of all features’ explanatory power). Larger bars indicate that the feature is more important to the model. Negative values indicate the feature reduces the likelihood that the county is better off than average; positive values indicate the feature increases the likelihood of a smaller reduction in lights than the average. Panel (a) shows the feature importances for the model including those counties receiving combined aid, Panel (b) shows the feature importances for the model including those receiving individual-level FEMA aid alone. Comparative importances are provided for four periods (3, 6, and 12 months after the storm as well as the overall timeline, which can range from 52 to 16 months, dependent on the storm. Compare to figure 1).

At the six month time window, and especially in the case of individual-level aid alone, we see a strong trend in most factors predicting negatively. This can be explained by the fact that the overall proportion of counties considered to be better off are fewest for this time horizon (supplementary figure 22).

Viewing all timelines and treatments in aggregate, the three FEMA aid related variables (total cost, individual applications, and individual cost) consistently affect the likelihood of a county being better off.

Reconstructing the analysis omitting variables relating to the quantity of FEMA aid, we find that higher income has a consistent positive influence on the likelihood that a county is better off, which is more pronounced for those counties having received individual level FEMA aid (full results are included in supplementary figure 23.) This suggests that income has a moderating effect on the reduction of light due to hurricanes, especially in those areas most significantly affected by the disaster.

## 4. Discussion and conclusions

Our results shed light on the dynamics and motivators of disaster recovery. Using NTL data, we find evidence that those areas of the United States affected by Category 4 and 5 hurricanes between the years 2014 and 2020 generally do not show a full return to previous radiance levels, especially three years or longer after the storm. This finding supports existing literature suggesting that disaster-hit areas do not demonstrate full economic recovery, though this research addresses significantly longer timescales (20 years post-disaster) [9]. Given the time granularity of our data, we are able to investigate recovery dynamics month by month to provide a more detailed timescale than previously studied. We find that many areas affected by hurricanes demonstrate large fluctuations in radiance in the short term (both positive and negative) but those storms with the longest time frames do not return to the levels of NTL radiance they demonstrated previous to the storm.

VIIRS NTL data have been shown to be well correlated to economic activity and population density [24–27, 47]. Our results suggest that areas affected by hurricanes may experience a lasting reduction in economic activity, population density, or both. These findings support previous evidence on the recovery-dampening effects of out-migration from hurricane-affected areas [8]. It is difficult to distinguish between these motivators without further data on migration or business activity, or more granular data on the extent of damage in the areas affected by these storms. Without these data, it is possible that part of the reduction in radiance we see is due to ‘rebuilding better’ where areas affected by the storm do experience a full recovery, but simply emit less light due to new, more energy-efficient infrastructure.

Data on the extent of damage due to the storm exogenous to both economics and population density (not derived from the costs of reconstruction or insurance claims) would enable a more precise identification of those counties affected by the storm. Eligibility for FEMA aid is a strong but imperfect proxy for the effect of the storm. It is possible that counties neighboring those eligible for FEMA aid were damaged by the storm, but at a level that did not warrant FEMA aid [48]. While this may muddy the difference between our treatment and control groups, this leakage would lead to an underestimation of the effect of the storm on the treatment counties, indicating that the true effect of the storm was larger than presented here.

Of those factors contributing to a below-average light reduction after a storm, we find that for all periods after three months, the quantity of FEMA aid received by the county is most influential. Given NTL radiance’s strong correlation to economic activity, it may be that aid to public entities drives immediate recovery. Similarly using VIIRS NTL data, Chen and

Nordhaus [26] suggest that the types of economic activities best captured by NTL are those dependent on power for lighting at night, such as service, retail and transportation [4]. This could mean that these industries are fastest to recover with an influx of aid, and may motivate the rebound in radiance we see in the first few months after Hermine, Matthew, Michael and Dorian. The change in direction after three months suggests that FEMA aid may be integral to very short term recovery, as has been found in previous work [15, 16], but does not create sustainable recovery in the long term. This highlights the importance of analyzing longer time scales for a comprehensive overview of recovery dynamics.

When considering just those counties in which individual-level post-disaster aid was available, we find the factors most important in determining the likelihood of being worse-off than average are related to individual-level FEMA aid (total number of individual-level FEMA applications and the total cost of those applications). Similarly, in the case of those areas demonstrating an increase in NTL radiance after the storm, we find larger amounts of aid to be correlated with an even larger increase in light than the average (supplementary figure 21). This supports the hypothesis that while short term recovery is motivated by firms, longer term recovery is sustained by individuals.

When examining the full available timeline, economic and demographic factors such as employment and the proportion of elderly in the population usurp those related to individual-level aid, suggesting that socioeconomic status and age may play a role in the decision to rebuild or move from areas affected by a hurricane. Our finding that areas with larger elderly populations are more likely to demonstrate less NTL reduction than the average suggests that elderly populations may be more likely to return to hurricane-hit areas than younger residents. Though highly moderated by levels of severe housing damage, Fussell *et al* [49] also found this trend in studying the return to New Orleans after Hurricane Katrina, where two thirds of individuals over 40 returned after 14 months, compared to only half of those under 40 [49]. Similar to our results indicating that employment and income are positively related to the likelihood of a county recovering, they also find a reduction in the likelihood of migration with an increase in socioeconomic status. Using education as a proxy for socioeconomic status, they find that half of individuals with college degrees returned within four months of the storm, compared to 14 months for individuals without.

Some areas considered in this study were hit by several hurricanes during the timeframe of our analysis. As we do not have data on the number of hurricanes experienced by each area previous to the beginning of our dataset, it is difficult to determine the possible accumulated effect of

multiple hurricanes over time on the recovery or out-migration process without biasing our data on more recent storms. In the case of Hurricane Harvey which hit Texas, an area less frequently hit by hurricanes, we see a comparatively small, though negative, long-term change in NTL radiance. This suggests that the reduction in NTL radiance we see due to other storms in more frequently affected areas may indeed be motivated by accumulated disasters, perhaps by making individuals more likely to out-migrate or be reluctant to rebuild.

While we study a large geographic area, it is difficult to know how readily our results are generalizable to other regions affected by hurricanes, or other types of natural disasters. Especially given the importance of FEMA aid in determining county-level outcomes, it is difficult to generalize to regions with different disaster aid schemes (or none) or those areas with lower levels of development. Further research is required to understand whether, and to what extent, these results may apply to other regions.

Our results indicate the recovery of pre-storm levels of economic activity and population may be dependent on more long-term individual aid support. As extreme weather events become more frequent and intense, it is possible that this support must also be provided in new forms. Currently FEMA aid covers emergency disaster-related expenses and repair costs on the public and individual level. Our findings about the interaction of socioeconomic class and age in long-term recovery suggest that new forms of aid, such as those that encourage sustainable, disaster-informed rebuilding or that encourage individuals to take proactive measures in areas frequented by hurricanes, may be key to preserving future economic prosperity and manage hurricane-induced migration.

### Data availability statement

The dataset was constructed from publicly available data [36–40]. The combined dataset, containing nighttime light radiance values, socio-economic and demographic information, as well as disaster-related data from FEMA is available upon request.

### Code availability

The code used for the analysis and plots is available upon request.

### Acknowledgments

We gratefully acknowledge funding from the Volkswagen foundation (Europe and Global Challenges Initiative). We thank Maximilian Kotz, Sven Kunze, Robbin Middlanis and Annika Stechemesser for helpful discussion and feedback.

### Author contributions

K B H and L W designed the research. K B H constructed the dataset, analyzed the data, and constructed the models. K B H, and L W analyzed the results. K B H wrote the manuscript, and all authors provided feedback.

### Conflict of interest

The authors declare no competing interests.

### ORCID iDs

Kelsey Barton-Henry  <https://orcid.org/0000-0002-5274-2845>

Leonie Wenz  <https://orcid.org/0000-0002-8500-1568>

### References

- [1] Chen D *et al* 2021 Framing, context, and methods *Climate Change 2021: the Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* accepted
- [2] Fischer E M and Knutti R 2015 Anthropogenic contribution to global occurrence of heavy precipitation and high-temperature extremes *Nat. Clim. Change* **5** 560–4
- [3] Karmalkar A and Bradley R S 2017 Consequences of global warming of 1.5 °C and 2 °C for regional temperature and precipitation changes in the contiguous United States *PLoS One* **12** e0168697
- [4] King A D, Karoly D J and Henley B J 2017 Australian climate extremes at 1.5 °C and 2 °C of global warming *Nat. Clim. Change* **7** 412–6
- [5] Emergency Events Database (EM-DAT) 2021 Centre for research on the epidemiology of disasters (CRED) (Brussels: School of Public Health of the Université catholique de Louvain) (available at: [www.emdat.be](http://www.emdat.be))
- [6] Middelani R, Willner S N, Otto C, Kuhla K, Quante L and Levermann A 2021 Wave-like global economic ripple response to Hurricane Sandy *Environ. Res. Lett.* **16** 124049
- [7] Sebastian A, Gori A, Blessing R B, van der Wiel K and Bass B 2019 Disentangling the impacts of human and environmental change on catchment response during hurricane harvey *Environ. Res. Lett.* **14** 124023
- [8] Keenan J M and Hauer M E 2020 Resilience for whom? Demographic change and the redevelopment of the built environment in Puerto Rico *Environ. Res. Lett.* **15** 074028
- [9] Hsiang S and Jina A 2014 The causal effect of environmental catastrophe on long-run economic growth: evidence from 6,700 cyclones NBER Working Paper 20352 [10.3386/w20352](https://doi.org/10.3386/w20352)
- [10] Felbermayr G, Gröschl J, Sanders M, Schippers V and Steinwachs T 2022 The economic impact of weather anomalies *World Dev.* **151** 105745
- [11] Klomp J 2016 Economic development and natural disasters: a satellite data analysis *Glob. Environ. Change* **36** 67–88
- [12] Tveit T, Skoufias E and Strobl E 2022 Using VIIRS nightlights to estimate the impact of the 2015 Nepal earthquakes *Geoenviron. Disasters* **9** 00204–z
- [13] Skoufias E, Strobl E and Tveit T 2021 Can we rely on VIIRS nightlights to estimate the short-term impacts of natural disasters? Evidence from Five Southeast Asian Countries *Geomat. Nat. Haz. Risk* **12** 381–404
- [14] Gillespie T, Frankenberg E, Chum K F and Thomas D 2014 Nighttime lights time series of tsunami damage, recovery, and economic metrics in Sumatra, Indonesia *Remote Sens. Lett.* **5** 286–94

- [15] Paul B K 2005 Evidence against disaster-induced migration: the 2004 tornado in north central Bangladesh *Disasters* **29** 370–85
- [16] DeWaard J, Curtis K and Fussell E 2016 Population recovery in New Orleans after Hurricane Katrina: exploring the potential role of stage migration in migration systems *Popul. Environ.* **37** 449–63
- [17] Myers C, Slack T and Singlemann J 2008 Social vulnerability and migration in the wake of disaster: the case of Hurricanes Katrina and Rita *Popul. Environ.* **29** 271–91
- [18] Nawrotzki R J, Hunter L M, Runfola D M and Riosmena F 2015 Climate change as a migration driver from rural and urban Mexico *Environ. Res. Lett.* **10** 114023
- [19] Sheldon T L and Zhan C 2020 The impact of hurricanes and floods on domestic migration *J. Environ. Econ. Manage.* **115** 102726
- [20] Finch C, Emrich C T and Cutter S L 2010 Disaster disparities and differential recovery in New Orleans *Popul. Environ.* **31** 179–202
- [21] Skoufias E, Strobl E and Tveit T 2017 Natural disaster damage indices based on remotely sensed data: an application to Indonesia World Bank Policy Research Paper vol 8188 (Poverty and Equity Global Practice Group)
- [22] Hodler R and Raschky P 2014 Regional favoritism *Q. J. Econ.* **129** 995–1033
- [23] Zhao M, Cheng W, Zhou C, Li M, Wang N and Liu Q 2017 GDP spatialization and economic differences in south China based on NPP-VIIRS nighttime light imagery *Remote Sens.* **9** 673
- [24] Stathakis D, Tselios V and Faraslis I 2015 Urbanization in European regions based on night lights *Remote Sens. Appl.* **2** 26–34
- [25] Zhao N, Liu Y, Hsu F-C, Samson E L, Letu H, Liang D and Cao G 2020 Time series analysis of VIIRS-DNB nighttime lights imagery for change detection in urban areas: a case study of devastation in Puerto Rico from hurricanes Irma and Maria *Appl. Geogr.* **120** 102222
- [26] Chen X and Nordhaus W 2015 A test of the new VIIRS lights data set: population and economic output in Africa *Remote Sens.* **7** 4937–47
- [27] Ma T, Zhou C, Pei T, Haynie S and Fan J 2014 Responses of Suomi-NPP VIIRS derived nighttime lights to socioeconomic activity in China's cities *Remote Sens. Lett.* **5** 165–74
- [28] Schneider A, Friedl M A and Potere D 2008 A new map of global urban extent from MODIS satellite data *Environ. Res. Lett.* **4** 044003
- [29] Zhou Y, Smith S J, Zhao K, Imhoff M, Thomson A, Bond-Lamberty B, Asrar G R, Zhang X, He C and Elvidge C D 2015 A global map of urban extent from nightlights *Environ. Res. Lett.* **10** 054011
- [30] Small C, Pozzi F and Elvidge C 2005 Spatial analysis of global urban extent from DMSP-OLS night lights *Remote Sens. Environ.* **96** 277–91
- [31] Zegarra M A, Schmid J P, Palomino L and Seminario B 2020 Impact of Hurricane Dorian in the Bahamas: a view from the sky (Inter-American Development Bank)
- [32] Del Valle A, Elliott R J R, Strobl E and Tong M 2018 The short-term economic impact of tropical cyclones: satellite evidence from Guangdong Province *Econ. Disasters Clim. Change* **2** 225–35
- [33] Mohana P and Strobl E 2017 The short-term economic impact of tropical Cyclone Pam: an analysis using VIIRS nighttime satellite imagery *Int. J. Remote Sens.* **38** 1–15
- [34] Huang X, Wang C and Lu J 2019 Understanding the spatiotemporal development of human settlement in hurricane-prone areas on the US Atlantic and Gulf coasts using nighttime remote sensing *Nat. Hazards Earth Syst. Sci.* **19** 2141–55
- [35] Eker S, Garcia D, Valin H and van Ruijven B 2021 Using social media audience data to analyse the drivers of low-carbon diets *Environ. Res. Lett.* **16** 074001
- [36] Yin H, Xiao L, Youwen S, Ke Li M G, Zheng B and Lui C 2021 Unprecedented decline in summertime surface ozone over eastern China in 2020 comparably attributable to anthropogenic emission reductions and meteorology *Environ. Res. Lett.* **16** 124069
- [37] Lundberg S, Erion G, Chen H, DeGrave A, Prutkin J M, Nair B, Katz R, Himmelfarb J, Bansal N and Lee S-I 2020 From local explanations to global understanding with explainable AI for trees *Nat. Mach. Intell.* **2** 56–67
- [38] Lundberg S and Lee S 2017 A unified approach to interpreting model predictions *31st Conf. on Neural Information Processing Systems (NIPS 2017)* (arXiv:1705.07874)
- [39] Elvidge C D, Baugh K E, Zhizhin M and Hsu F-C Visible infrared imaging radiometer suite monthly cloud-free day/night band composite (Earth Observation Group, Payne Institute for Public Policy, Colorado School of Mines (available at: <https://eogdata.mines.edu/products/vnl/#monthly>))
- [40] Disasters and Assistance Federal Emergency Management Agency, United States' Department of Homeland Security (available at: [www.fema.gov/disaster/declarations](http://www.fema.gov/disaster/declarations)) (Accessed January 2022)
- [41] National Oceanic and Atmospheric Administration 2021 National Centers for Environmental Information *Historical hurricane tracks* (available at: <https://coast.noaa.gov/digitalcoast/data/hurricanes.html>) (Accessed 12 July 2022)
- [42] Urban-Rural Classification Scheme for Counties 2017 National center for health statistics (available at: [www.cdc.gov/nchs/data\\_access/urban\\_rural.htm](http://www.cdc.gov/nchs/data_access/urban_rural.htm)) (Accessed January 2022)
- [43] County-Level Shapefiles 2011 Center for disease control and prevention (available at: [www.cdc.gov/epiinfo/support/downloads/shapefiles.html#USA\\_Shapefiles](http://www.cdc.gov/epiinfo/support/downloads/shapefiles.html#USA_Shapefiles)) (Accessed January 2022)
- [44] Employment by County, Metro, and Other Areas 2021 Bureau of economic analysis (available at: [www.bea.gov/data/employment/employment-county-metro-and-other-areas](http://www.bea.gov/data/employment/employment-county-metro-and-other-areas)) (Accessed January 2022)
- [45] County Population by Characteristics: 2010–2019 2021 United States Census Bureau (available at: [www.census.gov/data/tables/timeseries/demo/popest/2010s-counties-detail.html](http://www.census.gov/data/tables/timeseries/demo/popest/2010s-counties-detail.html)) (Accessed January 2022)
- [46] Angrist J D and Pischke J S 2009 *Mostly Harmless Econometrics: An Empiricist's Companion* vol 1 (New Jersey: Princeton University Press) pp 165–87
- [47] Chen X and Nordhaus W 2019 VIIRS nighttime lights in the estimation of cross-sectional and time-series GDP *Remote Sens.* **11** 1057
- [48] Eichenauer V Z, Fuchs A, Kunze S and Strobl E 2020 Distortions in aid allocation of United Nations flash appeals: evidence from the 2015 Nepal Earthquake *World Dev.* **136** 105023
- [49] Fussell E, Sastry N and VanLandingham M 2009 Race, socioeconomic status, and return migration to New Orleans after Hurricane Katrina *Popul. Environ.* **31** 150–75